

CLOTRISURE (Clotrimazole 1% Cream)

Active Ingredient: Clotrimazole BP 1%w/w

Excipient: Cetomacrogol 1000 BP, Cetostearyl alcohol BP, White soft paraffin BP, Light Liquid Paraffin BP, Benzyl alcohol BP, Glycerol BP, Polysorbate 20 BP, Purified Water BP

Dosage Form: Cream (External Preparation)

Product Description: White smooth cream, odourless.

Pharmacodynamics:

Pharmacotherapeutic group: Antifungals for topical use – imidazole and triazole derivatives, ATC code: D01A C01

Mechanism of action

Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the cytoplasmic membrane.

Pharmacodynamic Effects

Clotrimazole has a broad antimycotic spectrum of action in vitro and in vivo, which includes dermatophytes, yeasts, moulds, etc. Under appropriate test conditions, the MIC values for these types of fungi are in the region of less than 0.062-8.0 µg/ml substrate.

The mode of action of clotrimazole is primarily fungistatic or fungicidal depending on the concentration of clotrimazole at the site of infection. In vitro activity is limited to proliferating fungal elements; fungal spores are only slightly sensitive.

In addition to its antimycotic action, clotrimazole also acts on gram-positive microorganisms (Streptococci/ Staphylococci/ Gardnerella vaginalis), and gram-negative microorganisms (Bacteroides). In vitro clotrimazole inhibits the multiplication of Corynebacteria and gram-positive cocci - with the exception of Enterococci - in concentrations of 0.5-10 µg/ml substrate.

Primarily resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

Pharmacokinetics:

Pharmacokinetic investigations after dermal application have shown that clotrimazole is minimally absorbed from the intact or inflamed skin into the human blood circulation. The resulting peak serum concentrations of clotrimazole were below the detection limit of 0.001 mcg/ml, suggesting that clotrimazole applied topically is unlikely to lead to measurable systemic effects or side effects.

Indication: For the treatment of:

- i. All dermatomycoses due to moulds and other fungi (e.g. Trichophyton species)
- ii. All dermatomycoses due to yeasts (Candida species). These include ringworm (tinea) infections (e.g. athlete's foot), paronychia, pityriasis versicolor, erythrasma and intertrigo.
- iii. Skin diseases showing secondary infection with these fungi.
- iv. Candidal nappy rash, vulvitis and balanitis.

Recommended Dose: The cream should be applied thinly and evenly to the affected area 2 – 3 times daily and rubbed in gently. A strip of cream (½ cm long) is enough to treat an area of about the size of the hand.

If the feet are infected, they should be thoroughly washed and dried, especially between the toes, before applying the cream.

Treatment should be continued for at least one month for dermatophyte infections, or for at least two weeks for candidal infections.

Route of Administration: For topical route of administration.

Contraindication: Hypersensitivity to the active substance or to any of the excipients. Do not use the cream to treat nail or scalp infections.

Special warnings and precautions for use: This product contains cetostearyl alcohol, which

may cause local skin reactions (e.g. contact dermatitis). The cream also contains benzyl alcohol which may cause allergic reactions and mild local irritation.

Interactions with other medicaments: Laboratory tests have suggested that, when used together, this product may cause damage to latex contraceptives. Consequently the effectiveness of such contraceptives may be reduced. Patients should be advised to use alternative precautions for at least five days after using this product.

Fertility, pregnancy and lactation:

Fertility: No human studies of the effects of clotrimazole on fertility have been performed; however, animal studies have not demonstrated any effects of the drug on fertility.

Pregnancy:

There is a limited amount of data from the use of clotrimazole in pregnant women. Animal studies with clotrimazole have shown reproductive toxicity at high oral doses. At the low systemic exposures of clotrimazole following topical treatment, harmful effects with respect to reproductive toxicity are not predicted. Clotrimazole can be used during pregnancy but only under the supervision of a physician or midwife.

Lactation:

There are no data on the excretion of clotrimazole into human milk. However, systemic absorption is minimal after administration and is unlikely to lead to systemic effects. Clotrimazole may be used during lactation.

Side Effect:

As the listed undesirable effects are based on spontaneous reports, assigning an accurate frequency of occurrence for each is not possible.

Immune system disorders: anaphylactic reaction, angioedema, hypersensitivity.

Vascular disorders: syncope, hypotension.

Respiratory, thoracic and mediastinal disorders: dyspnoea.

Skin and subcutaneous tissue disorders: blisters, dermatitis contact, erythema, paresthesia, skin exfoliation, pruritus, rash, urticaria, stinging skin/burning sensation skin.

General disorders and administration site conditions: application site irritation, application site reaction, oedema, pain.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Symptoms and Treatment of Overdose:

No risk of acute intoxication is seen as it is unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favourable to absorption) or inadvertent oral ingestion. There is no specific antidote.

However, in the event of accidental oral ingestion, routine measures such as gastric lavage should be performed only if clinical symptoms of overdose become apparent (e.g. dizziness, nausea or vomiting). Gastric lavage should be carried out only if the airway can be protected adequately.

Effect on ability to drive and use machine: Clotrimazole cream has no or negligible influence on the ability to drive or use machines.

Instruction for Use: If Clotrimazole 1% Cream has been prescribed for you by your doctor, follow any instructions he/she may have given you. If you purchased this product without a prescription, follow these directions closely:

- Before use, pierce the tube seal by inverting the cap over the end of the tube and press.
- If the feet are infected, they should be washed and dried thoroughly, especially between the toes, before applying the cream.
- Clotrimazole 1% Cream should be applied thinly and evenly to the affected areas two or three times daily and rubbed in gently.

- A strip of cream (1/2 cm long) is enough to treat an area of about the size of the hand.
- The duration of the treatment depends upon the type of infection. Generally, a minimum of two weeks is required, although up to four weeks may be necessary.
- If you have athlete's foot, it may help to use an antifungal dusting powder as well. Ask your doctor or pharmacist to recommend one.

Storage Condition: Store below 30°C.

Shelf Life: 36 months from date of manufacture.

Therapeutic Code: ATC code: D01A C01

Presentation: 20g Collapsible Aluminum Tube having white colour cap along with Package insert packed in multicolor carton.

Date of revision: 29th July, 2022

Manufacturer:

ICPA Health Products Limited
286/287/288, GIDC, Ankleshwar 393002, India.
E. mail: wecare@icpahealth.com
Call us at 02646 221915

Product Registration Holder:

Healol Pharmaceuticals Sdn Bhd,
74-3, Jalan Wangsa Delima 6,
KLSC Wangsa Maju, 53300,
Kuala Lumpur, Malaysia.

™ Trade mark of ICPA