

Malasone

Ear drops, suspension

Treatment of canine otitis external

COMPOSITION

Each mL contains:

Marbofloxacin	3 mg
Clotrimazole	10 mg
Dexamethasone acetate.....	1 mg

DESCRIPTION

The finished product is ivory white to yellow oily suspension

PHARMACOLOGY

Pharmacodynamic properties

The preparation combines three active ingredients:

-Marbofloxacin, a synthetic bactericidal agent belonging to the fluoroquinolone family that acts by inhibiting DNA gyrase. It exhibits a broad spectrum of activity against Gram-positive bacteria (e.g. *Staphylococcus intermedius*) and against Gram-negative organisms (*Pseudomonas aeruginosa*, *Escherichia coli* and *Proteus mirabilis*).

-Clotrimazole, an anti-fungal agent that belongs to the imidazole family and which acts by causing changes in membrane permeability, allowing intracellular compounds to leak from the cell and thus inhibiting cellular molecular synthesis. It exhibits a wide spectrum of activity and is aimed, in particular, at *Malassezia pachydermatis*;

-Dexamethasone acetate, a synthetic glucocorticoid exhibiting anti-inflammatory and anti-pruritic activity.

Pharmacokinetic particulars

Pharmacokinetics studies in dogs at the therapeutic dosage have shown that:

Marbofloxacin plasma concentrations peak at 0.06 µg/ml on the 14th day of treatment.

Marbofloxacin binds weakly to plasma proteins (< 10% in dogs) and is eliminated slowly, mainly in the active form, over 2/3 in urine and over 1/3 in faeces.

Clotrimazole absorption is extremely poor (plasma concentration < 0.04 mcg/ml).

Dexamethasone acetate plasma concentration reaches 1.25 ng/ml on the 14th day of treatment. Dexamethasone resorption is not increased by the inflammatory process induced by otitis.

INDICATIONS

Treatment of otitis externa of both bacterial and fungal origin respectively due to bacteria sensitive to marbofloxacin, and fungi especially *Malassezia pachydermatis* sensitive to clotrimazole.

The product should be used based on susceptibility testing.

TARGET ANIMALS

Dogs

ROUTE OF ADMINISTRATION

Ear drops, suspension

DOSAGE AND ADMINISTRATION

Shake well before use.

Apply ten drops into the ear once daily for 7 to 14 days. After 7 days of treatment, the veterinary surgeon should evaluate the necessity to extend the treatment another week.

One drop of the preparation contains 71mcg marbofloxacin, 237mcg clotrimazole and 23.7mcg dexamethasone acetate.

After application, the base of the ear may be massaged briefly and gently to allow the preparation to penetrate to the lower part of the ear canal.

When the product is indicated for use in several dogs, use one cannula per dog.

CONTRAINDICATIONS

Do not administer to dogs suffering from perforation of the tympanic membrane. Do not administer to dogs with known hypersensitivity to any of the ingredients.

Do not administer to pregnant or lactating bitches.

OVERDOSE (symptoms, emergency procedures, antidotes), if necessary

Changes in biochemical and haematological parameters (such as increase of alkaline phosphatase, aminotransferase, some limited neutrophilia, eosinopenia, lymphopenia) are observed with three fold the recommended dosage; such changes are not serious and will reverse once the treatment has stopped.

ADVERSE REACTIONS (FREQUENCY AND SERIOUSNESS)

Usual adverse reactions associated with corticosteroid drugs may be observed (changes in biochemical and hematological parameters, such as increase of alkaline phosphatase, and of aminotransferase, some limited neutrophilia).

Prolonged and intensive use of topical corticosteroid preparations is known to trigger local and systemic effects, including suppression of adrenal function, thinning of the epidermis and delayed wound healing.

On rare occasions, the use of this product may be associated with deafness, mainly in elderly dogs and mostly of a transient nature.

SPECIAL WARNINGS FOR EACH TARGET SPECIES

None

SPECIAL PRECAUTIONS FOR USE IN ANIMALS

Heavy reliance on a single class of antibiotic may result in the induction of resistance in a bacterial population.

It is prudent to reserve the fluoroquinolones for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antibiotics.

Before treating with the product, the integrity of the tympanic membrane must be verified.

The external ear canal should be meticulously cleaned and dried before treatment.

SPECIAL PRECAUTIONS TO BE TAKEN BY THE PERSON ADMINISTERING THE MEDICINAL PRODUCTS TO ANIMALS

Wash hands carefully after applying the product.

Avoid contact with eyes. If splashed in the eye, rinse with copious amounts of water.

Persons with known hypersensitivity to compounds in the product should avoid any contact with the product.

- Keep out of reach of children

- Jauhkan daripada kanak-kanak

USE DURING PREGNANCY, LACTATION OR LAY

See "Contraindications".

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

None known

WITHDRAWAL TIME: Not applicable.

STORAGE

Store in a cool dry place not exceeding 30°C.

SHELF LIFE: 2 years

SHELF LIFE AFTER OPENING: 2 months

PACKAGING: 10 ml dropper bottle.

REVISED DATE: Aug 10, 2022