

pharmaniaga®

IDAMAN METFORMIN TABLET
500MG

DESCRIPTION

A white to off-white, round, biconvex and film-coated tablet with breakline on one side

COMPOSITION

Each tablet contains Metformin Hydrochloride 500mg.

PHARMACODYNAMICS

Pharmacotherapeutic group: Blood glucose lowering drugs. Biguanides; ATC code: A10BA02

Mechanism of action

Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and postprandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

Metformin may act via 3 mechanisms:

- (1) reduction of hepatic glucose production by inhibiting gluconeogenesis and glycogenolysis.
- (2) in muscle, by increasing insulin sensitivity, improving peripheral glucose uptake and utilization.
- (3) and delay of intestinal glucose absorption.

Metformin stimulates intracellular glycogen synthesis by acting on glycogen synthase. Metformin increases the transport capacity of all types of membrane glucose transporters (GLUTs) known to date.

Pharmacodynamic effects

Based on published clinical data, use of metformin was associated with either a stable body weight or modest weight loss.

In humans, independently of its action on glycaemia, metformin has favourable effects on lipid metabolism. Metformin reduces total cholesterol, LDL cholesterol and triglyceride levels at therapeutic doses.

Clinical efficacy:

The published prospective randomised study (UKPDS) has established the long-term benefit of intensive blood glucose control in adult patients with type 2 diabetes.

Analysis of the results for overweight patients treated with metformin after failure of diet alone showed a significant reduction of the absolute risk for:

- diabetes-related complication in the metformin group (29.8 events/1000 patient-years) versus diet alone (43.3 events/1000 patient-years), and versus the combined sulfonylurea and insulin monotherapy groups (40.1 events/1000 patient-years);
- diabetes-related mortality: metformin (7.5 events/1000 patient-years), diet alone (12.7 events/1000 patient-years);
- overall mortality: metformin (13.5 events/1000 patient-years) versus diet alone (20.6 events/1000 patient-years), and versus the combined sulfonylurea and insulin monotherapy groups (18.9 events/1000 patient-years);
- myocardial infarction: metformin (11 events/1000 patient-years), diet alone (18 events/1000 patient-years).

Benefit regarding the outcome has not been shown for metformin used as second-line therapy, in combination with a sulfonylurea. In type 1 diabetes, the combination of metformin and insulin has been used in selected patients, but the clinical benefit of this combination has not been formally established.

Paediatric population

Controlled clinical studies in a limited paediatric population aged 10-16 years treated during 1 year demonstrated a similar response in glycaemic control to that seen in adults.

PHARMACOKINETICS

Absorption:

After an oral dose of metformin hydrochloride tablet, maximum plasma concentration (C_{max}) is reached in approximately 2.5 hours (t_{max}). Absolute bioavailability of a 500 mg metformin hydrochloride tablet is approximately 50-60% in healthy patients. After an oral dose, the non-absorbed fraction recovered in faeces was 20-30%. After oral administration, metformin absorption is saturable and incomplete. It is assumed that the pharmacokinetics of metformin absorption is non-linear. At the recommended metformin doses and dosing schedules, steady state plasma concentrations are reached within 24 to 48 hours and are generally less than 1 microgram/ml. Maximum metformin plasma levels (C_{max}) did not exceed 5 microgram/ml, even at maximum doses. Food decreases the extent and slightly delays the absorption of metformin.

Following oral administration of a 850 mg tablet, a 40% lower plasma peak concentration, a 25% decrease in AUC (area under the curve) and a 35 minute prolongation of the time to peak plasma concentration were observed. The clinical relevance of these findings is unknown.

Distribution:

Protein binding in plasma is negligible.

Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution.

The mean volume of distribution (V_d) ranged between 63-276 L.

Metabolism:

Metformin is excreted unchanged in the urine.

No metabolites have been identified in humans.

Elimination:

Renal clearance of metformin is >400 ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

Characteristics in specific groups of patients

Renal impairment

The available data in subjects with moderate renal insufficiency are scarce and no reliable estimation of the systemic exposure to metformin in this subgroup as

compared to subjects with normal renal function could be made. Therefore, the dose adaptation should be made upon clinical efficacy/tolerability considerations.

Paediatrics population

Single dose study: As per reported data, after single doses of metformin hydrochloride 500 mg, paediatric patients have shown similar pharmacokinetic profile to that observed in healthy adults.

Multiple dose study: Available data are restricted to one study. After repeated doses of 500 mg twice daily for 7 days in paediatric patients, the peak plasma concentration (C_{max}) and systemic exposure (AUC_{0-t}) were reduced by approximately 33% and 40%, respectively compared to diabetic adults who received repeated doses of 500 mg twice daily for 14 days. As the dose is individually titrated based on glycaemic control, this is of limited clinical relevance.

Preclinical safety data

Published preclinical data reveal no special hazard for humans based on conventional studies on safety, pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and reproductive toxicity

INDICATION

Treatment of type 2 diabetes mellitus, when dietary management and exercise alone does not result in adequate glycaemic control.

- In adults, IDAMAN METFORMIN TABLET 500MG may be used as monotherapy or in combination with other oral anti-diabetic agents or with insulin.
- In children from 10 years of age and adolescents, IDAMAN METFORMIN TABLET 500MG may be used as monotherapy or in combination with insulin.

A reduction of diabetic complications has been shown in overweight type 2 diabetic adult patients treated with metformin as first-line therapy after diet failure.

RECOMMENDED DOSAGE

Adults with normal renal function (GFR ≥ 90 mL/min):

Monotherapy and combination with other oral antidiabetic agents:

- The usual starting dose is one tablet 2 or 3 times daily given during or after meals. After 10 to 15 days the dose should be adjusted on the basis of blood glucose measurements. A slow increase of dose may improve gastrointestinal tolerability.
- The maximum recommended dose of metformin hydrochloride is 3 g daily, taken as 3 divided doses.
- If transfer from another oral anti-diabetic agent is intended: discontinue the other agent and initiate metformin at the dose indicated above.

Combination with insulin:

Metformin and insulin may be used in combination therapy to achieve better blood glucose control. Metformin hydrochloride is given at the usual starting dose of one tablet 2 or 3 times daily, while insulin dosage is adjusted on the basis of blood glucose measurements.

Elderly

Due to the potential for decreased renal function in elderly subjects, the metformin dosage should be adjusted based on renal function. Regular assessment of renal function is necessary.

Patients with renal impairment

A GFR should be assessed before initiation of treatment with metformin containing products and at least annually thereafter. In patients at an increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3-6 months.

GFR mL/min	Total maximum daily dose (to be divided into 2-3 daily doses)	Additional considerations
60-89	3000 mg	Dose reduction may be considered in relation to declining renal function.
45-59	2000 mg	Factors that may increase the risk of lactic acidosis should be reviewed before considering initiation of metformin. The starting dose is at most half of the maximum dose.
30-44	1000 mg	
< 30	-	Metformin is contraindicated.

Paediatric population:

Monotherapy and combination with insulin

- IDAMAN METFORMIN TABLET 500MG can be used in children from 10 years of age and adolescents.
- The usual starting dose is 500 mg or 850 mg metformin hydrochloride once daily, given during or after meals.
- After 10 to 15 days the dose should be adjusted on the basis of blood glucose measurements. A slow increase of dose may improve gastrointestinal tolerability. The maximum recommended dose of metformin hydrochloride is 2g daily, taken as 2 or 3 divided doses.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

- Hypersensitivity to metformin or to any of the excipients.
- Any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis).
- Diabetic pre-coma.
- Severely reduced kidney function (GFR <30 mL/min).
- Acute conditions with the potential to alter renal function such as: dehydration, severe infection, shock.
- Disease which may cause tissue hypoxia (especially acute disease, or worsening chronic disease) such as : decompensated heart failure, respiratory failure, recent myocardial infarction, shock.
- Hepatic insufficiency, acute alcohol intoxication, alcoholism.

WARNINGS & PRECAUTIONS

Lactic acidosis Lactic acidosis, a very rare but serious metabolic complication, most often occurs at acute worsening of renal function or cardiorespiratory illness or sepsis. Metformin accumulation occurs at acute worsening of renal function and increases the risk of lactic acidosis. In case of dehydration (severe diarrhoea or vomiting, fever or reduced fluid intake), metformin should be temporarily discontinued and contact with a health care professional is recommended. Medicinal products that can acutely impair renal function (such as antihypertensives, diuretics and NSAIDs) should be initiated with caution in metformin-treated patients. Other risk factors for lactic acidosis are excessive alcohol intake, hepatic insufficiency, inadequately controlled diabetes, ketosis, prolonged fasting and any conditions associated with hypoxia, as well as concomitant use of medicinal products that may cause lactic acidosis. Patients and/or care-givers should be informed of the risk of lactic acidosis. Lactic

acidosis is characterised by acidotic dyspnoea, abdominal pain, muscle cramps, asthenia and hypothermia followed by coma. In case of suspected symptoms, the patient should stop taking metformin and seek immediate medical attention. Diagnostic laboratory findings are decreased blood pH (<7.35), increased plasma lactate levels (>5 mmol/L) and an increased anion gap and lactate/pyruvate ratio.

Renal function:

GFR should be assessed before treatment initiation and regularly thereafter. Metformin is contraindicated in patients with GFR <30 mL/min and should be temporarily discontinued in the presence of conditions that alter renal function.

Surgery:

Metformin must be discontinued at the time of surgery under general, spinal or epidural anesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and provided that renal function has been re-evaluated and found to be stable.

Administration of Iodinated Contrast Agents

Intravascular administration of iodinated contrast agents may lead to contrast induced nephropathy, resulting in metformin accumulation and an increased risk of lactic acidosis. Metformin should be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable.

Cardiac Function

Patients with heart failure are more at risk of hypoxia and renal insufficiency. In patients with stable chronic heart failure, metformin may be used with a regular monitoring of cardiac and renal function.

For patients with acute and unstable heart failure, metformin is contraindicated.

Paediatric population:

The diagnosis of type 2 diabetes mellitus should be confirmed before treatment with metformin is initiated.

No effect of metformin on growth and puberty has been detected during controlled clinical studies of one-year duration but no long-term data on these specific points are available. Therefore, a careful follow-up of the effect of metformin on these parameters in metformin-treated children, especially prepubescent children, is recommended.

Children aged between 10 and 12 years:

Based on available controlled clinical studies conducted in children and adolescents whereby 15 subject aged 10 and 12 years were included, particular caution is recommended when prescribing to children aged between 10 and 12 years although efficacy and safety of metformin in these children did not differ from efficacy and safety in older children and adolescents.

Other precautions:

All patients should continue their diet with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet. The usual laboratory tests for diabetes monitoring should be performed regularly. Metformin alone does not cause hypoglycemia, but caution is advised when it is used concurrently with insulin or other oral antidiabetics (e.g. sulfonylureas or meglitinides).

Metformin may reduce vitamin B12 serum levels. The risk of low vitamin B12 levels increases with increasing metformin dose, treatment duration, and/or in patients with risk factors known to cause vitamin B12 deficiency. In case of suspicion of vitamin B12 deficiency (such as anemia or neutropathy), vitamin B12 serum levels should be monitored. Periodic vitamin B12 monitoring could be necessary in patients with risk factors for vitamin B12 deficiency. Metformin therapy should be continued for as long as it is tolerated and not contraindicated and appropriate corrective treatment for vitamin B12 deficiency provided in line with current clinical guidelines.

INTERACTIONS WITH OTHER MEDICAMENTS

Concomitant use not recommended:

Alcohol:

Alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in case of: fasting, malnutrition or hepatic impairment.

Iodinated contrast agents:

Metformin must be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable.

Combinations requiring precautions for use:

Some medicinal products can adversely affect renal function which may increase the risk of lactic acidosis, e.g. NSAIDs, including selective cyclo-oxygenase (COX) II inhibitors, ACE inhibitors, angiotensin II receptor antagonists and diuretics, especially loop diuretics. When starting or using such products in combination with metformin, close monitoring of renal function is necessary.

Medicinal products with intrinsic hyperglycemic activity (e.g. glucocorticoids (systemic and local routes) and sympathomimetics).

More frequent blood glucose monitoring may be required, especially at the beginning of treatment. If necessary, adjust the metformin dosage during therapy with the respective medicinal product and upon its discontinuation.

Organic cation transporters (OCT):

Metformin is a substrate of both transporters OCT1 and OCT2. Co-administration of metformin with:

- Inhibitors of OCT1 (such as verapamil) may reduce efficacy of metformin.
- Inducers of OCT1 (such as rifampicin) may increase gastrointestinal absorption and efficacy of metformin.
- Inhibitors of OCT2 (such as cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazole) may decrease the renal elimination of metformin and thus lead to an increase in metformin plasma concentration.
- Inhibitors of both OCT1 and OCT2 (such as crizotinib, olaparib) may alter efficacy and renal elimination of metformin.

Caution is therefore advised, especially in patients with renal impairment, when these drugs are co-administered with metformin, as metformin plasma concentration may increase. If needed, dose adjustment of metformin may be considered as OCT inhibitors/inducers may alter the efficacy of metformin

PREGNANCY AND LACTATION

Pregnancy:

Uncontrolled diabetes during pregnancy (gestational or permanent) is associated with increased risk congenital abnormalities and perinatal mortality. A limited amount of data from the use of metformin in pregnant women does not indicate an increased risk of congenital abnormalities.

Based on available animal studies, they do not indicate harmful effects with respect to pregnancy, embryonic or fetal development, parturition or postnatal development.

When the patient plans to become pregnant and during pregnancy, it is recommended that diabetes is not treated with metformin but insulin be used to maintain blood glucose levels as close to normal as possible, to reduce the risk of malformations of the foetus.

Breast-feeding:

Metformin is excreted into human breast milk. No adverse effects were observed in breastfed Newborns/ infants. However, as only limited data are available, breast-feeding is not recommended during metformin treatment. A decision on whether to discontinue breast-feeding should be made, taking into account the benefit of breastfeeding and the potential risk to adverse effects on the child.

Fertility

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

SIDE EFFECTS

During treatment initiation, the most common adverse reactions are nausea, vomiting, diarrhea, abdominal pain and loss of appetite which resolve spontaneously in most cases. To prevent them, it is recommended that metformin be taken in 2 or 3 daily doses and to increase slowly the doses.

Gastrointestinal disorders:

Very common: Gastrointestinal disorders such as nausea, vomiting, diarrhea, abdominal pain and loss of appetite. These undesirable effects occur most frequently during initiation of therapy and resolve spontaneously in most cases. To prevent them, it is recommended that metformin be taken in 2 or 3 daily doses during or after meals. A slow increase of the dose may also improve gastrointestinal tolerability.

Metabolism and nutrition disorders:

Common: Vitamin B12 decrease/deficiency
Very rare: Lactic acidosis

Nervous system disorders:

Common: Taste disturbance.

Skin and subcutaneous tissue disorders:

Very rare: Skin reactions such as erythema, pruritus, urticaria.

Hepatobiliary disorders:

Very rare: Isolated reports of liver functions tests abnormalities or hepatitis resolving upon metformin discontinuation.

Paediatric Population

Adverse event reporting was similar in nature and severity to that reported in adults.

SYMPTOMS AND TREATMENT OF OVERDOSE

Hypoglycaemia has not been seen with metformin hydrochloride doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose of metformin or concomitant risks may lead to lactic acidosis. Lactic acidosis is a medical emergency and must be treated in hospital. The most effective method to remove lactate and metformin is hemodialysis.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Metformin monotherapy does not cause hypoglycemia and therefore has no effect on the ability to drive or to use machines. However, patients should be alerted to the risk of hypoglycemia when metformin is used in combination with other antidiabetic agents (e.g. sulfonylureas, insulin or meglitinides).

INSTRUCTION OF USE

Metformin can be taken during or after meals.

STORAGE

Store below 30°C.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PACKING

Blister Pack of 10's
a) 50 strips of 10 tablets
b) 10 strips of 10 tablets

REGISTRATION NO: MAL23026018AZ

Lt-319.01

For further information, please consult your doctor or your pharmacist.

**KEEP MEDICINES OUT OF REACH OF CHILDREN
JAUHI UBAT-UBATAN DARI KANAK-KANAK**

Revision No: 01

Revision Date : 20 Jun 2024

**PRODUCT REGISTRATION HOLDER & MANUFACTURER
IDAMAN PHARMA MANUFACTURING SDN BHD (661901P)
LOT 24 & 25, JALAN PERUSAHAAN LAPAN,
BAKAR ARANG INDUSTRIAL ESTATE, 08000 SUNGAI PETANI,
KEDAH DARUL AMAN, MALAYSIA.**