

BETA FUSIDIC CREAM

MAL23056038AZ

DESCRIPTION:

CREAM

Colour : White

Odour : Natural

CONTENT:

Fusidic Acid 2.0% w/w

Betamethasone 17-Valerate equivalent to Betamethasone 0.1% w/w

Chlorocresol 0.1% w/w

PHARMACODYNAMIC:

Beta Fusidic Cream combines the well-known anti-inflammatory and antipruritic effects of betamethasone with the potent topical antibacterial action of Fusidic Acid.

Betamethasone Valerate is a topical steroid rapidly effective in those inflammatory dermatoses which normally respond to this form of therapy.

When applied topically, Fusidic Acid is effective against *Staphylococcus aureus*, *Streptococci*, *Corynebacteria*, *Neisseria* and certain *Clostridia* and *Bacteroides*. The antibacterial activity of Fusidic Acid is not diminished in the presence of betamethasone.

PHARMACOKINETICS:

There are no data which define the pharmacokinetics of Fusidic Acid, following topical administration in man.

However, in vitro studies show that Fusidic Acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to Fusidic Acid and the condition of the skin. Fusidic Acid is excreted mainly in the bile with little excreted in the urine.

Betamethasone is absorbed following topical administration. The degree of absorption is dependent on various factors including skin condition and site of application. Betamethasone is metabolised largely in the liver but also to a limited extent in the kidneys, and the inactive metabolites are excreted with the urine.

INDICATIONS:

Beta Fusidic Cream is indicated in inflammatory dermatoses where bacterial infection is present or likely to occur. Inflammatory dermatoses include atopic eczema, discoid eczema, stasis eczema, seborrhoeic dermatitis, contact dermatitis, lichen simplex chronicus, psoriasis, discoid lupus erythematosus.

RECOMMENDED DOSE:

Uncovered lesion : Apply 2-3 daily

Covered lesion : Less frequent application may be adequate

ROUTE OF ADMINISTRATION:

Topical

CONTRAINDICATIONS:

Hypersensitivity to Fusidic Acid, Betamethasone Valerate or to any of the excipients.

Due to the content of corticosteroid, Beta Fusidic Cream is contraindicated in the following conditions:

Systemic fungal infections.

Primary skin infections caused by fungi, virus or bacteria, either untreated or uncontrolled by appropriate treatment (refer section on Warnings and Precautions below).

Skin manifestations in relation to tuberculosis or syphilis, either untreated or uncontrolled by appropriate therapy.

Perioral dermatitis and rosacea.

WARNINGS AND PRECAUTIONS:

Long-term continuous topical therapy with Beta Fusidic Cream should be avoided.

Depending on the application site, possible systemic absorption of Betamethasone Valerate should always be considered during treatment with Beta Fusidic Cream.

Due to the content of corticosteroid, Beta Fusidic Cream should be used with care near the eyes. Avoid getting Beta Fusidic Cream into the eyes.

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Reversible hypothalamic-pituitary-adrenal (HPA) axis suppression may occur following systemic absorption of topical corticosteroids. Beta Fusidic Cream should be used with care in children as paediatric patients may demonstrate greater susceptibility to topical corticosteroids-induced HPA axis suppression and Cushing's syndrome than adult patients. Avoid large amounts, occlusion and prolonged treatment.

Due to the content of Betamethasone Valerate, prolonged topical use of Beta Fusidic Cream may cause skin atrophy.

Bacterial resistance has been reported to occur with the topical use of Fusidic Acid. As with all antibiotics, extended or recurrent use of Fusidic Acid may increase the risk of developing antibiotic resistance. Limiting therapy with topical Fusidic Acid and Betamethasone Valerate to no more than 14 days at a time will minimise the risk of developing resistance.

This also prevents the risk that the immunosuppressive action of corticosteroid might mask any potential symptoms of infections due to antibiotic-resistant bacteria.

Due to the content of corticosteroid having immunosuppressant effect, Beta Fusidic Cream may be associated with increased susceptibility to infection, aggravation of existing infection, and activation of latent infection. It is advised to switch to systemic treatment if infection cannot be controlled with topical treatment.

Beta Fusidic Cream contains Chlorocresol as excipients. Chlorocresol may cause allergic reactions.

Instruct patients not to smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

DRUG INTERACTIONS:

No interaction studies have been performed. Interactions with systemically administered medicinal products are considered minimal.

PREGNANCY AND LACTATION:

Pregnancy

Fusidic Acid:

No effects during pregnancy are anticipated, since systemic exposure to Fusidic Acid is negligible. Studies in animals have not shown teratogenic effects with Fusidic Acid. Limited studies in animals have shown negligible systemic absorption of topical Fusidic Acid.

Betamethasone Valerate:

There are no or limited amount of data from the use of topical Betamethasone Valerate in pregnant women. Studies in animals have shown reproductive toxicity/foetal abnormalities

Beta Fusidic Cream should not be used during pregnancy unless clearly necessary.

Breast-feeding

No effects on the breast-fed newborn/infant are anticipated since the systemic exposure of the topically applied Fusidic Acid and Betamethasone Valerate to a limited area of skin of the breast-feeding woman is negligible. Beta Fusidic Cream can be used during breast-feeding but should not be applied on the breasts to avoid accidental ingestion by the infant.

SIDE EFFECT / ADVERSE REACTIONS:

The most frequently reported adverse reaction during treatment with Fusidic Acid and Betamethasone Valerate is pruritus.

	Undesirable effects	Frequency
Immune system disorders	Hypersensitivity	Uncommon
Eye disorder	Vision, blurred*	Not known
Skin and subcutaneous tissue disorders	Dermatitis contact Eczema (condition aggravated) Skin burning sensation Pruritus Dry skin	Uncommon
	Erythema Urticaria Rash (including rash erythematous and rash generalised)	Rare
General disorders and administration site conditions	Application site pain & irritation	Uncommon
	Application site swelling & vesicles	Rare

*See also the section **Warnings and Precautions**.

Systemic undesirable class effects of corticosteroids like Betamethasone Valerate include adrenal suppression especially during prolonged topical administration (see section **Warnings and Precautions**).

Raised intra-ocular pressure, glaucoma or cataract may also occur after topical use of corticosteroids near the eyes, particularly with prolonged use and in patients predisposed to developing glaucoma and cataract (see section on **Warnings and Precautions**).

Dermatological undesirable class effects of potent corticosteroids include: Atrophy, dermatitis (including dermatitis contact and dermatitis acneiform), perioral dermatitis, skin striae, telangiectasia, rosacea, erythema, hypertrichosis, hyperhidrosis and depigmentation. Ecchymosis may also occur with prolonged use of topical corticosteroids.

Class effects for corticosteroids have been uncommonly reported for Beta Fusidic Cream as described in the frequency table above.

Paediatric population

The observed safety profile is similar in children and adults (see section on **Warnings and Precautions**).

SYMPTOMS AND TREATMENT OVERDOSE:

For topically applied Fusidic Acid, no information concerning potential symptoms and signs due to overdose administration is available. Cushing's syndrome and adrenocortical insufficiency may develop following topical application of corticosteroids in large amounts and for more than 3 weeks.

Systemic consequences of an overdose of the active substances after accidental oral intake are unlikely to occur. The amount of Fusidic Acid in one tube of Beta Fusidic Cream does not exceed the oral daily dose of systemic treatment. A single oral overdose of corticosteroids is rarely a clinical problem.

PACKING / PACK SIZE(S):

Aluminium tubes of 15 g and 50x15 g.

JAUHI UBAT DARIPADA KANAK-KANAK

KEEP OUT OF REACH OF CHILDREN

Keep Container Tightly Closed

Do Not Store Above 30°C

Protect From Light

FOR EXTERNAL USE ONLY

MANUFACTURER / PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN BHD 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, MALAYSIA

PI1PE B001

Latest Revision: 00-May 2023