

BENATUSSIL FORTE EXPECTORANT

COMPOSITION

Each 5ml syrup contains:

Diphenhydramine HCL	14.0mg
Ammonium Chloride	135.0mg
Methylparaben (as preservative)	10.0mg

PRODUCT DESCRIPTION

A brown colour liquid having raspberry flavour.

PHARMACODYNAMICS

The combination of Diphenhydramine Hydrochloride, an antihistamine and Ammonium Chloride, an expectorant, will provide rapid relief for cough which is associated with common cold influenza or allergic conditions.

PHARMACOKINETICS

Diphenhydramine hydrochloride is well absorbed from the gastrointestinal tract. Peak plasma concentrations are achieved about 1 to 4 hours after oral doses. Diphenhydramine is widely distributed throughout the body including the CNS. Diphenhydramine is highly bound to plasma proteins. Metabolism is extensive. Diphenhydramine is excreted mainly in the urine as metabolites; little is excreted as unchanged drug. The elimination half-life has been reported to range from 2.4 to 9.3 hours. Ammonium chloride is absorbed from the gastrointestinal tract.

The ammonium ion is converted into urea in the liver, the anion thus liberated into the blood and extracellular fluid causes a metabolic acidosis and decreases the pH of the urine; this is followed by transient diuresis.

INDICATION

It is indicated as an expectorant for control of coughs due to colds or allergy.

RECOMMENDED DOSAGE

Unless otherwise prescribed by the physician

Adults and children over 12 years: 5 to 10 ml every 4 or 6 hours (not to exceed 60 ml in 24 hours).

Children 6 to 12 years: 2.5 to 5 ml every 4 to 6 hours (not to exceed 30 ml in 24 hours).

Children 2 to 5 years: 2.5 ml every 4 to 6 hours (not to exceed 20 ml in 24 hours).

Do not exceed the stated dose.

ROUTE OF ADMINISTRATION

Oral administration

CONTRAINDICATIONS

Hypersensitivity to antihistamines or to any of the ingredients and Patients who are receiving Monoamine Oxidase Inhibitors (MAOIs) therapy.

WARNINGS AND PRECAUTIONS

Benatussil Forte Expectorant may enhance the sedative effect of central depressants including alcohol, barbiturates, hypnotics, sedatives and tranquilisers.

When used for treatment of cough and cold:

a) Not to be used in children less than 2 years of age.

b) To be used with caution and doctor's/ pharmacists advice in children 2 to 6 years of age.

Effects on Ability to Drive and Operate Machinery

Benatussil Forte Expectorant may cause drowsiness and the patients so affected should not drive or operate machinery.

DRUG INTERACTION

Alcohol, tranquilizers, sedative-hypnotics and CNS depressants - increase CNS depression. MAO inhibitors - increases anticholinergic effects.

PREGNANCY AND LACTATIONPregnancy

Reproduction studies in rats and rabbits given up to 5 times the human dose have shown no evidence of impaired fertility or harm to the fetus; use during pregnancy only if clearly needed.

Lactation

Contraindicated for use in nursing mothers due to the increased risk of antihistaminic side effects in newborns and infants.

ADVERSE EFFECTS

The most common side effect is sedation. Other side effects include gastro intestinal disturbances such as nausea, vomiting, diarrhoea or constipation and epigastric pain. It may produce headache, blurred vision, tinnitus, elation or depression, irritability, nightmare, anorexia, difficulty in micturition, dryness of the mouth, tightness of the chest and tingling, heaviness and weakness of the hands.

OVERDOSE AND TREATMENT

In case of accidental overdosage, professional assistance must be considered or poison control centre should be contacted immediately

STORAGE CONDITIONS

Store below 30°C. Keep container tightly closed. Protect from light

PRESENTATION

HDPE Bottles of 60ml, 90ml, 100ml and 120ml

MANUFACTURED BY

NORIPHARMA SDN. BHD. (200701034604 (792633-A)

Lot 5030, Jalan Teratai. 51/2 Mile Off Jalan Meru,
41050 Klang, Selangor, Malaysia

DATE OF REVISION

May 2023