

BENATUSSIL PAEDIATRIC EXPECTORANT

COMPOSITION

Each 5ml syrup contains :

Diphenhydramine HCL 7.0mg

Ammonium Chloride 67.5mg

Methylparaben (as preservative) 9.0mg

Propylparaben (as preservative) 0.1mg

PRODUCT DESCRIPTION

A clear, red liquid having raspberry flavour.

PHARMACODYNAMICS

The combination of Diphenhydramine Hydrochloride, an antihistamine and Ammonium Chloride, an expectorant, will provide rapid relief for cough which is associated with common cold influenza or allergic conditions.

PHARMACOKINETICS

Diphenhydramine hydrochloride is well absorbed from the gastro-intestinal tract, though high first-pass metabolism appears to affect systematic availability. Peak plasma concentrations are achieved about 1 to 4 hours after administration by mouth. A mean peak plasma-diphenhydramine concentration of 110 ng per mL has been reported after administration of 50 mg four times daily by mouth. Diphenhydramine is widely distributed throughout the body including the CNS. It crosses the placenta and has been detected in breast milk. Diphenhydramine is highly bound to plasma proteins. Metabolism is extensive. Diphenhydramine is excreted mainly in the urine as metabolites; little is excreted as unchanged drug. Excretion is almost complete within 24 hours of administration. Ammonium chloride is absorbed from the gastro-intestinal tract. The ammonium ion is converted into urea in the liver; the anion thus liberated into the blood stream and extracellular fluid causes a metabolic acidosis and decreases the pH of the urine; this is followed transient diuresis.

INDICATION

It is indicated as an expectorant for control of coughs due to colds or allergy.

RECOMMENDED DOSAGE

Unless otherwise prescribed by the physician:

To be taken 4-6 hourly.

Children 2-5 years : 5ml, not to exceed 40 ml in 24 hours.

Children 6-12 years : 5-10 ml, not to exceed 60 ml in 24 hours.

Children above 12 years : 10 - 20 ml, not to exceed 120 ml in 24 hours.

ROUTE OF ADMINISTRATION

Oral administration

CONTRAINDICATIONS

Hypersensitivity to antihistamines or to any of the ingredients and Patients who are receiving Monoamine Oxidase Inhibitors (MAOIs) therapy.

WARNINGS AND PRECAUTIONS

BENATUSSIL PAEDIATRIC may enhance the sedative effect of central depressant including alcohol, barbiturates, hypnotics, sedative and tranquilisers. It may cause drowsiness and dulling of mental

alertness. Patients undergoing treatment should not take charge of vehicles, other means of transport and machinery where loss of attention may lead to accidents.

When used for treatment of cough and cold:

- (a) Not to be used in children less than 2 years of age
- (b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age

Pregnancy and Lactation

A pregnant woman who has receiving diphenhydramine hydrochloride 150mg daily for pruritic rash gave birth to an infant who developed diarrhoea and generalised tremulousness 5 days later. The delay in appearance of withdrawal symptoms was considered to be due to absence of full activity of glucuronyl conjugating enzymes in the first few days of life.

Effects on Ability to Drive and Operate Machinery

BENATUSSIL PAEDIATRIC may cause drowsiness and the patients so affected should not drive or operate machinery.

DRUG INTERACTION

There has been a report suggesting that a reduction in temazepam, metabolism caused by diphenhydramine may have contributed to perinatal death after ingestion of this drug by the mother.

ADVERSE EFFECTS

The most common side effect is sedation.

Other side effects include gastric intestinal disturbances such as nausea, vomiting, diarrhea or constipation and epigastric pain. It may produce headache, blurred vision, tinnitus, elation or depression, irritability, nightmare, anorexia, difficulty in micturition, dryness of the mouth, tightness of chest and tingling, heaviness and weakness of the hands.

OVERDOSE AND TREATMENT

In case of accidental overdosage, professional assistance must be considered or poison control centre should be contacted immediately.

STORAGE CONDITIONS

Store below 30°C.

Keep container tightly closed.

Protect from light

PRESENTATION

HDPE Bottles of 60ml, 90ml, 100ml and 120ml

PRODUCT REGISTRATION HOLDER & MANUFACTURER

NORIPHARMA SDN. BHD. 200701034604 (792633-A)

Lot 5030, Jalan Teratai. 5½ Mile Off Jalan Meru,

41050 Klang, Selangor, Malaysia

DATE OF REVISION

May 2023