

ADE FORT INJECTION

Description and Composition

ADE FORT INJECTION is a light yellow clear oily solution. Each ml contains 500,000iu Vitamin A, 50,000iu Vitamin D₃, and 50mg Vitamin E and 1mg Chlorocresol as preservative.

Pharmacodynamics

Vitamin A (retinol) has a role in growth by intervening in the development of muscle and bone tissue. It helps maintain the integrity of epithelial tissues promoting resistance to infectious and parasitic attacks. Vitamin D₃ (cholecalciferol) is involved in phospho-calcic homeostasis and in the mineralization of bones in growth and in the adult stage. Vitamin E (alpha-tocopherol) acts as an antioxidant by protecting membrane unsaturated fatty acids against lipoperoxidation phenomena.

Pharmacokinetics

After intramuscular or subcutaneous administration, Vitamins A, D₃ and E, which are fat-soluble, are stored in hepatic tissue and in adipose tissue. The elimination of vitamin A takes place via the faecal and urinary routes in the form of derivatives. Vitamin E is little or not metabolized in the body, its elimination is mainly biliary. The elimination of vitamin D takes place via the faecal route.

Indication

For the treatment and prevention of vitamin A, D and E deficiencies in cattle, sheep and pigs.

Recommended Dose

By intramuscular injection only.

Cows: 5ml: At least twice per year and 30 days prior to calving

Calves: 1ml: During the first week after birth

Sheep: 1ml to 2ml: At least twice per year

Lambs: 0.5ml: As soon as possible after birth

Pigs: 2ml to 3ml: Sows: One injection upon placement in farrowing house and a second injection during the first week after farrowing.

Boars: One injection 1 to 2 weeks prior to breeding season.

Piglets: 0.25ml to 0.5ml: During the first week after birth.

Route of Administration

By intramuscular

Contraindication

Do not use in food-producing animals with adequate vitamin A intake due to possible accumulation in edible tissues.

Warning and Precautions

Special warning for target species:

None

Special precautions for use:

Special precautions for use in animals

Avoid carcass damage.

Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.

Maintain cleanliness at all times.

Keep needles sharp and clean. Replace frequently.

Use needles of appropriate gauge and length.

As far as possible, avoid injection of animals in wet weather or under dusty conditions.

This product must be injected only into muscle tissue. If possible, inject into muscle tissue on side of neck.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

In the event of accidental self-injection, a risk of vitamin A-related hypervitaminosis cannot be excluded. Therefore, administration should be performed with extreme caution. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Studies of vitamin A in laboratory animals have shown evidence of teratogenic effects. Therefore, this veterinary medicinal product should not be administered by pregnant women.

Interaction with Other Medicinal Product

None known.

Pregnancy and Lactation

Studies carried out on laboratory animals have shown the teratogenic effects of vitamin A. The use of the specialty during pregnancy is reserved for cases of proven deficiency.

Side Effects

None known.

Symptoms and Treatment of Overdose

None known.

Storage Condition

Store below 25°C. Protect from light.

Shelf life

3 years. Following withdrawal of the first dose, use the product within 24 hours. Any unused material should be discarded.

Withdrawal Periods

Meat: 0 days; Milk: 0 days

Packing

20ml, 50ml, 100ml, 250ml.

Revision date: 25.11.2022

	Manufacturer & Product Registration Holder:
	Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com