

SEPTIGRAS SPRAY 1% w/v

Active Ingredient:

Chlorhexidine Gluconate
1% w/v.

Product Description:

Clear colourless to slight
yellowish and odourless
solution.

Indication:

As an antiseptic
for treatment of skin lesions,
cuts, abrasions, infected wounds
and burns.

Directions of Use:

Clean and dry the affected area.
Then spray the solution
3 times a day.

Route of Administration:

Topical.

Contraindications:

Hypersensitivity to
Chlorhexidine.

Warnings and Precautions:

TOPSEAL SEPTIGRAS SPRAY
1% w/v contains
chlorhexidine. Chlorhexidine
is known to induce
hypersensitivity, including
generalized allergic
reactions and anaphylactic
shock. The prevalence
of chlorhexidine hypersensitivity
is unknown, but available
literature suggests this is likely to
be very rare. TOPSEAL SEPTIGRAS

SPRAY 1% w/v should
not be administered to anyone
with a possible history
of an allergic reaction
to chlorhexidine.

If any signs or symptoms
of a suspected hypersensitivity
reaction such as itching, skin
rash, redness, swelling, breathing
difficulties, light headedness and
rapid heart rate develop,
immediately stop using the
product. Appropriate therapeutic
countermeasures must be
instituted as clinically indicated.

For external use only.

If symptoms persist, please
consult doctor.

**Interactions with Other
Medicaments:**

The cationic nature of the
compound results in activity
being reduced by soap and
anionic compounds as they can
form insoluble salts. Anions to be
wary of include borates,
bicarbonates, carbonates,
chlorides, citrates, nitrates,
phosphates and sulphates with
due attention being paid to the
presence of hard water.

Pregnancy and Lactation:

No special precautions need to
be taken when used in
pregnancy and lactation.

Adverse Effects / Undesirable Effects:

Immune system disorders.
Frequency not known:
Hypersensitivity including anaphylactic shock.

Symptoms and Treatment of Overdose:

Chlorhexidine Gluconate taken orally is poorly absorbed into the blood. Accidental ingestion may require gastric lavage using milk, egg white, gelatine or mild soap. Employ supportive measures as appropriate. Haemolysis has been reported following ingestion or accidental intravenous administration. Haemolysis may necessitate blood transfusion.

Effects on Ability to Drive and Use Machines:
Not known

Pharmacodynamics:

Chlorhexidine is effective against a wide range of Gram negative and Gram positive vegetative bacteria, yeasts, dermatophyte fungi and lipophilic viruses. It is inactive against bacterial spores except at elevated temperatures.

Pharmacokinetics:

Chlorhexidine is cationic nature, it binds strongly to skin mucosa and other tissues and is thus very poorly absorbed. There are, as a consequence, no general pharmacological studies on chlorhexidine available and its effects on internal organs are minimal. No detectable blood levels have been found in man following oral use and percutaneous absorption, if it occurs at all, is insignificant.

Storage Conditions:
Store below 30°C. Protect from light. Keep out of reach of children.

Shelf Life:
2 years.

Dosage Form:
Topical, Spray.

Packing Size:
30 ml

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

Manufacturer:
Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor,
Malaysia.

Product Registration Holder:
AA Alive Sdn Bhd
PS-G-2, Blok Pelangi Sentral,
PSN Surian, Pelangi Damansara,
47400 Petaling Jaya, Selangor.

Date of Revision: August 2023.