

Remeco

PEPTICON

Double Action
Oral Suspension

COMPOSITION
Each 10ml of suspension contains 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate.

PRODUCT DESCRIPTION
Pepticon is an opaque, off-white creamish color, peppermint odour, viscous suspension.

PHARMACODYNAMICS
Pepticon Double Action Suspension is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate. It works in two ways:
1) Neutralising excess stomach acid to relieve the pain and discomfort
2) Forming a protective barrier over the stomach contents to soothe the burning pain in your chest

PHARMACOKINETICS
Absorption:
The mechanism of action of the medicinal product is physical and does not depend on absorption into the systemic circulation.

Distribution:
No data available.

Biotransformation:
No data available.

Elimination:
No data available.

Linear /Non-linear situation:
No data available.

INDICATION
Treatment of symptoms resulting from the reflux of acid into the oesophagus such as acid regurgitation, heartburn and indigestion, for example following meals or during pregnancy, and for symptoms of excess stomach acid (hyperacidity).

RECOMMENDED DOSAGE AND ADMINISTRAION
For oral use.

Shake well before use.

Adults and children over 12 years: Take 10-20ml (Two to four 5ml spoonfuls) after meal and at bedtime, up to four times a day.

Children under 12 years: Not be given unless recommended by your doctor.

CONTRAINDICATIONS
PEPTICON should not be used in patients with known or suspected hypersensitivity to any of the active ingredients or other excipients (parabens) contained in the suspension.

WARNINGS AND PRECAUTIONS
Each 10 ml dose has a sodium content of 124.33 mg. This should be taken into account when a highly restricted salt diet is recommended, e.g. in some cases of congestive cardiac failure and renal impairment.

Each 10 ml contains 130.12 mg calcium. Care needs to be taken in treating patients with hypercalcaemia, nephrocalcinosis and recurrent calcium renal calculi.

This medicinal product contains methyl paraben (E 218) and propyl paraben (E 216), which may cause allergic reactions (possibly delayed).

If symptoms do not improve after seven days, the clinical situation of patient should be reviewed by doctor. Avoid to use more than you should.

Like as other antacids, antacid-alginate combinations can mask more serious underlying situations.

Do not use PEPTICON in following situations:
• Patients with severe / partial renal function / failure
• Hypophosphatemic patients

It is possible for this medicine to reduction in activity for the patient who has very low gastric acid levels.
Children with renal failure or gastroenteritis are under a serious hypernatrem-ia risk.
Except medical advice, it is generally not recommended in the treatment of children under 12 years.

INTERACTIONS WITH OTHER MEDICAMENTS
It is recommended to wait 2 hours before using other medicinal products (especially H2 antihistamines, tetracyclines, digoxin, fluoroquinolone, iron salts, ketoconazole, neuroleptics, penicillamine, beta blockers (atenolol, metoprolol, propranolol), glucocorticoid, chloroquine, bisphosphonates and estramustine (for prostate gland cancer)) since the calcium carbonate in the product has antacid properties.

Since the absorption of medicine containing levothyroxine is impaired when taken together with PEPTICON, it is recommended to wait minimum 2 hours between taking these products.

Additional information on special populations
No interaction studies have been conducted.

Paediatric population:
No interaction studies have been conducted.

PREGNANCY AND LACTATION
General advice
Pregnancy category is B.

Women of childbearing potential / birth control (contraception)
No data is available that proposes additional contraceptive method while PEPTICON treatment.

There is no evidence that the effects on contraception methods.

Pregnancy
Open, controlled study conducted on 281 pregnant women showed that no significant adverse effect of sodium alginate, sodium bicarbonate and calcium carbonate combination on pregnancy or on the foetus / new-born child's health.

Based on such as studies, PEPTICON can be used the pregnancy period.

Breast feeding
Open, controlled study showed no significant adverse effect of sodium alginate, sodium bicarbonate and calcium carbonate combination on new-born child's health.
It can be used during breast feeding.

Fertility
The effect of PEPTICON on reproductive ability / fertility is unknown.

SIDE EFFECTS
Adverse reactions have been ranked under headings of frequency using the following convention: very common (≥ 1/10), common (<1/10 and ≥1/100), uncommon (<1/100 and ≥1/1000), rare (<1/1000 and ≥1/10,000), very rare (<1/10,000) and not known (cannot be estimated from the available data).

Immune System Disorders
Very rare: Allergic symptoms such as urticaria, anaphylactic or anaphylactoid reactions may develop very rarely in patients sensitive to the ingredients of the product.

Metabolism and Nutrition Disorders
Not known: Excess amount of calcium carbonate consumption can cause alkalosis, hypercalcemia, acid rebound, milk alkali syndrome or constipation. Generally, these side effects can occur when taken more than normal dose.

Respiratory, Thoracic and Mediastinal Disorders
Very rare: Respiratory effects such as bronchospasm.

Gastrointestinal Disorders
Not Known: Constipation

OVERDOSAGE
In the event of overdose, symptomatic treatment should be given. The patient may notice abdominal distension.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE
It has no effect on ability to drive and use machines.

STORAGE CONDITION
Store at room temperature below 30°C. Keep in original package, avoid from moisture. Do not keep in refrigerator or do not freeze.

PACKING
100 ml or 200 ml of amber colour glass bottle.

PRODUCT REGISTRATION HOLDER
Abio Marketing Sdn Bhd
No. 2, Jalan SS 13/5, 47500 Subang Jaya, Malaysia.

MANUFACTURER
Vefa İlaç Sanayi ve Ticaret Anonim Şirketi
Beylikdüzü OSB Mah. 2. Cadde No: 3/2 Beylikdüzü/İSTANBUL/Turkey.

REVISION DATE
December 2024

Ürün Adı	Remeco PEPTICON Double Action Oral Suspension KT		
Renk	Siyah		
Teknik Özellikler	60gr. 1.Hamur Kırımsız		
Ebat	Boy 240mm	Yükseklik	170mm