

DYNA CWC SOLUTION

MALXXXXXXXXX

DESCRIPTION:

Colourless in colour liquid solution, clear with natural odour

CONTENT:

Salicylic Acid17% w/v
Lactic Acid 5% w/v

PHARMACODYNAMIC:

Topically applied salicylic acid is keratolytic. The keratolytic activity produces desquamation by solubilising the intercellular cement in the stratum corneum resulting in the shedding of skin scales.

Lactic acid affects the keratinisation process, reducing the hyperkeratosis which is characteristic of corns, warts, and calluses. At high concentrations it can cause epidermolysis, leading to the destruction of the keratotic tissue, and in the case of warts, of the causative virus. It also has antiseptic properties.

Flexible collodion provides a viscous vehicle that allows accurate application of the active ingredients to the corn, wart or callus. It also forms a film that helps to hydrate and promote the destruction of hyperkeratotic tissue.

PHARMACOKINETICS:

Absorption

Salicylic acid is absorbed through the skin; where detectable, maximum plasma levels are found 6 to 12 hours after application. Systemic absorption of salicylic acid has been reported to range from 9% to 25% after topical application of other salicylic acid-containing preparations. The extent of absorption is variable depending on the duration of contact and the vehicle. Despite percutaneous absorption, the systemic exposure is low given the low dose topically administered to small, localised areas of hyperkeratotic tissue.

Human abdominal skin in a flow-through diffusion system was used to assess the in vitro percutaneous absorption of lactic acid. At a pH of 3, the amount of radioactivity detected in the receptor fluid, stratum corneum, epidermis, and dermis was 3.6, 6.3, 6.6, and 13.9%, respectively.

Distribution

Following percutaneous absorption, salicylic acid is distributed in the extracellular space; approximately half of which is protein bound to albumin.

Metabolism

Salicylates are metabolised in the liver by microsomal enzymes to salicyluric acid and phenolic glucuronides of salicylic acid. That which is not metabolised is excreted in the urine as unchanged salicylic acid.

Elimination

Within 24 hours of salicylic acid being absorbed and distributed in the intercellular space, approximately 95% of the absorbed dose can be recovered in the urine.

INDICATIONS:

For the removal of corns, warts and calluses.

APPLICATIONS:

Apply Dyna CWC solution twice daily directly to the affected area. Make sure the affected area is clean and dry before applying. Avoid applying to the healthy surrounding skin.

Procedure for application:

Step 1 : Soak the corn, wart and callus in warm water for 5 minutes and dried thoroughly with a clean towel.

Step 2 : Using nail file, pumice stone, emery board or coarse washcloth, carefully rub along the surface of the corn, wart and callus.

Step 3 : Apply a thin layer of Dyna CWC solution by using the dropper directly to the corn, wart and callus.

Step 4 : Allow the solution to dry thoroughly and cover with plaster.

For adults, the elderly and children aged 2 years and over. Use in children should be under the supervision of an adult.

Patients should be advised to consult a pharmacist or doctor if skin irritation develops.

ROUTE OF ADMINISTRATION:

Topical

CONTRAINDICATIONS:

Hypersensitive to the ingredients of the medication.

WARNINGS AND PRECAUTIONS:

The preparation is not recommended for prolonged use for the treatment over large areas of the body, in high concentrations, or on broken or inflamed areas of the body. Avoid contact with mouth, eyes, and other mucous membrane.

INTERACTIONS WITH OTHER MEDICAMENTS:

Topical Salicylic Acid and Lactic Acid solution may increase the absorption of other topically applied medicines. Therefore, concomitant use of Dyna CWC Solution and other topical medicines on the treated area should be avoided.

PREGNANCY AND LACTATION:

The safety of Salicylic Acid and lactic acid during pregnancy and lactation is not been established. Salicylate are excreted in human milk. Dyna CWC Solution is not recommended during pregnancy and lactation.

SIDE EFFECT / ADVERSE REACTIONS:

Salicylic acid is a mild irritation and application of salicylic acid preparation to the skin may cause dermatitis. Symptom of acute systemic salicylate poisoning have been reported after the application of salicylic acid to large areas of the body. Usage according to the recommended dose, would rarely cause overdose. Should accidental ingestion occurs, seek medical treatment immediately.

SYMPTOMS AND TREATMENT OVERDOSE:

Usage according to the recommended dose, would rarely cause overdose. Should accidental ingestion occurs, seek medical treatment immediately.

PACKING / PACK SIZE(S):

Amber glass bottle of 15ml

**JAUHKAN UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN
FOR EXTERNAL USE ONLY**

MANUFACTURER / PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN BHD 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, MALAYSIA

Latest Revision: 00-Nov 2023