

Duloxpra® Enteric Microencapsulated Capsule 30mg

Duloxpra® Enteric Microencapsulated Capsule 60mg

Active Ingredient:

Each enteric microencapsulated capsule contains:

Duloxetine	30mg (as hydrochloride)
Duloxetine	60mg (as hydrochloride)

Pharmacodynamics:

Pharmacotherapeutic group: Other antidepressants. ATC code: N06AX21

Mechanism of action

Duloxetine is a combined serotonin (5-HT) and noradrenaline (NA) reuptake inhibitor. It weakly inhibits dopamine reuptake, with no significant affinity for histaminergic, dopaminergic, cholinergic, and adrenergic receptors. It was reported that duloxetine dose-dependently increases extracellular levels of serotonin and noradrenaline in various brain areas.

Pharmacokinetics

Duloxetine is administered as a single enantiomer. Duloxetine is extensively metabolised by oxidative enzymes (CYP1A2 and the polymorphic CYP2D6), followed by conjugation.

Absorption

Duloxetine is well absorbed after oral administration, with a C_{max} occurring 6 hours post-dose. The absolute oral bioavailability of duloxetine ranged from 32% to 80%. Food delays the time to reach the peak concentration from 6 to 10 hours and it marginally decreases the extent of absorption.

Distribution

Duloxetine is approximately 96% bound to human plasma proteins. Duloxetine binds to both albumin and alpha1-acid glycoprotein. Protein binding is not affected by renal or hepatic impairment.

Metabolism

Duloxetine is extensively metabolised and the metabolites are excreted principally in urine. Both cytochromes P450-2D6 and 1A2 catalyse the formation of the two major metabolites, glucuronide conjugate of 4-hydroxy duloxetine and sulfate conjugate of 5-hydroxy, 6-methoxy duloxetine. The pharmacokinetics of duloxetine in patients who are poor metabolisers with respect to CYP2D6 has not been specifically investigated. Limited data suggest that the plasma levels of duloxetine are higher in these patients.

Elimination

The elimination half-life of duloxetine ranges from 8 to 17 hours. After an oral dose the apparent plasma clearance of duloxetine ranges from 33 to 261 l/hr.

Special Populations

Gender: Pharmacokinetic differences have been identified between males and females. However, gender-based pharmacokinetic differences do not justify the recommendation for using a lower dose for female patients.

Age: Pharmacokinetic differences have been identified between younger and elderly females (≥65 years), although the magnitude of these changes is not sufficient to justify adjustments to the dose. As a general recommendation, caution should be exercised when treating the elderly.

Renal impairment: Pharmacokinetic data on duloxetine is limited in patients with mild or moderate renal impairment.

Hepatic impairment: Moderate liver disease (Child-Pugh Class B) affected the pharmacokinetics of duloxetine. The pharmacokinetics of duloxetine and its metabolites have not been studied in patients with mild or severe hepatic insufficiency.

Breast-feeding mothers: Lactation did not influence duloxetine pharmacokinetics.

Paediatric population: The model-predicted duloxetine steady-state plasma concentrations in paediatric patients were mostly within the concentration range observed in adult patients.

Indication(s):

Duloxpra is indicated for:

- Treatment of major depressive disorder.
- Management of neuropathic pain associated with diabetic peripheral neuropathy in adults.
- Treatment of generalised anxiety disorder.

Dosage and Administration(s):

Major Depressive Disorder

The starting and recommended maintenance dose is 60 mg once daily with or without food. Dosages above 60 mg once daily, up to a maximum dose of 120 mg per day have

been evaluated from a safety perspective in clinical trials. However, there is no clinical evidence suggesting that patients not responding to the initial recommended dose may benefit from dose up-titrations.

Therapeutic response is usually seen after 2-4 weeks of treatment.

After consolidation of the antidepressive response, it is recommended to continue treatment for several months, in order to avoid relapse. In patients responding to duloxetine, and with a history of repeated episodes of major depression, further long-term treatment at a dose of 60 to 120 mg/day could be considered.

Generalised Anxiety Disorder

The recommended starting dose in patients with generalised anxiety disorder is 30 mg once daily with or without food. In patients with insufficient response the dose should be increased to 60 mg, which is the usual maintenance dose in most patients.

In patients with co-morbid major depressive disorder, the starting and maintenance dose is 60 mg once daily (please see also dosing recommendation above).

Doses up to 120 mg per day have been shown to be efficacious and have been evaluated from a safety perspective in clinical trials. In patients with insufficient response to 60 mg, escalation up to 90 mg or 120 mg may therefore be considered. Dose escalation should be based upon clinical response and tolerability.

After consolidation of the response, it is recommended to continue treatment for several months, in order to avoid relapse.

Diabetic Peripheral Neuropathic Pain

The starting and recommended maintenance dose is 60 mg daily with or without food. Dosages above 60 mg once daily, up to a maximum dose of 120 mg per day administered in evenly divided doses, have been evaluated from a safety perspective in clinical trials. The plasma concentration of duloxetine displays large inter-individual variability. Hence, some patients that respond insufficiently to 60 mg may benefit from a higher dose.

Response to treatment should be evaluated after 2 months. In patients with inadequate initial response, additional response after this time is unlikely.

The therapeutic benefit should be reassessed regularly (at least every three months)

Special Populations

Elderly

No dosage adjustment is recommended for elderly patients solely on the basis of age. However, as with any medicine, caution should be exercised when treating the elderly, especially with duloxetine 120 mg per day for major depressive disorder or generalised anxiety disorder, for which data are limited.

Hepatic Impairment

Duloxetine must not be used in patients with liver disease resulting in hepatic impairment.

Renal Impairment

No dosage adjustment is necessary for patients with mild or moderate renal dysfunction (creatinine clearance 30 to 80 ml/min). Duloxetine must not be used in patients with severe renal impairment (creatinine clearance <30 ml/min).

Paediatric population

Duloxetine should not be used in children and adolescents under the age of 18 years for the treatment of major depressive disorder because of safety and efficacy concerns.

The safety and efficacy of duloxetine for the treatment of generalised anxiety disorder in paediatric patients aged 7-17 years have not been established.

The safety and efficacy of duloxetine for the treatment of diabetic peripheral neuropathic pain has not been studied. No data are available.

Discontinuation of Treatment

Abrupt discontinuation should be avoided. When stopping treatment with duloxetine, the dose should be gradually reduced over a period of at least one to two weeks in order to reduce the risk of withdrawal reactions. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose, but at a more gradual rate.

Route of Administration(s):

Duloxpra is administered orally. It may be taken with or without food.

Contraindication(s):

- Hypersensitivity to the active substance or to any of the excipients listed.
- Concomitant use of duloxetine with non-selective, irreversible monoamine oxidase inhibitors (MAOIs) is contraindicated.
- Liver disease resulting in hepatic impairment.
- Duloxetine should not be used in combination with fluvoxamine, ciprofloxacin or enoxacin (i.e. potent CYP1A2 inhibitors) since the combination results in elevated plasma concentrations of duloxetine.
- Severe renal impairment (creatinine clearance <30 ml/min).
- The initiation of treatment with duloxetine is contraindicated in patients with uncontrolled hypertension that could expose patients to a potential risk of hypertensive crisis.

Warning and Precaution(s):

Mania and Seizures

Duloxetine should be used with caution in patients with a history of mania or a diagnosis of bipolar disorder, and/or seizures.

Mydriasis

Mydriasis has been reported in association with duloxetine, therefore, caution should be used when prescribing duloxetine to patients with increased intraocular pressure or those at risk of acute narrow-angle glaucoma.

Blood Pressure and Heart Rate

Duloxetine has been associated with an increase in blood pressure and clinically significant hypertension in some patients. Blood pressure monitoring is recommended, especially during the first month of treatment. Duloxetine should be used with caution in patients whose conditions could be compromised by an increased heart rate or by an increase in blood pressure. Caution should also be exercised when duloxetine is used with medicinal products that may impair its metabolism. For patients who experience a sustained increase in blood pressure while receiving duloxetine either dose reduction or gradual discontinuation should be considered. In patients with uncontrolled hypertension duloxetine should not be initiated.

Renal Impairment

Increased plasma concentrations of duloxetine occur in patients with severe renal impairment on haemodialysis (creatinine clearance <30 ml/min).

Serotonin syndrome

Serotonin syndrome, a potentially life-threatening condition, may occur with duloxetine treatment, particularly with concomitant use of other serotonergic agents (including SSRIs, SNRIs, tricyclic antidepressants or triptans), with agents that impair metabolism of serotonin such as MAOIs, or with antipsychotics or other dopamine antagonists that may affect the serotonergic neurotransmitter systems.

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhoea).

If concomitant treatment with duloxetine and other serotonergic agents that may affect the serotonergic and/or dopaminergic neurotransmitter systems is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases.

St John's Wort

Adverse reactions may be more common during concomitant use of duloxetine and herbal preparations containing St John's Wort (*Hypericum perforatum*).

Suicide

Major Depressive Disorder and Generalised Anxiety Disorder: Depression is associated with an increased risk of suicidal thoughts, self-harm, and suicide (suicide-related events). This risk persists until significant remission occurs. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored until such improvement occurs. It is general clinical experience that the risk of suicide may increase in the early stages of recovery.

Other psychiatric conditions for which duloxetine is prescribed can also be associated with an increased risk of suicide-related events. In addition, these conditions may be co-morbid with major depressive disorder. The same precautions observed when treating patients with major depressive disorder should therefore be observed when treating patients with other psychiatric disorders.

Patients with a history of suicide-related events or those exhibiting a significant degree of suicidal thoughts prior to commencement of treatment, are known to be at greater risk of suicidal thoughts or suicidal behaviour, and should receive careful monitoring during treatment.

Cases of suicidal thoughts and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation.

Close supervision of patients, and in particular those at high risk, should accompany medicinal product therapy, especially in early treatment and following dose changes. Patients (and caregivers of patients) should be alerted about the need to monitor for any clinical worsening, suicidal behaviour or thoughts, and unusual changes in behaviour, and to seek medical advice immediately if these symptoms present.

Diabetic Peripheral Neuropathic Pain: As with other medicinal products with similar pharmacological action (antidepressants), isolated cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation. Concerning risk factors for suicidality in depression, see above. Physicians should encourage patients to report any distressing thoughts or feelings at any time.

Use in Children and Adolescents Under 18 Years of Age

Duloxetine should not be used in the treatment of children and adolescents under the age of 18 years. Suicide-related behaviours (suicide attempts and suicidal thoughts) and hostility (predominantly aggression, oppositional behaviour, and anger) were more frequently observed in clinical trials among children and adolescents treated with antidepressants compared to those treated with placebo. If, based on clinical need, a decision to treat is nevertheless taken, the patient should be carefully monitored for the appearance of suicidal symptoms. In addition, long-term safety data in children and

adolescents concerning growth, maturation, and cognitive and behavioural development are lacking.

Suicidality in Children and Adolescents

- Antidepressants increase the risk of suicidal thinking and behavior (suicidality) in children and adolescents with major depressive disorder (MDD) and other psychiatric disorders.
- Anyone considering the use of an antidepressant in a child or adolescent for any clinical use must balance the risk of increased suicidality with the clinical need.
- Patients who are started on therapy should be observed closely for clinical worsening, suicidality, or unusual changes in behavior.
- Families and caregivers should be advised to closely observe the patient and to communicate with the prescriber.
- The indication(s) approved in paediatric for the particular drug should be clearly stated / included.

Haemorrhage

There have been reports of bleeding abnormalities, such as ecchymoses, purpura, and gastrointestinal haemorrhage, with selective serotonin reuptake inhibitors (SSRIs) and serotonin/noradrenaline reuptake inhibitors (SNRIs), including duloxetine. Duloxetine may increase the risk of postpartum haemorrhage. Caution is advised in patients taking anticoagulants and/or medicinal products known to affect platelet function (e.g., NSAIDs or acetylsalicylic acid (ASA)), and in patients with known bleeding tendencies.

Hyponatraemia

Hyponatraemia has been reported when administering duloxetine. Hyponatraemia may be due to a syndrome of inappropriate anti-diuretic hormone secretion (SIADH). Caution is required in patients at increased risk for hyponatraemia, such as elderly, cirrhotic, or dehydrated patients, or patients treated with diuretics.

Discontinuation of Treatment

Withdrawal symptoms when treatment is discontinued are common, particularly if discontinuation is abrupt.

The risk of withdrawal symptoms seen with SSRIs and SNRIs may be dependent on several factors, including the duration and dose of therapy and the rate of dose reduction. Generally, these symptoms are mild to moderate, however, in some patients they may be severe in intensity. They usually occur within the first few days of discontinuing treatment, but there have been very rare reports of such symptoms in patients who have inadvertently missed a dose. Generally, these symptoms are self-limiting and usually resolve within 2 weeks, though in some individuals they may be prolonged (2-3 months or more). It is therefore advised that duloxetine should be gradually tapered when discontinuing treatment over a period of no less than 2 weeks, according to the patient's needs.

Elderly

Caution should be exercised when treating the elderly with the maximum dosage.

Akathisia/Psychomotor Restlessness

The use of duloxetine has been associated with the development of akathisia, characterised by a subjectively unpleasant or distressing restlessness and need to move, often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.

Medicinal Products Containing Duloxetine

Duloxetine is used under different trademarks in several indications (treatment of diabetic neuropathic pain, major depressive disorder, generalised anxiety disorder and stress urinary incontinence). The use of more than one of these products concomitantly should be avoided.

Hepatitis/Increased Liver Enzymes

Cases of liver injury, including severe elevations of liver enzymes (>10 times upper limit of normal), hepatitis, and jaundice have been reported with duloxetine. Most of them occurred during the first months of treatment. The pattern of liver damage was predominantly hepatocellular. Duloxetine should be used with caution in patients treated with other medicinal products associated with hepatic injury.

Sexual dysfunction

Selective serotonin reuptake inhibitors (SSRIs)/serotonin norepinephrine reuptake inhibitors (SNRIs) may cause symptoms of sexual dysfunction. There have been reports of long-lasting sexual dysfunction where the symptoms have continued despite discontinuation of SSRIs/SNRIs.

Sucrose

Duloxpra contains sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

Interaction with Other Medicaments(s):

Monoamine Oxidase Inhibitors (MAOIs):

Due to the risk of serotonin syndrome, duloxetine should not be used in combination with non-selective, irreversible monoamine oxidase inhibitors (MAOIs) or within at least 14

days of discontinuing treatment with an MAOI. Based on the half-life of duloxetine, at least 5 days should be allowed after stopping duloxetine before starting an MAOI.

The concomitant use of duloxetine with selective, reversible MAOIs, like moclobemide, is not recommended. The antibiotic linezolid is a reversible non-selective MAOI and should not be given to patients treated with duloxetine.

Inhibitors of CYP1A2: Because CYP1A2 is involved in duloxetine metabolism, concomitant use of duloxetine with potent inhibitors of CYP1A2 is likely to result in higher concentrations of duloxetine. Therefore, duloxetine should not be administered in combination with potent inhibitors of CYP1A2 like fluvoxamine.

CNS Medicinal Products: Caution is advised when duloxetine is taken in combination with other centrally-acting medicinal products or substances, including alcohol and sedative medicinal products (e.g., benzodiazepines, morphinomimetics, antipsychotics, phenobarbital, sedative antihistamines).

Serotonergic agents: In rare cases, serotonin syndrome has been reported in patients using SSRIs/SNRIs concomitantly with serotonergic agents. Caution is advisable if duloxetine is used concomitantly with serotonergic agents like SSRIs, SNRIs, tricyclic antidepressants like clomipramine or amitriptyline, MAOIs like moclobemide or linezolid, St John's Wort (*Hypericum perforatum*) or triptans, tramadol, pethidine, and tryptophan.

Effect of Duloxetine on Other Medicinal Products

Medicinal products metabolised by CYP1A2: The pharmacokinetics of theophylline, a CYP1A2 substrate, were not significantly affected by co-administration with duloxetine (60 mg twice daily).

Medicinal products metabolised by CYP2D6: Duloxetine is a moderate inhibitor of CYP2D6. Caution is advised if duloxetine is co-administered with medicinal products that are predominantly metabolised by CYP2D6 (risperidone, tricyclic antidepressants [TCAs], such as nortriptyline, amitriptyline, and imipramine), particularly if they have a narrow therapeutic index (such as flecainide, propafenone, and metoprolol).

Oral contraceptives and other steroidal agents: Reported studies demonstrate that duloxetine does not induce the catalytic activity of CYP3A. Specific drug interaction studies have not been performed.

Anticoagulants and antiplatelet agents: Caution should be exercised when duloxetine is combined with oral anticoagulants or antiplatelet agents due to a potential increased risk of bleeding attributable to a pharmacodynamic interaction. Furthermore, increases in INR values have been reported when duloxetine was co-administered to patients treated with warfarin.

Effects of Other Medicinal Products on Duloxetine

Antacids and H₂ antagonists: Co-administration of duloxetine with aluminium- and magnesium-containing antacids, or duloxetine with famotidine, had no significant effect on the rate or extent of duloxetine absorption after administration of a 40 mg oral dose.

Inducers of CYP1A2: Population pharmacokinetic analyses have shown that smokers have almost 50% lower plasma concentrations of duloxetine compared with non-smokers.

Pregnancy and Lactation(s):

Fertility

Duloxetine had no effect on male fertility, and effects in females were only evident at doses that caused maternal toxicity.

Pregnancy

There are no adequate data on the use of duloxetine in pregnant women. Studies in animals have shown reproductive toxicity at systemic exposure levels (AUC) of duloxetine lower than the maximum clinical exposure.

The potential risk for humans is unknown.

Epidemiological data have suggested that the use of SSRIs in pregnancy, particularly in late pregnancy, may increase the risk of persistent pulmonary hypertension in the newborn (PPHN). Although no studies have investigated the association of PPHN to SNRI treatment, this potential risk cannot be ruled out with duloxetine taking into account the related mechanism of action (inhibition of the re-uptake of serotonin).

Observation data indicate an increased risk (less than 2-fold) of postpartum haemorrhage following SSRI/SNRI exposure within the month prior to birth. As with other serotonergic medicinal products, discontinuation symptoms may occur in the neonate after maternal duloxetine use near term. Discontinuation symptoms seen with duloxetine may include hypotonia, tremor, jitteriness, feeding difficulty, respiratory distress and seizures. The majority of cases have occurred either at birth or within a few days of birth. Duloxetine should be used in pregnancy only if the potential benefit justifies the potential risk to the foetus. Women should be advised to notify their physician if they become pregnant, or intend to become pregnant, during therapy.

Breast-Feeding

Duloxetine is very weakly excreted into human milk. However, as the safety of duloxetine in infants is not known, the use of duloxetine while breast-feeding is not recommended.

Side Effect(s):

a. Summary of the safety profile

The most commonly reported adverse reactions in patients treated with duloxetine were nausea, headache, dry mouth, somnolence and dizziness. However, the majority of common adverse reactions were mild to moderate; they usually started early in therapy, and most tended to subside even as therapy was continued.

b. Summary of adverse reactions

Infections and infestations

Uncommon: Laryngitis

Immune system disorders

Rare: Anaphylactic reaction, hypersensitivity disorder

Endocrine disorders

Rare: Hypothyroidism

Metabolism and nutrition disorders

Common: Decreased appetite

Uncommon: Hyperglycaemia (reported especially in diabetic patients)

Rare: Dehydration, hyponatraemia, SIADH

Psychiatric disorders

Common: Insomnia, agitation, libido decreased, anxiety, orgasm abnormal, abnormal dreams

Uncommon: Suicidal ideation, sleep disorder, bruxism, disorientation, apathy

Rare: Suicidal behaviour, mania, hallucinations, aggression and anger

Nervous system disorders

Very common: Headache, somnolence

Common: Dizziness, lethargy, tremor, paraesthesia

Uncommon: Myoclonus, akathisia, nervousness, disturbance in attention, dysgeusia, dyskinesia, restless legs syndrome, poor quality sleep

Rare: Serotonin syndrome, convulsion, psychomotor restlessness, extra-pyramidal symptoms

Eye disorders

Common: Blurred vision

Uncommon: Mydriasis, visual impairment

Rare: Glaucoma

Ear and labyrinth disorders

Common: Tinnitus

Uncommon: Vertigo, ear pain

Cardiac disorders

Common: Palpitations

Uncommon: Tachycardia, supraventricular arrhythmia (mainly atrial fibrillation)

Vascular disorders

Common: Blood pressure increased, flushing

Uncommon: Syncope, hypertension, orthostatic hypotension, peripheral coldness

Rare: Hypertensive crisis

Respiratory, thoracic and mediastinal disorders

Common: Yawning

Uncommon: Throat tightness, epistaxis

Rare: Interstitial lung disease, eosinophilic pneumonia

Gastrointestinal disorders

Very common: Nausea, dry mouth

Common: Constipation, diarrhoea, abdominal pain, vomiting, dyspepsia, flatulence

Uncommon: Gastrointestinal haemorrhage, gastroenteritis, eructation, gastritis, dysphagia

Rare: Stomatitis, haematochezia, breath odour, microscopic colitis

Hepato-biliary disorders

Uncommon: Hepatitis, elevated liver enzymes (ALT, AST, alkaline phosphatase), acute liver injury

Rare: Hepatic failure, jaundice

Skin and subcutaneous tissue disorders

Common: Sweating increased, rash

Uncommon: Night sweats, urticaria, dermatitis contact, cold sweat, photosensitivity reactions, increased tendency to bruise

Rare: Stevens-Johnson Syndrome, angioneurotic oedema

Very rare: Cutaneous vasculitis

Musculoskeletal and connective tissue disorders

Common: Musculoskeletal pain, muscle spasm

Uncommon: Muscle tightness, muscle twitching

Rare: Trismus

Renal and urinary disorders

Common: Dysuria, Pollakiuria

Uncommon: Urinary retention, urinary hesitation, nocturia, polyuria, urine flow decreased

Rare: Urine odour abnormal

Reproductive system and breast disorders

Common: Erectile dysfunction, ejaculation disorder, ejaculation delayed

Uncommon: Gynaecological haemorrhage, menstrual disorder, sexual dysfunction, testicular pain

Rare: Menopausal symptoms, galactorrhoea, hyperprolactinaemia, postpartum haemorrhage

General disorders and administration site conditions

Common: Falls, fatigue

Uncommon: Chest pain, feeling abnormal, feeling cold, thirst, chills, malaise, feeling hot, gait disturbance

Investigations:

Common: Weight decrease

Uncommon: Weight increase, blood creatine phosphokinase increased, blood potassium increased

Rare: Blood cholesterol increased

Effects on Ability to Drive and Use Machine:

No studies on the effects on the ability to drive and use machines have been performed. Duloxetine may be associated with sedation and dizziness. Patients should be instructed that if they experience sedation or dizziness they should avoid potentially hazardous tasks such as driving or operating machinery.

Symptoms and Treatment of Overdose(s):

Cases of overdoses, alone or in combination with other medicinal products, with duloxetine doses of 5400 mg were reported. Some fatalities have occurred, primarily with mixed overdoses, but also with duloxetine alone at a dose of approximately 1000 mg. Signs and symptoms of overdose (duloxetine alone or in combination with other medicinal products) included somnolence, coma, serotonin syndrome, seizures, vomiting and tachycardia.

No specific antidote is known for duloxetine, but if serotonin syndrome ensues, specific treatment (such as with cyproheptadine and/or temperature control) may be considered. A free airway should be established. Monitoring of cardiac and vital signs is recommended, along with appropriate symptomatic and supportive measures. Gastric lavage may be indicated if performed soon after ingestion or in symptomatic patients. Activated charcoal may be useful in limiting absorption. Duloxetine has a large volume of distribution and forced diuresis, haemoperfusion, and exchange perfusion are unlikely to be beneficial.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Shelf Life(s):

30 months from the date of manufacture.

Product Description and Packing(s)

Duloxpra E.M. Capsule 30mg:

A size #3 dark blue cap and white body hard gelatin capsule, cap and body are imprinted with a mark "YSP" on one side and "DUCp" on the other side.

Available in 10's x 10 Alu-Alu blister pack.

Duloxpra E.M. Capsule 60mg:

A size #1 dark blue cap and light green body hard gelatin capsule, cap and body are imprinted with a mark "YSP" on one side and "DUCp6" on the other side.

Available in 10's x 10 Alu-Alu blister pack.



Y. S. P.

Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679