
UNIDINE

Gargle and Mouthwash (Povidone Iodine 1% w/v)

Name and Strength of Active Ingredient:

Povidone Iodine	10.00mg
(Equivalent to 1mg of available iodine)	
Ethanol 96%	71.21mg

Product Description:

A deep brown liquid

Pharmacodynamics:

UNIDINE gargle and mouthwash contains povidone-iodine which has broad spectrum germicidal activity. The germicidal activity is maintained in the presence of blood, pus, serum and necrotic tissue. Povidone Iodine kills bacteria, viruses, fungi, spores, and protozoa.

Pharmacokinetics:

Not Applicable

Indications:

- For the relief, prevention, and treatment of painful infection of the mouth and pharynx e.g., pharyngitis, oral moniliasis, tonsillitis, aphthous ulcers, stomatitis, gingivitis, and other inflammatory conditions.
- For oral hygiene prior to and after oral and dental surgery.
- For minor irritation of the mouth and throat.
- Helps to reduce or eliminate mouth odour.

Recommended Dose:

Use full strength or diluted with an equal volume of water. Gargle or rinse for at least 30 seconds. Repeat every 6 hours or as directed.

Route of administration:

For External Use Only. Not To Be Swallowed.

Contraindications:

Hypersensitivity to povidone-Iodine. People with thyroid disorder. Application to a large areas of broken skin should be avoided.

Warning and Precautions:

For external use only. Do not swallow.
Discontinue use in rare case of local sensitivity

Interactions with Other Medicaments:

Use with concurrent lithium therapy has been shown to exhibit additive hypothyroidic effects. Absorption of iodine from povidone iodine through either intact skin or broken skin may interfere with thyroid function tests.

Pregnancy and Lactation:

Iodine freely crosses the placenta and is secreted in breast milk. Povidone Iodine should not be used during pregnancy.

Adverse Effects:

Local irritation, skin burns and sensitivity reactions have been reported rarely. Anaphylactic reactions, anaphylactoid reactions and anaphylactic shock have been reported uncommonly with products containing povidone-iodine or povidone. Excess iodine can produce goitre and hypothyroidism or hyperthyroidism. Such effects have occasionally been seen with extensive or prolonged use of povidone iodine. Other effects that have been reported are metabolic acidosis and acute renal failure.

Symptoms and Treatment of Overdose:

Deliberate or accidental ingestion of large quantities of povidone iodine will result in high blood concentrations of iodine and gastrointestinal corrosive effects including vomiting, diarrhea and abdominal pain. Systemic toxicity may result in shock, hypotension, tachycardia, fever, metabolic acidosis and renal impairment. Symptomatic and supportive treatment should be started with special attention to monitoring electrolyte balance, renal function, thyroid function and liver function. Haemodialysis effectively clears iodine and should be employed in severe cases of iodine poisoning particularly if renal failure is present. Continuous venovenous hemodiafiltration is less effective than haemodialysis.

Storage Condition:

Store below 30°C. Protect from light.

Dosage form and packaging available

100ml, 200ml and 500ml brown HDPE bottle with red HDPE screw cap

120ml Amber PET bottle with red HDPE screw cap

Registration Number:

XXXX

Manufacturer & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas

Pulau Pinang, Malaysia.

Date of Revision:

11 August 2023