

Product Name: DIO-1	Market: Malaysia	Mfg. Location: Unit-3
----------------------------	-------------------------	------------------------------

Mfg. Location: Unit-3

Size: 260 x 120 mm
Colour: P P Black C

ARTWORK APPROVAL FORM

Product Name: DIO-1

Market: Malaysia

Mfg. Location: Unit-3

260 mm

derivatives.

patients with end-stage renal disease with GFR (Glomerular Filtration Rate) below 15 ml/min.

Warnings and Precautions

Alcohol

At therapeutic doses, no clinically significant interactions have been demonstrated with alcohol (for a blood alcohol level of 0.5 g/L). Nevertheless, precaution is recommended if alcohol is taken concomitantly.

Increased risk of urinary retention

Caution should be taken in patients with predisposition factors of urinary retention (e.g. spinal cord lesion, prostatic hyperplasia) as cetirizine may increase the risk of urinary retention.

Patients at risk of convulsions

Caution in epileptic patients and patients at risk of convulsions is recommended.

Skin reactions

Pruritus and/or urticaria may occur when cetirizine is stopped, even if those symptoms were not present before treatment initiation. In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

Children

The use of the film-coated tablet formulation is not recommended in children aged less than 6 years since this formulation does not allow for appropriate dose adaptation. It is recommended to use a paediatric formulation of cetirizine.

Allergy skin tests

Allergy skin tests are inhibited by antihistamines and a wash-out period of 3 days is recommended before performing them.

Food

The extent of absorption of cetirizine is not reduced with food, although the rate of absorption is decreased.

Activities requiring mental alertness

In clinical trials, the occurrence of somnolence has been reported in some patients taking Cetirizine. Due caution should therefore be exercised when driving a car or operating potentially dangerous machinery.

Interaction with other medicinal products and other forms of interaction

Due to the pharmacokinetic, pharmacodynamic and tolerance profile of cetirizine, no interactions are expected with this antihistamine. Actually, neither pharmacodynamic nor significant pharmacokinetic interaction was reported in drug-drug interactions studies performed, notably with pseudoephedrine or theophylline (400 mg/day).

The extent of absorption of cetirizine is not reduced with food, although the rate of absorption is decreased. In sensitive patients, the concurrent use of alcohol or other CNS depressants may cause additional reductions in alertness and impairment of performance, although cetirizine does not potentiate the effect of alcohol (0.5 g/L blood levels).

Pregnancy and Lactation

Pregnancy

For cetirizine prospectively collected data on pregnancy outcomes do not suggest potential for maternal or foetal/embryonic toxicity above background rates.

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/fetal development, parturition or postnatal development. Caution should be exercised when prescribing to pregnant women.

Breast-feeding

Cetirizine is excreted in human milk at concentrations representing 25% to 90% those measured in plasma, depending on sampling time after administration. Therefore, caution should be exercised when prescribing cetirizine to lactating women.

Fertility

Limited data is available on human fertility but no safety concern has been identified.

Animal data show no safety concern for human reproduction.

Effects on ability to drive and use machines

Objective measurements of driving ability, sleep latency and assembly line performance have not demonstrated any clinically relevant effects at the recommended dose of 10 mg.

However, patients who experience somnolence should refrain from driving, engaging in potentially hazardous activities or operating machinery. They should not exceed the recommended dose and should take their response to the medicinal product into account.

Undesirable Effects Undesirable effects are described according to MedDRA System Organ Class and by estimated frequency based on post-marketing experience.

Frequencies are defined as follows: Very common

(≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000); very rare (<1/10,000), not known (cannot be estimated from the available data)

Blood and lymphatic disorders:

Very rare: thrombocytopenia

Immune system disorders:

Rare: hypersensitivity

Very rare: anaphylactic shock

Metabolism and nutrition disorders:

Not known: increased appetite

Psychiatric disorders:

Uncommon: agitation

Rare: aggression, confusion, depression, hallucination, insomnia

Very rare: tics

Not known: suicidal ideation, nightmare

Nervous system disorders:

Uncommon: paraesthesia

Rare: convulsions

Very rare: dysgeusia, syncope, tremor, dystonia, dyskinesia

Not known: amnesia, memory impairment

Eye disorders:

Very rare: accommodation disorder, blurred vision, oculogyration

Ear and labyrinth disorders:

Not known: vertigo

Cardiac disorders:

Rare: tachycardia

Gastro-intestinal disorders:

Uncommon: diarrhoea

Hepatobiliary disorders:

Rare: hepatic function abnormal (increased transaminases, alkaline phosphatase, γ-GT and bilirubin)

Not known: hepatitis

Skin and subcutaneous tissue disorders:

Uncommon: pruritus, rash

Rare: urticaria

Very rare: angioneurotic oedema, fixed drug eruption

Not known: acute generalized exanthematous pustulosis

Musculoskeletal and connective tissue disorders

Not known: arthralgia, myalgia

Renal and urinary disorders:

Very rare: dysuria, enuresis

Not known: urinary retention

General disorders and administration site conditions:

Uncommon: asthenia, malaise

Rare: oedema

Investigations:

Rare: weight increased

Description of selected adverse reactions

After discontinuation of cetirizine, pruritus (intense itching) and/or urticaria have been reported.

Overdose and treatment

Symptoms

Symptoms observed after an overdose of cetirizine are mainly associated with CNS effects or with effects that could suggest an anticholinergic effect.

Adverse events reported after an intake of at least 5 times the recommended daily dose are: confusion, diarrhoea, dizziness, fatigue, headache, malaise, mydriasis, pruritus, restlessness, sedation, somnolence, stupor, tachycardia, tremor, and urinary retention.

Management

There is no known specific antidote to cetirizine.

Should overdose occur, symptomatic or supportive treatment is recommended.

Cetirizine is not effectively removed by haemodialysis.

Dosage Forms and Packaging Available

Tablets are available in Alu-PVC/PVDC blister pack

Pack size: 10 x 10 Alu-PVC/PVDC & 50 x 10 Alu-PVC/PVDC

Storage Condition

Store at a temperature not exceeding 30°C.

Keep the medicine out of reach of children

Date of revision: June-2024

Manufactured by:
UNISON PHARMACEUTICALS PVT. LTD.
Unit-III, C-7,8,9, Steel Town, Opp. Nova
Petrochemicals, Village Moraiya, Sanand,
Ahmedabad, Gujarat 382213, INDIA

Product registration holder:
HEALOL PHARMACEUTICALS SDN BHD,
74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju,
53300, Kuala Lumpur, Malaysia

XXXXXXXXXX

120 mm

Font Type : Times New Roman
Font Size : 7 Brand Font Size : 8

Size: 260 x 120 mm
Colour: P P Black C

DATE: 04/06/2024 VERSION: 00

INSERT/PIL/OUTSERT SPECIFICATION

Artwork Code XXXXXXXXXXXX

Open Size 260 x 120 mm

Close Size 120 x 32.5 mm

Floding Condition Folded

Type of Paper Bible

GSM of Paper 40 gsm

Colour of Paper White

Colour of Matter P P Black C

Pharmacode 714

Any other special process N.A.

Old Artwork Code N.A.

Old Pharmacode N.A.

Reference Change Control no. N.A.

of Plant N.A

Review & Approved By Contract Giver/Customer/Authority/MA Holder (If Applicable) :

Sign & Date :

Name :

Designation :

Department :

F:\Artworks\General Export\MALAYSIA\HEALOL PHARMA\DIO-1\XXXXXXXXXX_DIO-1_PIL(Draft)_Malaysia_04-06-2024.cdr