

POMPEZO 40mg

Lyophilized Powder for Solution for Injection or Infusion

QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains Esomeprazole Sodium 42.5mg equivalent to 40 mg esomeprazole.

For a full list of excipients see “List of Excipients.”

PHARMACEUTICAL FORM

Lyophilized powder for solution for injection or infusion.

White colored lyophilized powder.

After reconstitution/dilution: clear, colorless or light-yellow colored solution

CLINICAL PARTICULARS

Therapeutic indications

Pompezo for injection and infusion is indicated for

- Gastric antisecretory treatment when the oral route is not possible, such as:
 - gastroesophageal reflux disease in patients with esophagitis and/or severe symptoms of reflux.
 - healing of gastric ulcers associated with NSAID therapy.
 - prevention of gastric and duodenal ulcers associated with NSAID therapy, in patients at risk.
- Prevention of rebleeding following therapeutic endoscopy for acute bleeding gastric or duodenal ulcers.

Posology and method of administration

Gastric antisecretory treatment when the oral route is not possible

Patients who cannot take oral medication may be treated parenterally with 20-40 mg once daily.

Patients with reflux oesophagitis should be treated with 40 mg once daily. Patients treated symptomatically for reflux disease should be treated with 20 mg once daily.

For healing of gastric ulcers associated with NSAID therapy the usual dose is 20 mg once daily.

For prevention of gastric and duodenal ulcers associated with NSAID therapy, patients at risk should be treated with 20 mg once daily.

Usually, the intravenous treatment duration is short and transfer to oral treatment should be made as soon as possible. Treatment with Esomeprazole IV can be given for up to 10 days as part of a full treatment period for the specified indications.

Prevention of rebleeding of gastric and duodenal ulcers

Following therapeutic endoscopy for acute bleeding gastric or duodenal ulcers, 80 mg should be administered as a bolus infusion over 30 minutes, followed by a continuous intravenous infusion of 8 mg/h given over 3 days (72 hours).

The parenteral treatment period should be followed by oral acid suppression therapy.

Method of administration:

For preparation of reconstituted solution, please see section Instruction for use and handling.

*Injection*40 mg dose

5 ml of the reconstituted solution (8 mg/ml) should be given as an intravenous injection over a period of at least 3 minutes

20 mg dose

2.5 ml or half of the reconstituted solution (8 mg/ml) should be given as an intravenous injection over a period of at least 3 minutes. Any unused solution should be discarded.

*Infusion*40 mg dose

The reconstituted solution should be given as an intravenous infusion over a period of 10 to 30 minutes.

20 mg dose

Half of the reconstituted solution should be given as an intravenous infusion over a period of 10 to 30 minutes. Any unused solution should be discarded.

80 mg bolus dose

The reconstituted solution should be given as a continuous intravenous infusion over 30 minutes.

8 mg/hour dose

The reconstituted solution should be given as a continuous intravenous infusion over a period of 71.5 hours (calculated rate of infusion of 8 mg/h).

Children and adolescents

Esomeprazole should not be used in children since no data is available.

Impaired renal function

Dose adjustment is not required in patients with impaired renal function. Due to limited experience in patients with severe renal insufficiency, such patients should be treated with caution.

Impaired hepatic function

GERD: Dose adjustment is not required in patients with mild to moderate liver impairment. For patients with severe liver impairment, a maximum daily dose of 20 mg Esomeprazole should not be exceeded.

Bleeding ulcers: Dose adjustment is not required in patients with mild to moderate liver impairment. For patients with severe liver impairment, following an initial bolus dose of 80 mg Esomeprazole for infusion, a continuous intravenous infusion dose of 4 mg/h for 71.5 hours may be sufficient.

Elderly

Dose adjustment is not required in the elderly.

Contraindications

It is contraindicated in patients with hypersensitivity to esomeprazole or benzimidazole derivatives or to any of the excipients in the formula.

Esomeprazole should not be used concomitantly with nelfinavir

Special warnings and precautions for use

In the presence of any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, dysphagia, haematemesis or melaena) and when gastric ulcer is suspected or present, malignancy should be excluded, as treatment with Pompezo may alleviate symptoms and delay diagnosis.

Gastrointestinal infections

Treatment with proton pump inhibitors may lead to slightly increased risk of gastrointestinal infections such as Salmonella and Campylobacter.

Absorption of vitamin B12

Esomeprazole, as all acid-blocking medicines, may reduce the absorption of Vitamin B12 (cyanocobalamin) due to hypo- or achlorhydria. This should be considered in patients with reduced body stores or risk factors for reduced Vitamin B12 absorption on long-term therapy.

Hypomagnesaemia

Severe hypomagnesaemia has been reported in patients treated with PPI like esomeprazole for at least three months, and in most cases for a year. Serious manifestations of hypomagnesaemia such as fatigue, tetany, delirium, convulsions, dizziness, and ventricular arrhythmia can occur but they may begin insidiously and be overlooked. In most affected patients, hypomagnesaemia improved after magnesium replacement and discontinuation of the PPI.

For patients expected to be on prolonged treatment or who take PPI with digoxin or drugs that may cause hypomagnesaemia (e.g. diuretics), health care professionals should consider measuring magnesium levels before starting PPI treatment and periodically during treatment

Fracture

Proton pump inhibitors, especially if used in high doses and over long durations (>1 year), may modestly increase the risk of hip, wrist and spine fracture, predominantly in the elderly or in presence of other recognised risk factors. Observational studies suggest that proton pump inhibitors may increase the overall risk of fracture by 10-40%. Some of this increase may be due to other risk factors. Patients at risk of osteoporosis should receive care according to current clinical guidelines and they should have an adequate intake of vitamin D and calcium.

Clostridium Difficile Diarrhea

Published observational studies suggest that PPI therapy may be associated with an increased risk of Clostridium difficile associated diarrhea, especially in hospitalized patients. This diagnosis should be considered for diarrhea that does not improve. Patients should use the lowest dose and shortest duration of PPI therapy appropriate to the condition being treated.

Vitamin B12 Deficiency

Daily treatment with any acid-suppressing medications over a long period of time (e.g., longer than 3 years) may lead to malabsorption of cyanocobalamin (vitamin B12) caused by hypo- or achlorhydria. Rare reports of cyanocobalamin deficiency occurring with acid-suppressing therapy have been reported in the literature. This diagnosis should be considered if clinical symptoms consistent with cyanocobalamin deficiency are observed.

Subacute Cutaneous Lupus Erythematosus (SCLE)

Proton pump inhibitors are associated with very infrequent cases of subacute cutaneous lupus erythematosus (SCLE). If lesions occur, especially in sun-exposed areas of the skin, and if accompanied by arthralgia, the patient should seek medical help promptly and the health care professional should

consider stopping esomeprazole. SCLE after previous treatment with a proton pump inhibitor may increase the risk of SCLE with other proton pump inhibitors.

Serious cutaneous adverse reactions (SCARs)

Serious cutaneous adverse reactions (SCARs) such as erythema multiforme (EM), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS), which can be life-threatening, have been reported very rarely in association with esomeprazole treatment.

Patients should be advised of the signs and symptoms of the severe skin reaction EM/SJS/TEN/DRESS and should seek medical advice from their physician immediately when observing any indicative signs or symptoms.

Esomeprazole should be discontinued immediately upon signs and symptoms of severe skin reactions and additional medical care/close monitoring should be provided as needed.

Re-challenge should not be undertaken in patients with EM/SJS/TEN/DRESS.

Combination with other medicinal products

Co-administration of esomeprazole with atazanavir is not recommended. If the combination of atazanavir with a proton pump inhibitor is judged unavoidable, close clinical monitoring is recommended in combination with an increase in the dose of atazanavir to 400 mg with 100 mg of ritonavir; esomeprazole 20 mg should not be exceeded.

Esomeprazole is a CYP2C19 inhibitor. When starting or ending treatment with esomeprazole, the potential for interactions with drugs metabolised through CYP2C19 should be considered. An interaction is observed between clopidogrel and esomeprazole. The clinical relevance of this interaction is uncertain. As a precaution, concomitant use of esomeprazole and clopidogrel should be discouraged.

Interference with laboratory tests

Increased Chromogranin A (CgA) level may interfere with investigations for neuroendocrine tumours. To avoid this interference, esomeprazole treatment should be stopped for at least 5 days before CgA measurements. If CgA and gastrin levels have not returned to reference range after initial measurement, measurements should be repeated 14 days after cessation of PPI treatment.

Regular Surveillance

Patients on proton pump inhibit treatment (particularly those treated for long term) should be kept under regular surveillance.

Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

Effects of esomeprazole on the pharmacokinetics of other drugs

Medicinal products with pH dependent absorption

Gastric acid suppression during treatment with esomeprazole and other PPIs might decrease or increase the absorption of medicinal products drugs with a gastric pH dependent absorption. As with other medicinal products that decrease the intragastric acidity, the absorption of medicinal products, such as ketoconazole, itraconazole and erlotinib and the absorption of digoxin can increase during treatment with esomeprazole. Concomitant treatment with omeprazole (20 mg daily) and digoxin in healthy subjects increased the bioavailability of digoxin by 10% (up to 30% in two out of ten subjects). Digoxin toxicity

has been rarely reported. However, caution should be exercised when esomeprazole is given at high doses in elderly patients. Therapeutic drug monitoring of digoxin should then be reinforced.

Omeprazole has been reported to interact with some protease inhibitors. The clinical importance and the mechanisms behind these reported interactions are not always known. Increased gastric pH during omeprazole treatment may change the absorption of the protease inhibitors. Other possible interaction mechanisms are via inhibition of CYP 2C19. For atazanavir and nelfinavir, decreased serum levels have been reported when given together with omeprazole and concomitant administration is not recommended. Co-administration of omeprazole (40 mg once daily) with atazanavir 300 mg/ritonavir 100 mg to healthy volunteers resulted in a substantial reduction in atazanavir exposure (approximately 75% decrease in AUC, C_{max} and C_{min}). Increasing the atazanavir dose to 400 mg did not compensate for the impact of omeprazole on atazanavir exposure. The co-administration of omeprazole (20 mg qd) with atazanavir 400 mg/ritonavir 100 mg to healthy volunteers resulted in a decrease of approximately 30% in the atazanavir exposure as compared with the exposure observed with atazanavir 300 mg/ritonavir 100 mg qd without omeprazole 20 mg qd. Co-administration of omeprazole (40 mg qd) reduced mean nelfinavir AUC, C_{max} and C_{min} by 36–39 % and mean AUC, C_{max} and C_{min} for the pharmacologically active metabolite M8 was reduced by 75–92%. For saquinavir (with concomitant ritonavir), increased serum levels (80–100%) have been reported during concomitant omeprazole treatment (40 mg qd). Treatment with omeprazole 20 mg qd had no effect on the exposure of darunavir (with concomitant ritonavir) and amprenavir (with concomitant ritonavir). Treatment with esomeprazole 20 mg qd had no effect on the exposure of amprenavir (with and without concomitant ritonavir). Treatment with omeprazole 40 mg qd had no effect on the exposure of lopinavir (with concomitant ritonavir). Due to the similar pharmacodynamic effects and pharmacokinetic properties of omeprazole and esomeprazole, concomitant administration with esomeprazole and atazanavir is not recommended and concomitant administration with esomeprazole and nelfinavir is contraindicated.

Drugs metabolised by CYP2C19

Esomeprazole inhibits CYP2C19, the major esomeprazole metabolising enzyme. Thus, when esomeprazole is combined with drugs metabolised by CYP2C19, such as diazepam, citalopram, imipramine, clomipramine, phenytoin etc., the plasma concentrations of these drugs may be increased and a dose reduction could be needed. Concomitant oral administration of 30 mg esomeprazole resulted in a 45% decrease in clearance of the CYP2C19 substrate diazepam. Concomitant oral administration of 40 mg esomeprazole and phenytoin resulted in a 13% increase in trough plasma levels of phenytoin in epileptic patients. It is recommended to monitor the plasma concentrations of phenytoin when treatment with esomeprazole is introduced or withdrawn. Omeprazole (40 mg once daily) increased voriconazole (a CYP2C19 substrate) C_{max} and AUC_τ by 15% and 41%, respectively.

Concomitant oral administration of 40 mg esomeprazole to warfarin-treated patients in a clinical trial showed that coagulation times were within the accepted range. However, postmarketing of oral esomeprazole, a few isolated cases of elevated INR of clinical significance have been reported during concomitant treatment. Monitoring is recommended when initiating and ending concomitant esomeprazole treatment during treatment with warfarin or other coumarine derivatives.

In healthy volunteers, concomitant oral administration of 40 mg esomeprazole and cisapride resulted in a 32% increase in area under the plasma concentration-time curve (AUC) and a 31% prolongation of elimination half-life ($t_{1/2}$) but no significant increase in peak plasma levels of cisapride. The slightly prolonged QTc interval observed after administration of cisapride alone, was not further prolonged when cisapride was given in combination with esomeprazole.

Esomeprazole has been shown to have no clinically relevant effects on the pharmacokinetics of amoxicillin or quinidine.

No in vivo interaction studies have been performed with the high dose intravenous regimen (80mg+8mg/h). The effect of esomeprazole on drugs metabolised by CYP2C19 may be more pronounced during this regimen, and patients should be monitored closely for adverse effects, during the 3-day intravenous treatment period.

Results from studies in healthy subjects have shown a pharmacokinetic (PK)/ pharmacodynamic (PD) interaction between clopidogrel (300 mg loading dose/75 mg daily maintenance dose) and esomeprazole (40 mg p.o.daily) resulting in decreased exposure to the active metabolite of clopidogrel by an average of 40% and resulting in decreased maximum inhibition of (ADP induced) platelet aggregation by an average of 14%.

When clopidogrel was given together with a fixed dose combination of esomeprazole 20 mg + ASA 81 mg compared to clopidogrel alone in a study in healthy subjects there was a decreased exposure by almost 40% of the active metabolite of clopidogrel. However, the maximum levels of inhibition of (ADP induced) platelet aggregation in these subjects were the same in the clopidogrel and the clopidogrel + the combined (esomeprazole + ASA) product groups.

Inconsistent data on the clinical implications of a PK/PD interaction of esomeprazole in terms of major cardiovascular events have been reported from both observational and clinical studies. As a precaution concomitant use of clopidogrel should be discouraged.

Omeprazole as well as esomeprazole act as inhibitors of CYP 2C19. Omeprazole, given in doses of 40 mg to healthy subjects in a cross-over study, increased C_{max} and AUC for cilostazol by 18% and 26% respectively, and one of its active metabolites by 29% and 69%, respectively.

Unknown mechanism

Concomitant administration of esomeprazole has been reported to increase the serum levels of tacrolimus.

When given together with proton pump inhibitors, methotrexate levels have been reported to increase in some patients. In high-dose methotrexate administration a temporary withdrawal of esomeprazole may need to be considered.

Effects of other drugs on the pharmacokinetics of esomeprazole

Esomeprazole is metabolised by CYP2C19 and CYP3A4. Concomitant oral administration of esomeprazole and a CYP3A4 inhibitor, clarithromycin (500 mg b.i.d.), resulted in a doubling of the exposure (AUC) to esomeprazole. Concomitant administration of esomeprazole and a combined inhibitor of CYP2C19 and CYP 3A4 may result in more than doubling of the esomeprazole exposure. The CYP2C19 and CYP3A4 inhibitor voriconazole increased omeprazole AUC_τ by 280%. A dose adjustment of esomeprazole is not regularly required in either of these situations. However, dose adjustment should be considered in patients with severe hepatic impairment and if long-term treatment is indicated.

Drugs known to induce CYP2C19 or CYP3A4 or both (such as rifampicin and St. John's wort) may lead to decreased esomeprazole serum levels by increasing the esomeprazole metabolism.

Pregnancy and lactation

For esomeprazole limited data on exposed pregnancies are available. Animal studies with esomeprazole do not indicate direct or indirect harmful effects with respect to embryonal/fetal development. Animal studies with the racemic mixture do not indicate direct or indirect harmful effects with respect to pregnancy, parturition or postnatal development. Caution should be exercised when prescribing Esomeprazole to pregnant women.

It is not known whether esomeprazole is excreted in human breast milk. No studies in lactating women have been performed. Therefore, Esomeprazole should not be used during breastfeeding.

Effects on ability to drive and use machines

It is not likely to affect the ability to drive or use machines.

Undesirable effects

The following adverse drug reactions have been identified or suspected for esomeprazole administered orally or intravenously and post-marketing when administered orally. The reactions are classified according to frequency: very common; common; uncommon; rare; very rare; not known.

Blood and lymphatic system disorders

Rare: Leukopenia, thrombocytopenia.

Very rare: Agranulocytosis, pancytopenia.

Immune system disorders

Rare: Hypersensitivity reactions e.g. fever, angioedema and anaphylactic reaction/shock.

Metabolism and nutrition disorders

Uncommon: Peripheral edema.

Rare: Hyponatraemia.

Not known: Hypomagnesaemia; severe hypomagnesaemia may result in hypocalcaemia, Hypomagnesaemia may also result in hypokalaemia.

Psychiatric disorders

Uncommon: Insomnia.

Rare: Agitation, confusion, depression.

Very rare: Aggression, hallucination.

Nervous system disorders

Common: Headache.

Uncommon: Dizziness, paresthesia, somnolence.

Rare: Taste disturbance

Eye disorders

Rare: Blurred vision.

Ear disorders

Uncommon: Vertigo.

Respiratory, thoracic and mediastinal disorders

Rare: Bronchospasm.

Gastrointestinal disorders

Common: Abdominal pain, constipation, diarrhea, flatulence, nausea/vomiting, fundic gland polyps (benign)

Uncommon: Dry mouth.

Rare: Stomatitis, gastrointestinal candidiasis.

Not known: Microscopic colitis.

Hepatobiliary disorder

Uncommon: Increased liver enzymes.

Rare: Hepatitis with or without jaundice.

Very rare: Hepatic failure, encephalopathy in patients with pre-existing liver disease

Skin and subcutaneous tissue disorders

Common: Administration site reactions*

Uncommon: Dermatitis, pruritus, rash, urticaria

Rare: Alopecia, photosensitivity

Very rare: Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (TEN)

Not known: Subacute cutaneous lupus erythematosus (SCLE)

Musculoskeletal, connective tissue and bone disorders

Uncommon: Fracture of the hip, wrist, or spine

Rare: Arthralgia, myalgia

Very rare: Muscular weakness

Renal and urinary system disorders

Very rare: Interstitial nephritis.

Reproductive system and breast disorders

Very rare: Gynecomastia.

General disorders and administration site conditions

Rare: malaise, increased sweating

*Administration site reactions have mainly been observed in a study with high-dose exposure over 3 days (72 hours).

Irreversible visual impairment has been reported in isolated cases of critically ill patients who have received omeprazole (the racemate) intravenous injection, especially at high doses, but no causal relationship has been established.

Other possible side effects include *Clostridium difficile* associated diarrhoea and Vitamin B12 deficiency (with unknown frequency).

Overdose and treatment

There is very limited experience to date with deliberate overdose. The symptoms described in connection with an oral dose of 280 mg were gastrointestinal symptoms and weakness. Single oral doses of 80 mg esomeprazole and intravenous doses of 308 mg esomeprazole over 24 hours were uneventful. No specific antidote is known. Esomeprazole is extensively plasma protein bound and is therefore not readily dialyzable. As in any case of overdose, treatment should be symptomatic and general supportive measures should be utilised.

PHARMACOLOGICAL PROPERTIES**Pharmacodynamic properties**

Pharmacotherapeutic group: Proton pump inhibitor

ATC Code: A02B C05

Esomeprazole is the S-isomer of omeprazole and reduces gastric acid secretion through a specific targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. Both the R- and S-isomer of omeprazole have similar pharmacodynamic activity.

Site and mechanism of action

Esomeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the secretory canaliculi of the parietal cell, where it inhibits the enzyme H⁺, K⁺-ATPase - the acid pump and inhibits both basal and stimulated acid secretion.

Effect on gastric acid secretion:

After 5 days of oral dosing with 20 mg and 40 mg of esomeprazole, intragastric pH above 4 was maintained for a mean time of 13 hours and 17 hours, respectively over 24 hours in symptomatic GERD patients. The effect is similar irrespective of whether esomeprazole is administered orally or intravenously.

Using AUC as a surrogate parameter for plasma concentration, a relationship between inhibition of acid secretion and exposure has been shown after oral administration of esomeprazole.

During intravenous administration of 80 mg esomeprazole as a bolus infusion over 30 minutes followed by a continuous intravenous infusion of 8 mg/h for 23.5 hours, intragastric pH above 4, and pH above 6 was maintained for a mean time of 21 hours, and 11-13 hours, respectively, over 24 hours in healthy individuals.

Therapeutic effects of acid inhibition:

Healing of reflux esophagitis with esomeprazole 40 mg occurs in approximately 78% of patients after 4 weeks, and in 93% after 8 weeks of oral treatment.

Other effects related to acid inhibition

During treatment with antisecretory medicinal products serum gastrin increases in response to the decreased acid secretion. Also, CgA increases due to decreased gastric acidity. The increased CgA level may interfere with investigations for neuroendocrine tumours. \

Available published evidence suggests that proton pump inhibitors should be discontinued between 5 days and 2 weeks prior to CgA measurements. This is to allow CgA levels that might be spuriously elevated following PPI treatment to return to reference range.

An increased number of ECL cells possibly related to the increased serum gastrin levels, have been observed in both children and adults during long term treatment with orally administered esomeprazole. The findings are considered to be of no clinical significance.

During long-term oral treatment with antisecretory drugs gastric glandular cysts have been reported to occur at a somewhat increased frequency. These changes are a physiological consequence of pronounced inhibition of acid secretion, are benign and appear to be reversible.

Decreased gastric acidity due to any means including proton pump inhibitors, increases gastric counts of bacteria normally present in the gastrointestinal tract. Treatment with proton pump inhibitors may lead to slightly increased risk of gastrointestinal infections such as Salmonella and Campylobacter and, in hospitalised patients, possibly also *Clostridium difficile*.

Pharmacokinetic properties

Distribution

The apparent volume of distribution at steady state in healthy individuals is approximately 0.22 l/kg body weight. Esomeprazole is 97% plasma protein bound.

Metabolism and excretion

Esomeprazole is completely metabolised by the cytochrome P450 system (CYP). The major part of the metabolism of esomeprazole is dependent on the polymorphic CYP2C19, responsible for the formation of the hydroxy- and desmethyl metabolites of esomeprazole. The remaining part is dependent on another specific isoform, CYP3A4, responsible for the formation of esomeprazole sulphone, the main metabolite in plasma.

The parameters below reflect mainly the pharmacokinetics in individuals with a functional CYP2C19 enzyme, extensive metabolisers.

Total plasma clearance is about 17 l/h after a single dose and about 9 l/h after repeated administration. The plasma elimination half-life is about 1.3 hours after repeated once-daily dosing. Total exposure (AUC) increases with repeated administration of esomeprazole. This increase is dose-dependent and results in a non-linear dose-AUC relationship after repeated administration. This time- and dose dependency is due to a decrease of first pass metabolism and systemic clearance probably caused by inhibition of the CYP2C19 enzyme by esomeprazole and/or its sulphone metabolite.

Esomeprazole is completely eliminated from plasma between doses with no tendency for accumulation during once-daily administration.

Following repeated doses of 40 mg administered as intravenous injections, the mean peak plasma concentration is approx. 13.6 micromol/l. The mean peak plasma concentration after corresponding oral doses is approx. 4.6 micromol/l. A smaller increase (of approx. 30%) can be seen in the total exposure after intravenous administration compared to oral administration.

There is a dose-linear increase in total exposure following intravenous administration of esomeprazole as a 30-minute infusion (40 mg, 80 mg or 120 mg) followed by a continuous infusion (4 mg/h or 8 mg/h) over 23.5 hours.

The major metabolites of esomeprazole have no effect on gastric acid secretion. Almost 80% of an oral dose of esomeprazole is excreted as metabolites in the urine, the remainder in the faeces. Less than 1% of the parent drug is found in urine.

Special patient populations

Approximately 2.9±1.5% of the population lacks a functional CYP2C19 enzyme and is called poor metabolisers. In these individuals the metabolism of esomeprazole is probably mainly catalysed by CYP3A4. After repeated once-daily administration of 40 mg oral esomeprazole, the mean total exposure was approximately 100% higher in poor metabolisers than in individuals with a functional CYP2C19 enzyme (extensive metabolisers). Mean peak plasma concentrations were increased by about 60%. Similar differences have been seen for intravenous administration of esomeprazole. These findings have no implications for the dosology of esomeprazole.

The metabolism of esomeprazole is not significantly changed in elderly individuals (71-80 years of age).

Following a single oral dose of 40 mg esomeprazole the mean total exposure is approximately 30% higher in females than in males. No gender difference is seen after repeated once-daily administration. Similar differences have been observed for intravenous administration of esomeprazole. These findings have no implications for the dosology of esomeprazole.

The metabolism of esomeprazole in patients with mild to moderate liver dysfunction may be impaired. The metabolic rate is decreased in patients with severe liver dysfunction resulting in a doubling of the total exposure of esomeprazole. Therefore, a maximum dose of 20 mg should not be exceeded in GERD patients with severe dysfunction. For patients with bleeding ulcers and severe liver impairment, following an initial bolus dose of 80 mg, a maximum continuous intravenous infusion dose of 4 mg/h for 71.5 hours may be sufficient. Esomeprazole or its major metabolites do not show any tendency to accumulate with once daily dosing.

No studies have been performed in patients with decreased renal function. Since the kidney is responsible for the excretion of the metabolites of esomeprazole but not for the elimination of the parent compound, the metabolism of esomeprazole is not expected to be changed in patients with impaired renal function.

PHARMACEUTICAL PARTICULARS

List of excipients

Disodium edetate

Sodium hydroxide (for pH adjustment)

Incompatibilities

This medicinal product should not be used with other medicinal products except those mentioned in Instruction for use and handling.

Shelf life

Unopened vial: 24 months

Shelf-life after reconstitution/dilution

Use the reconstituted solution within 12 hours when stored below 30°C.

Keep out of reach of children.

Special precautions for storage

POMPEZO IV should be stored in outer package for keeping from light.

Vials could be stored for 24 hours when they are removed from the box at normal room light.

Store at room temperature below 30°C.

Nature and content of container

8 ml transparent type I glass vial, 20 mm lyophilized bromobutyl grey single notched stopper, 20 mm transparent flip off cap.

Package type: 1 vial in cardboard box.

INSTRUCTIONS FOR USE AND HANDLING

The reconstituted solution should be inspected visually for particulate matter and discoloration prior to administration. Only clear solution should be used. For single use only.

If the entire reconstituted content of the vial is not required, any unused solution should be discarded in accordance with local requirements.

Injection 40 mg

A solution for injection 8 mg/ml is prepared by adding 5 ml of 0.9% sodium chloride for intravenous use to the esomeprazole 40mg vial.

The reconstituted solution for injection is clear, colorless or light-yellow colored.

Infusion 40 mg

A solution for infusion is prepared by dissolving the content of one vial with esomeprazole 40mg in up to 100 ml of 0.9% sodium chloride for intravenous use.

The reconstituted solution for infusion is clear, colorless or light-yellow colored.

Infusion 80 mg

A solution for infusion is prepared by dissolving the contents of two vials of esomeprazole 40 mg in up to 100 ml of 0.9% sodium chloride for intravenous use.

The reconstituted solution for infusion is clear, colorless or light-yellow colored.

MANUFACTURER

Mefar Ilac San A.S.
Ramazanoglu Mah. Ensar Cad.
No:20 TR-34906 Kurtkoy Pendik
Istanbul Turkey

PRODUCT REGISTRATION HOLDER

Medispec (M) Sdn Bhd
B-1-07, Jalan SS 25/22,
Taman Mayang,
47301 Petaling Jaya, Selangor, Malaysia

DATE OF REVISION OF THE TEXT

May 2024