




Please change the following settings in Acrobat Reader to view this file correctly.
For PC: Edit > Preferences > Categories: Page Display > Use Overprint Preview > Select “Always”
For MAC: Adobe Reader > Preferences > Categories: Page Display > Use Overprint Preview > Select “Always”

RE-TYPESET ARTWORK. Due to the nature of the files received, this artwork has been re-typeset.



inova
pharmaceuticals

ARTWORK

Job No. : JB-03171

UIC : A9207

SKU : 000000 105894

V3

20.09.24 - V0 - Artwork build - Koragot

25.09.24 - V1 - Amend - Suphapon

11.10.24 - V2 - Amend - Suphapon

26.02.25 - V3 - Amend - Suphapon

A9207_DURO-TUSS AX CHESTY LIQ 60ML, 120ML (SHARED)

Component : LEAFLET

Country : MY

Dieline Ref: PP-5134

EMO: INTERPHIL

Printer: JARDIS PRINTING CORP

G247 File Name:

105382_A9207_DURO-TUSS AX CHESTY LIQ 60ML, 120ML (SHARED)_MY_LT_V3

Colour

100%

Black



Guide

PLEASE NOTE

This HI RES PDF is not to be used as Final Artwork

Supplier information may be present on the packaging

EMO/Printing Vendor:

Please share the Artwork PDF to Printer for checks on:

- Colours - No. of colours, Sequence, CMYK/PMS
- Text - Line Thickness, Font Size
- Dieline - Size, Eyemark, Print Direction
- Coding - Size, Location, VFA, Format Bkg Colour

Moving forward when possible any additional information and technical check required by EMO on the Artwork BEFORE APPROVING.

2.88 mm 11.00 pt
Helvetica Neue LT Std 77 Bold Condensed
Helvetica Neue LT Std 57 Condensed
1.41 mm 5.50 pt Helvetica Neue LT Std 57 Condensed
1.54 mm 6.00 pt Helvetica Neue LT Std 57 Condensed

DURO-TUSS AX CHESTY COUGH Liquid

Each 5mL of DURO-TUSS AX CHESTY COUGH Liquid contains 30mg of Ambroxol hydrochloride. It also contains Methyl Paraben 0.12%w/v and Propyl Paraben 0.04%w/v as preservatives.
Product Description
A clear, very light yellow to colorless syrup with orange butterscotch flavour.
Pharmacodynamics

Interactions with Other Medicaments

If DURO-TUSS AX CHESTY COUGH Liquid is used in combination with antitussives in patients with pre-existing respiratory diseases such as cystic fibrosis or bronchiectasis who are affected by mucus hypersecretion, a reduced cough reflex may lead to (serious) accumulation of mucus.

Pregnancy and Lactation

Pregnancy

Ambroxol hydrochloride crosses the placental barrier. Non-clinical studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/ foetal development, labour and delivery or postnatal development. Extensive clinical experience after the 28th week of pregnancy has shown no evidence of harmful effects on the foetus. However, the usual precautions concerning the administration of any medicine during pregnancy should be observed. The use of DURO-TUSS AX CHESTY COUGH Liquid is not recommended during the first trimester of pregnancy in particular.

Breast-feeding

Ambroxol has been shown to pass into breast milk in animal studies. Use while breast-feeding is not recommended.

Fertility

Non-clinical studies do not indicate direct or indirect harmful effects with respect to fertility.

Side Effects

The following categories are normally used when reporting the frequencies of side effects:

Very common:	≥ 1/10
Common:	≥ 1/100 to < 1/10
Uncommon:	≥ 1/1,000 to < 1/100
Rare:	≥ 1/10,000 to < 1/1,000
Very rare:	< 1/10,000
Not known:	Frequency cannot be estimated from the available data

Immune system disorders

Rare: Hypersensitivity reactions

Not known: Anaphylactic reactions including anaphylactic shock, angioedema and pruritus

Skin and subcutaneous tissue disorders

Rare: Rash, urticaria

Not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Nervous system disorders

Common: Dysgeusia (e.g. changed taste)

Gastrointestinal disorders

Common: Nausea, oral hypoaesthesia

Uncommon: Vomiting, diarrhoea, dyspepsia, abdominal pain, dry mouth

Rare: Dry throat

Very rare: Sialorrhoea Respiratory, thoracic and mediastinal disorders

Common: Pharyngeal hypoaesthesia

Not known: Dyspnoea (as a symptom of a hypersensitivity reaction)

General disorders and administration site conditions

Uncommon: Fever, mucous membrane reactions

Symptoms and Treatment of Overdose

No specific symptoms of overdose have been reported to date. The symptoms observed in cases of accidental overdose or medication error are consistent with the known side effects at the recommended dose and may require symptomatic treatment.

Effects on Ability to Drive and Use Machine

There is no evidence from postmarketing data for an effect on the ability to drive and use machines. Studies on the effects on the ability to drive and use machines have not been performed.

Storage Conditions

Store below 30°C.

Dosage forms and packaging available

Available in pack size of 60ml and 120ml bottles

Manufactured by

Interphil Laboratories, Inc.
Canlubang Industrial Estate, Bo. Pittland
Cabuyao, Laguna 4025, Philippines

Date of revision

16 September 2024

A9207