

Peppermint

GastroRelief

Reflux Double Action Suspension

Active Ingredients:
Each 10ml contains -
Sodium Alginate 500mg
Sodium Bicarbonate 213mg
Calcium Carbonate 325mg
Preserved with Methyl Hydroxybenzoate 0.2%w/v, Propyl Hydroxybenzoate 0.02%w/v.

Presentations:
PET bottles of 120ml and 250ml.
Glass bottles of 120ml, 200ml and 250ml.

Product Descriptions:
Peppermint flavoured and white to off-white coloured suspension.

Pharmacology:
Pharmacodynamics:
GastroRelief Reflux Double Action Suspension is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate. On ingestion, the suspension reacts rapidly with gastric acid to form a raft of alginic acid gel having a near neutral pH and which floats on the stomach contents effectively (up to 4 hours) impeding gastro-oesophageal reflux. In severe cases, the raft itself may be refluxed into the oesophagus, in preference to the stomach contents, and exert a demulcent effect. Calcium carbonate neutralises gastric acid to provide fast relief from indigestion and heartburn. This effect is increased by the addition of sodium bicarbonate which also has a neutralising action. The total neutralising capacity of the product at the lowest dose of 10 mL dose is approximately 10 mEqH⁺.

Pharmacokinetics:
The mode of action of the medicinal product is physical and does not depend on absorption into the systemic circulation.

Indications:
For the relief of pain and discomfort resulting from gastro-oesophageal reflux (GERD) such as acid regurgitation, heartburn and indigestion, which can occur after meals or during pregnancy, and for symptoms of excess stomach acid (hyperacidity).

Dosage and Administration: For oral use only.
Adults and children above 12 years:
Take 10 to 20ml after meals and at bedtime, up to 4 times a day.

Children under 12 years:
Should be given only on medical advice.

Shake well before use.

Symptoms and Treatments of Overdose:
Symptoms are likely to be minor, some abdominal distension may be noticed. In the event of overdose, symptomatic treatment should be given.

Contraindications:
Contraindicated in patients with known hypersensitivity to sodium alginate, sodium bicarbonate, calcium carbonate or to any of the excipients listed in the product.
Should not be used in patients with moderate or severe renal insufficiency.

Precautions:
It should be used with caution in patients with congestive cardiac failure and renal impairment.
If symptoms persist after 7 days, consult the doctor.
Patient may feel bloated if taken too much. Please consult the doctor or pharmacist.
This product has no or negligible influence on the ability to drive and use machines.

Use in Pregnancy and Lactation:
Use in Pregnancy:
GastroRelief Reflux Double Action can be used during pregnancy.
It is recommended to limit the treatment duration as much as possible.
Use in Lactation:
No effects of the active substances have been shown in breastfed newborns/ infants of treated mothers.
This product can be used during breast-feeding.

Interactions with Other Medicaments:
Due to the presence of calcium and carbonates, a time-interval of 2 hours should be considered between intake of this product and the administration of other medicinal products, especially tetracyclines, fluoroquinolones, iron salts, thyroid hormones, chloroquine, bisphosphonates, and estramustine.

Undesirable Effects:
Hypersensitivity reactions and respiratory effects have been reported very rarely.

Storage:
Store below 30°C. Protect from light and moisture.
Keep out of reach of children.
Jauhkan daripada capaian kanak-kanak.

Shelf Life:
18 months from the date of manufacture.

Manufacturer and Product Registration Holder
WINWA MEDICAL SDN. BHD.
198101009677(75794-A)
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

MAL24076030X

Date of Revision: 02 December 2025

X.LDP-I.02

Size: 75mm x 100mm

Dark Blue (Pantone 2767U)

Checked by	
Approved by	