

**GAVELOX ANTACID
TABLET (DOUBLE STRENGTH)**

Active Ingredients:

Each tablet contains:
Dried Aluminium Hydroxide 400mg
Magnesium Hydroxide 400mg
Simethicone 40mg

Presentation:

Bottles of 12's.

Product Descriptions:

White coloured, round and flat chewable tablet with cool mint flavour. Source of lactose is bovine.

Pharmacology:

GaVelox Antacid Tablet when taken by mouth is non systemic. Aluminium Hydroxide and Magnesium Hydroxide react with hydrochloric acid and convert to their respective chloride salts in the stomach. They reduce the gastric acidity by neutralizing secreted gastric acid. The increased gastric pH and decreased pepsin production helps in the healing of peptic ulcers. The clinical use of Simethicone is based on its antifoam properties. Silicone antifoams spread on the surface of aqueous liquids, forming a film of low surface tension and thus causing collapse of foam bubbles. Simethicone reportedly allows mucus-surrounded gas bubbles in the gastrointestinal (GI) tract to coalesce and be expelled.

Aluminium Hydroxide reacts with Hydrochloric Acid in the stomach to form Aluminium Chloride. Absorbed Aluminium is eliminated in the urine. The majority of Aluminium remains in the GI tract and forms insoluble poorly absorbed Aluminium salts including hydroxide, phosphate, carbonate and fatty acid derivatives, which are excreted in the faeces. Magnesium Hydroxide reacts with hydrochloric acid in the stomach to produce Magnesium Chloride and water. About 30% of the Magnesium salts are absorbed from the small intestine and excreted in the urine (absorbed fraction) and the faeces (unabsorbed fraction). Simethicone is physiologically inert; it does not appear to be absorbed from the GI tract or to interfere with gastric secretion or absorption of nutrients. Following oral administration, the drug is excreted unchanged in faeces.

Indications:

For the relief of gastric pain and discomforts of peptic ulcer, heartburn, dyspepsia (indigestion) and acid reflux. It also helps to relieve gastrointestinal conditions caused by excessive gas (flatulence).

Dosage and Administration:

Adults and children above 12 years: Chew 1 to 2 tablets 4 times a day. Chew well before swallowing, preferably ½ hour before meals or 2 hours after meals and at bedtime or as directed by doctor.

Symptoms and Treatments of Overdose:

Symptoms of overdose may include gastrointestinal irritation, diarrhoea and constipation. Severe poisoning may produce hypermagnesaemia.

Treatment consists of the intravenous administration of Calcium Gluconate injection 10% in a dose of 10 to 20ml to counteract respiratory depression or heart block. If renal function is normal, adequate fluids should be given to assist removal of Magnesium from the body. Dialysis may be necessary in patients with renal impairment or severe hypermagnesaemia. Treatment should be symptomatic and supportive.

Contraindications:

Contraindicated in patients with known hypersensitivity to Aluminium Hydroxide, Magnesium Hydroxide or Simethicone. Should not be used in patients who severely debilitated or suffering from kidney failure.

Precautions:

It should be used with caution in patients with renal insufficiency. Unsuitable for phenylketonurics. If symptoms persist, please consult a doctor.

Use in Pregnancy and Lactation:

Antacids generally regarded as safe. No data is available to suggest any harmful effects of Aluminium Hydroxide and Magnesium Hydroxide associated with Simethicone in pregnant women. A small amount of Aluminium Hydroxide and Magnesium Hydroxide may be secreted in breast milk and the concentration is too small to be harmful. Secretion of Simethicone in the breast milk of nursing mothers has not been established.

Interactions with Other Medicaments:

May interfere with or reduce the absorption of anticholinergic agents, Barbiturates, Digoxin, Quinidine, Warfarin and Vitamins. Antacid can interfere with the absorption of iron preparations, Tetracyclines or other antibiotics. Hence, antacids should not be taken within 2 hours of other medications because the effectiveness of the other medications may be affected.

Undesirable Effects:

Antacids may occasionally cause constipation, diarrhoea, nausea and vomiting.

Storage:

Store below 30°C. Protect from light and moisture. Keep container tightly closed. Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Shelf Life:

2 years from the date of manufacture.

Registration No.: MAL00000000XC

X.TApGV-I.01

Date of Revision: 26 June 2024

Aetos

Product Registration Holder
AETOS PHARMA SDN. BHD.
No. 66B, Tingkat 2, Jalan Cerdas,
Taman Connaught,
56000 Kuala Lumpur,
Malaysia.

Manufacturer
WINWA MEDICAL SDN. BHD.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Size: 75mm x 115mm
■ Dark Blue (Pantone 2767U)