

MICO-GUARD 10MG/ML TOPICAL SPRAY

Description and Composition

MICO-GUARD 10MG/ML TOPICAL SPRAY is a pale yellow to light yellow clear solution. Each ml contains 10mg Miconazole (as nitrate) as its active ingredient and 5mg Phenoxyethanol as its preservative.

Pharmacodynamics

Topical miconazole has activity against dermatophytes and yeasts, which is effective treatment for *Malassezia* dermatitis. Lotions, sprays and creams are generally used for localized lesions associated with *Malassezia* or dermatophytes. Topical miconazole products are generally ineffective (or minimally effective) when used alone for dermatophytosis; adjunctive systemic treatment is usually required. Miconazole's actions are a result of altering permeability of fungal cellular membranes and interfering with peroxisomal and mitochondrial enzymes, leading to intracellular necrosis. Miconazole products are fungicidal with repeated application.

Pharmacokinetics

Absorption: There is little absorption through skin or mucous membranes when miconazole nitrate is applied topically.

Distribution: Data available indicates that over 90% of miconazole is bound to plasma proteins. Penetration into the cerebrospinal fluid and sputum is poor but miconazole diffuses well into infected joints. Animal studies indicate that the drug crossed the placenta but it is unknown whether miconazole nitrate is excreted in breast milk.

Metabolism: Miconazole is metabolised in the liver to inactive metabolites. Miconazole has an important first pass effect. Hepatic metabolism of miconazole is by O-dealkylation and oxidative N-dealkylation.

Indication

For the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum* and *Trichophyton mentagrophytes*.

Recommended Dose

Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

Route of Administration

For external use only.

Contraindications

There are no known contraindications to this drug when used as directed.

Warning and Precautions

Avoid contact with eyes. Wash hands after application or wear gloves when applying.

Interactions with Other Medicaments

None known.

Pregnancy and Lactation

Animal studies indicate that the drug crossed the placenta but it is unknown whether miconazole nitrate is excreted in breast milk.

Side Effects

Skin irritation is possible, but unlikely to occur, but in very inflamed, eroded to ulcerated skin, the pledgets, towelettes, spray containing alcohol can be severely irritating.

Symptoms and Treatment of Overdose

Since over 90% of miconazole is bound in plasma proteins, forced diuresis or dialysis is of no value in case of miconazole overdosage.

Storage Condition

Store below 30°C.

Shelf life

3 years.

Discard 6 months after first opening.

Packing

50ml, 100ml, 200ml, 500ml

Revised date: 27.9.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com