

Fusisure Pack Insert new ok
Malaysia Export Date: 31-07-2024, 02-08-2024

145 mm

148 mm

80 mm

FUSISURE (Fusidic acid cream 2%w/w)

Composition:
Fusidic acid BP 2%w/w
Preservative:
Potassium sorbate NF 0.27%w/w

Cream
White or almost White colored cream.

Pack Sizes: 5g, 15g.

Route of Administration: For External Use Only

Indications
Fusidic acid 2% is indicated for treatment of skin infections caused by staphylococci, streptococci, Propionibacterium acnes, Corynebacterium minutissimum, and other organisms sensitive to fusidic acid. The most important indications being:

Impetigo	Boils	Carbuncles	Paronychia	
Infected wounds	Sycosis barbae	Hidradenitis	Erythrasma	Folliculitis

Fusidic acid 2% is an invisible, non-staining preparation, which is cosmetically acceptable for treatment of face and scalp infections.

Application
Fusidic acid 2% is applied to the affected area 2–3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

Pharmacodynamics:
Pharmacotherapeutic group: other antibiotics for topical use, ATC code: D06AX01.

Mechanism of action:
Fusidic acid belongs to a unique group of antibiotics, the fusidanes, which act to inhibit bacterial protein synthesis by blocking the lengthening of factor G. This is to prevent it from associating with ribosomes and GTP, thus preventing energy supply to the synthesis process. As it is the only type of drug available in this family of drugs, there have been no reports of cross resistance to fusidic acid.

Clinical efficacy and safety is inserted:

Resistance mechanism(s)
Resistance for fusidic acid can vary geographically and information about local resistance patterns should be obtained through a local microbiology laboratory. In general, resistance occurs in 1-10 % of Staphylococcus aureus and 10-20 % of coagulase negative staphylococci. Cross-resistance between Fusidic acid 20mg/g cream and other antibiotics has not been reported.

Sensitivity:
The sensitivity of organisms to fusidic acid is based on the in vitro sensitivity and plasma concentrations that are achieved after systemic therapy. Local treatment causes higher peak concentrations as compared to plasma. However, it is not known how the kinetics of the cream after local application may change the effectiveness of the cream.

Pharmacokinetics:

Absorption:
In Vitro studies show that fusidic acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to fusidic acid and the condition of the skin.

Elimination:
Fusidic acid is excreted mainly in the bile with little excreted in the urine.

Contraindications
Infections caused by non-susceptible organisms. Patients with hypersensitivity to any of the ingredients.

Warning and Precautions
Bacterial resistance has been reported to occur with the use of fusidic acid. As with all antibiotics, extended or recurrent use may increase the risk of developing antibiotic resistance.
Fusidic acid should not be used in infections caused by non-susceptible organisms, in particular, Pseudomonas aeruginosa, Extended or recurrent use may increase the risk of developing contact sensitization. When Fusidic acid 20 mg/g cream is used on the face, care should be taken to avoid the eyes, because fusidic acid can cause irritation of the conjunctiva.

XX-000000
000000-00

Item-numb
See details

Space for text: 125 x 128 mm.

148 nm

000000-00

000000-XX

Excipients
Fusidic acid 20 mg/g cream contains butylhydroxyanisole, cetyl alcohol and potassium sorbate which may cause local skin reactions (e.g. contact dermatitis). Butylhydroxyanisole may also cause irritation to the eyes and mucous membranes.

There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that fusidic acid is devoid of teratogenic effects. Safety in nursing mothers has not been established.

Pregnancy:
No effects during pregnancy are anticipated, since systemic exposure to topically applied fusidic acid/sodium fusidate is negligible. Topical fusidic acid cream can be used during pregnancy.

Breast-feeding: No effects on the breastfed new-born/infant are anticipated since the systemic exposure of the breast-feeding woman is negligible. Topical Fusidic acid cream can be used during breast-feeding but it is recommended to avoid applying topical Fusidic acid cream on the breast.

Fusidic acid 2% is remarkably well tolerated, and there is an extremely low frequency of hypersensitivity reactions. The estimation of the frequency of undesirable effects is based on a pooled analysis of data from clinical trials and from spontaneous reporting. The most frequently reported adverse reactions during treatment are various skin reactions such as pruritus and rash, followed by application site conditions such as pain and irritation, which all occurred in less than 1% of patients. Hypersensitivity and angioedema have been reported. Undesirable effects are listed by MedDRA System Organ Class (SOC) and the individual undesirable effects are listed starting with the most frequently reported according to the following frequency convention:

System organ class	Frequency	Undesirable effects
Immune system disorders	Rare	Hypersensitivity
Eye disorders	Rare	Conjunctivitis
Skin and subcutaneous tissue disorders	Uncommon	Dermatitis (incl. contact dermatitis, eczema) Rash* Pruritus, Erythema
	Rare	Angioedema Urticaria Blister
General disorders and administration site conditions	Uncommon	Application site pain (incl. skin burning sensation), Application site irritation

Pediatric population:
Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

Overdose is unlikely.

Unless hypersensitivity to fusidic acid or any of the excipients exist, accidental ingestion of Fusidic acid cream is unlikely to cause any harm. The total quantity of fusidic acid will usually not exceed the approved total daily oral dose of fusidic acid containing products except in children aged less than 1 year and weighing $\leq 10\text{kg}$. Although in this instance a child of this particular age group is unlikely to ingest a whole tube of Fusidic acid cream. The concentration of the excipients is too low to constitute a safety risk.

Fusidic acid cream has no or negligible influence on the ability to drive and use machines.

No interaction studies have been performed.

Fucidic acid 2% is a ready-for-use preparation and should, therefore, not be mixed with other ingredients or preparations.

Store at temperature not exceeding 30°C.

ICPA HEALTH PRODUCTS LIMITED
286/287/288, GIDC, Ankleshwar, Gujarat, India - 393002.
This leaflet was last revised on 31/07/2024

Product Registration Holder:
Healol Pharmaceuticals Sdn Bhd
74-3, Jalan Wangsa Delima 6,
KLSC Wangsa Maju, 53300,
Kuala Lumpur, Malaysia



13001230

Space for text: 125 x 128 mm.