



Avenex® Antiseptic Cream

Active Ingredient: Cetrimide 0.5%w/w, Chlorhexidine Gluconate 0.1%w/w.

Preservative: Chlorocresol 0.2%w/w.

Product Description:

White colour, smooth application cream with characteristic odour.

Indication:

As an antiseptic for skin lesions, cuts, abrasions, infected wounds and burns.

Direction of Use:

Apply gently onto the affected area 2-3 times daily. Wash hand thoroughly before and after each application.

Route of Administration: Topical.

Pharmacodynamics:

Chlorhexidine gluconate is an effective antiseptic with a wide range of activity against micro-organisms, including gram positive and gram negative bacteria, fungi and viruses.

Cetrimide is a quaternary ammonium compound with surfactant and antiseptic properties.

Pharmacokinetics:

Chlorhexidine gluconate and cetrimide are poorly absorbed from the gastro-intestinal tract and skin.

Contraindications:

Patients with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions:

Avenex™ Antiseptic Cream contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalized allergic reactions and anaphylactic shock.

The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. **Avenex™ Antiseptic Cream** should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

For external use only. If symptoms persist, please consult a doctor.

Keep out of reach of children. *Jauhkan daripada kanak-kanak.*

Effects on Ability to Drive and Use Machines: Not known.

Interactions with Other Medicaments:

The cationic nature of the compound results in activity being reduced by soap and anionic compounds as they can form insoluble salts. Anions to be wary of include borates, bicarbonates, carbonates, chlorides, citrates, nitrates, phosphates and sulphates with due attention being paid to the presence of hard water.

Pregnancy & Lactation:

No special precautions need to be taken when used in pregnancy and lactation.

Adverse Effects / Undesirable Effects:

Immune system disorders

Frequency not known: Hypersensitivity including anaphylactic shock.

Symptoms and Treatment of Overdose:**Symptoms**

While accidental ingestion is unlikely to cause any systemic effects due to poor absorption of chlorhexidine gluconate and cetrimide, ingestion of high concentrations may cause esophageal damage and necrosis with symptoms such as nausea and vomiting.

Treatment

Treatment of poisoning is symptomatic; demulcents and diluents may be given if necessary but emesis and lavage should be avoided. Activated charcoal may be considered if the patient presents within an hour of ingestion.

Corticosteroids may reduce oropharyngeal edema.

Storage Condition: Store below 30°C. Protect from direct sunlight.

Shelf Life: Please refer to the packaging. Do not use beyond the expiry date.

Dosage Form: Cream.

Packing Size: 30g.

The information contained in this leaflet is limited. For further information, please consult your doctor or pharmacist.

Product Registration Holder:

Optimist Victory Sdn Bhd
No. 2-3, Jalan Radin Anum,
Bandar Baru Seri Petaling, 57000 Kuala Lumpur.

Manufacturer:

Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.



Date of Revision: September 2025.