

PACKAGE INSERT
Avobic Cream 1%w/w

COMPOSITION

Clotrimazole 1%w/w

PRODUCT DESCRIPTION

White soft cream

PHARMACODYNAMICS

Pharmacotherapeutic group: Antifungals for topical use – imidazole and triazole derivatives

ATC Code: D01A C01

Mechanism of Action

Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the fungal cytoplasmic membrane. Clotrimazole has a broad antimycotic spectrum of action in vitro and in vivo, which includes dermatophytes, yeasts, moulds, etc.

Under appropriate test conditions, the MIC values for these types of fungi are in the region of less than 0.062-8.0 µg/ml substrate. The mode of action of clotrimazole is primarily fungistatic or fungicidal depending on the concentration of clotrimazole at the site of infection. *In vitro* activity is limited to proliferating fungal elements; fungal spores are only slightly sensitive.

In addition to its antimycotic action, clotrimazole also acts on gram-positive microorganisms (Streptococci / Staphylococci / *Gardnerella vaginalis*), and gram-negative microorganisms (*Bacteroides*).

In vitro clotrimazole inhibits the multiplication of *Corynebacteria* and gram-positive cocci - with the exception of *Enterococci* - in concentrations of 0.5-10 µg/ml substrate.

Primarily resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

PHARMACOKINETICS

Pharmacokinetic investigations after dermal application have shown that clotrimazole is minimally absorbed from the intact or inflamed skin into the human blood circulation. The resulting peak serum concentrations of clotrimazole were below the detection limit of 0.001 mcg/ml, suggesting that clotrimazole applied topically is unlikely to lead to measurable systemic effects or side effects.

INDICATION

For the treatment of:

- (i) All dermatomycoses due to moulds and other fungi (e.g. *Trichophyton* species).
- (ii) All dermatomycoses due to yeasts (*Candida* species).
- (iii) Skin diseases showing secondary infection with these fungi.
- (iv) Candidal nappy rash.

ROUTE OF ADMINISTRATION

Topical.

RECOMMENDED DOSAGE

There is no separate dosage schedule for the young or elderly.

The cream should be applied thinly 2-3 times daily and rubbed in gently. A strip of cream (½ cm long) is enough to treat an area of about the size of the hand. Treatment should be continued for at least one month for dermatophyte infections and at least two weeks for candidal infections.

If the feet are infected they should be washed and dried, especially between the toes, before applying the cream.

CONTRAINDICATIONS

Hypersensitivity to the active substance or to any of the excipients.

Do not use the cream to treat nail or scalp infections.

WARNINGS AND PRECAUTIONS

This product contains cetostearyl alcohol, which may cause local skin reactions (e.g. contact dermatitis).

INTERACTIONS WITH OTHER MEDICAMENTS

Not known

PREGNANCY AND LACTATION

Fertility:

No human studies of the effects of clotrimazole on fertility have been performed; however, animal studies have not demonstrated any effects of the drug on fertility.

Pregnancy:

There is a limited amount of data from the use of clotrimazole in pregnant women. Animal studies with clotrimazole have shown reproductive toxicity at high oral doses. At the low systemic exposures of clotrimazole following topical treatment, harmful effects with respect to reproductive toxicity are not predicted. Clotrimazole can be used during pregnancy, but only under the supervision of a physician or midwife.

Lactation

Available pharmacodynamic/ toxicological data in animals have shown excretion of clotrimazole/ metabolites in milk after intravenous administration. A risk to the suckling child cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/ abstain from clotrimazole therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

SIDE EFFECTS

As the listed undesirable effects are based on spontaneous reports, assigning accurate frequency of occurrence for each is not possible.

Immune system disorders: allergic reaction (syncope, hypotension, dyspnea, urticaria).

Skin and subcutaneous tissue disorders: blisters, discomfort/ pain, oedema, erythema, irritation, peeling/ exfoliation, pruritus, rash, stinging/burning.

OVERDOSAGE

No risk of acute intoxication is seen as it is unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favourable to absorption) or inadvertent oral ingestion. There is no specific antidote.

However, in the event of accidental oral ingestion, gastric lavage is rarely required and should be considered only if a life-threatening amount of Clotrimazole has been ingested within the preceding hour or if clinical symptoms of overdose become apparent (e.g. dizziness, nausea or vomiting). Gastric lavage should be carried out only if the airway can be protected adequately.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Clotrimazole cream has no or negligible influence on the ability to drive or use machines.

INSTRUCTION OF USE

Do not accept if seal is broken. Puncture nozzle with piercing point of the cap. Keep the tube tightly closed after use. Keep the medicine out of reach of children. For external use only.

STORAGE CONDITION

Do not store above 30°C.

PACKING

15g collapsible aluminium tubes.

PRODUCT REGISTRATION HOLDER

Abio Marketing Sdn Bhd

No. 2, Jalan SS13/5, 47500 Subang Jaya, Selangor, Malaysia.

MANUFACTURER

S Kant Healthcare Ltd.

1802-1805, G.I.D.C., Phase-III, Vapi- 396 195, Gujarat.

REVISION DATE

September 2024