

Furimed Medicated Shampoo

Furimed Medicated Shampoo appears to be a yellowish to light brown color, opalescent viscous liquid.

Composition

Each ml contains

Chlorhexidine gluconate 20 mg (equivalent to chlorhexidine 11.26 mg)

Miconazole nitrate 20 mg (equivalent to miconazole 17.37 mg)

Pharmacodynamics

Chlorhexidine gluconate

Chlorhexidine gluconate is a Bisbiguanide antimicrobial agent against Gram-positive and Gram-negative bacteria. It is both bactericidal and bacteriostatic depending on the concentration used.

Growth inhibition is achieved by a direct effect on ATP-ase so interfering with the energy transport mechanisms. The bactericidal effect of chlorhexidine results from coagulation of the bacterial cell contents.

Chlorhexidine gluconate is incorporated in the product for its activity against *Staphylococcus intermedius*. *Staphylococcus intermedius* resistance to chlorhexidine has not been reported.

Miconazole nitrate

Miconazole nitrate is an imidazole antifungal agent with activity against yeasts such as *Malassezia pachydermatis*.

It is both fungicidal and fungistatic depending on the concentration used. Miconazole inhibits ergosterol incorporation into cell membranes so increasing concentrations of cytotoxic hydrogen peroxide within the fungal cell wall.

Miconazole nitrate has been incorporated in the product for its activity against *Malassezia pachydermatis*. *Malassezia pachydermatis* resistance to miconazole has not been reported.

Chlorhexidine and miconazole in combination:

In vitro studies have demonstrated synergistic activity against *Microsporum canis* between chlorhexidine and miconazole.

Pharmacokinetics

Chlorhexidine gluconate

High concentrations of chlorhexidine gluconate are achieved in the hair coat and on the skin for the 10 minute period following shampooing. These concentrations are well in excess of the MICs for *Staphylococcus intermedius*. Chlorhexidine gluconate is poorly absorbed from the gastrointestinal tract on ingestion. There is little or no percutaneous absorption. In humans, it has been shown that 26% remains on the skin at 29 hours after application.

Miconazole nitrate

High concentrations of miconazole nitrate are achieved in the hair coat and on the skin for the 10 minute period following shampooing. These concentrations are well in excess of the MICs for *Malassezia pachydermatis*. Miconazole nitrate is poorly absorbed from the skin and gastrointestinal tract.

Indications

Dogs: For the treatment and control of seborrheic dermatitis associated with *Malassezia pachydermatis* and *Staphylococcus intermedius*.

Cats: As an aid in the treatment of ringworm due to *Microsporum canis*.

Dosage and administration

Dogs: Shampoo twice weekly until the symptoms subside and then weekly or when necessary.

Cats: Shampoo twice weekly for a minimum period until coat brushings are negative for the culture of *M. canis*. The maximum length of the treatment period should not exceed 16 weeks.

Wet the animal thoroughly with clean water. Apply shampoo to the animal at several points and massage into the coat especially around the lips, under the tail and between the toes. Make sure that a sufficient amount of the

Cats

Furimed medicated shampoo should only be used in conjunction with griseofulvin when treating for ringworm. Shampooing cats may initially increase the recovery of *M. canis* by brush culture techniques.

Special precautions for use

Official, national and regional antimicrobial policies should be taken into account when the product is used.

i) Special precautions for use in animals

For external use only.

In case of accidental contact with eyes, rinse with plenty of water.

Do not allow the animal to lick itself during shampooing and rinsing, or before it is dried. Take care to avoid the animal inhaling the product or getting it into the nose or mouth during shampooing.

Puppies and kittens should not come into contact with nursing females after treatment until the coat has dried.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

- People with known hypersensitivity to chlorhexidine, miconazole or any of the excipients should avoid contact with the veterinary medicinal product.
- This product can cause eye irritation. Avoid contact with the eyes. In case of accidental contact with eyes, rinse with plenty of water. If irritation persists consult your doctor.
- Avoid excessive handling and stroking of treated animals immediately following treatment.
- Ringworm in the cat is infectious to humans and so it is advisable to wear gloves and have arms covered during clipping and shampooing of infected cats.
- When shampooing a cat, in order to avoid prolonged contact with the shampoo, wash and dry hands gently. Do not scrub.

Interactions with Other Medicaments

None known.

Statement on Usage During Pregnancy and Lactation

Pregnancy: Furimed medicated shampoo in conjunction with griseofulvin should not be used in pregnant cats as griseofulvin is contra-indicated in pregnancy.

Lactation: Please refer warning and precautions.

Side effects

Exceptionally a dog with atopy or a cat with allergic skin disease may develop a pruritic and/or erythematous reaction after treatment.

In very rare circumstances, dogs and cats may develop a skin reaction (itching, redness) after treatment.

Overdose and Treatment

Not applicable.

Withdrawal Period (s)

Not applicable.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/ Jauhkan dari kanak-kanak.

Any unused product or containers should be disposed of in accordance with local and national legislation.

Shelf-life as packaged for sale: 3 years

Shelf life after first opening: 3 months.

Packaging Available

250ml

shampoo is used to raise a lather. Allow the animal to stand for 10 minutes, then rinse off with clean water and leave to dry naturally in a warm, draught-free environment.

Contraindication

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

Warning and Precautions

Special warnings for each target species

Dogs and cats

In order to prevent reoccurrence of the infection, appropriate control methods should be employed in the animal's environment (e.g. cleaning and disinfection of kennels, beds).

Product holder/ Manufacturer

Nam Pharma Sdn. Bhd.
5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Date of revision

13-11-2024