

SOLUZIONE FIRST AID PVP-IODINE 10%w/v SPRAY

Description

Soluzione First Aid PVP-Iodine 10%w/v Spray is reddish brown solution with odor of Iodine.

Composition

Active ingredient: Povidone Iodine 10%w/v.

Non-active ingredient: Purified water, Lauryl alcohol ethoxylate, Citric acid, Trisodium phosphate, Glycerin.

Indications

For minor cuts, scrapes, wounds, burns and skin infections.

Recommended Dosage

Use undiluted. Clean the affected area. Spray a sufficient amount of this product directly to the affected area 2 to 3 times daily or as directed.

Route of Administration

Topical

Contraindication

Soluzione First Aid PVP-Iodine 10%w/v Spray is contraindicated in patients with:

- Hypersensitivity to iodine or to any of the ingredients in the product
- Thyroid disorders - History of abnormal thyroid function or goitre (in particular nodular colloid goitre, endemic goitre and Hashimoto's thyroiditis). Regular use should be avoided in patients on concurrent lithium therapy.

Warning and Precaution

Use of this preparation may interfere with tests of thyroid function. Iodine is absorbed through burns and broken skin and to a lesser extent through intact skin and may lead to toxic levels of iodine in the blood, particularly in patients with renal insufficiency. If used for long periods thyroid function tests should be performed. Particular care should be taken when treating patients with excessive burns. Use of this preparation in patients with renal impairment may possibly result in the development of toxic levels of iodine. Therefore, iodine levels should be monitored in these patients. Particular care should also be taken when treating neonates and very low birth weight infants. Avoid contact with eyes. In case of contact with the eyes, immediately flush the eyes liberally with water for at least 15 minutes and seek medical attention.

Interactions with Other Medicaments

Use of Soluzione First Aid Pvp-Iodine 10%W/V Spray should be avoided when you are receiving radioactive iodine treatment or using products containing mercury such as thiomersal. Other antiseptics or medications can affect how the product works. Avoid preparations containing hydrogen peroxide, silver, taurolidine or octenidine.

Side effects:

The side effects such as skin irritation, discolouration, redness, swelling, small blisters, itching, or dry skin, tiredness, weakness, confusion, rapid shallow breathing, passing little or no urine, fluid retention and swelling of legs, ankle and feet.

Visit your doctor or pharmacist immediately if you experience any side effects as listed below after using:

- Allergic reactions that occur quickly and include an itchy raised skin rash (hives), a body rash, sudden breathing problems, feeling faint, swelling of the mouth, face, lips, tongue or throat – seek medical help straight away
- Symptoms of a thyroid problem may include feeling nervous, moody, difficulty sleeping, hand tremors, fast or irregular heartbeat, trouble breathing, feeling very hot or sweaty, weight loss, weight gain and tiredness.

Symptoms and Treatment of Overdose

Excess iodine can produce goitre and overactive thyroid (hyperthyroidism) or underactive thyroid (hypothyroidism).

In an unlikely event, if you should experience any of the following symptoms: metallic taste in mouth, increased salivation, burning or pain in the throat or mouth, irritation and swelling of the eyes, difficulty in breathing, skin reactions, stomach pain and diarrhoea, any abnormalities due to metabolic disorder, unable to pass urine and kidney failure, immediately consult your doctor.

Therapy: Symptomatic and supportive treatment should be provided with special attention to electrolyte balance and renal and thyroid function.

Pack Size: Available in 15ml, 30ml, 50ml, 60ml, and 100ml.

Product Registration Holder & Manufacturer:

PAHANG PHARMACY SDN. BHD.

Registration No. 197101000448 (10826-K) Lot 5979, Jalan Teratai,

5 ½ Mile Off Jalan Meru, 41050 Klang,

Selangor Malaysia.

Date of Revision:

01/2025

