

THERAPEX ANTI-FUNGAL CLOTRIMAZOLE SPRAY 1% W/V (LEAFLET)
100w x 160h MM

100mm

160mm



ANTI-FUNGAL CLOTRIMAZOLE SPRAY 1% W/V

Composition:

Clotrimazole	1%w/v
Ethanol (Methylated Spirit)	60% v/v

Product Description:

Clear colourless solution with floral odour.

Indication:

- All dermatomycoses due to dermatophytes (Trichophyton species) e.g. athlete's foot (tinea pedis), jock itch (tinea cruris) and ringworm (tinea corporis).
- All dermatomycoses due to yeast (Candida species) e.g. candida vulvitis, candida balanitis
- Dermatomycoses due to moulds and other fungi.
- Skin diseases showing secondary infection with these fungi.

Recommended Dose:

General: The affected area should be washed, thoroughly dried and sprayed a thin layer of Therapex Clotrimazole Spray 1%w/v to the affected area twice daily. Treatment should continue for 2-4 weeks after the disappearance of all signs of infection to ensure mycological cure and prevent relapse.

Adult dosing based on type of infection:

- Candidiasis (topical): twice daily for up to 4 weeks
- Pityriasis versicolor (tinea versicolor): twice daily for up to 4 weeks
- Tinea corporis: twice daily for up to 4 weeks
- Tinea cruris: twice daily for up to 4 weeks
- Tinea pedis: twice daily for up to 4 weeks

Pharmacodynamics:

Clotrimazole is an imidazole derivative with a broad spectrum antimycotic activity.

- **Mechanism of Action**
Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the cytoplasmic membrane.

- **Pharmacodynamic effects**
Clotrimazole has a broad antimycotic spectrum of action in-vitro and in-vivo, which includes dermatophytes, yeasts, moulds, etc.

Under appropriate test conditions, the minimum inhibitory concentration (MIC) values for these types of fungi are in the region of less than 0.062-4(-8) mg/ml substrate. The mode of action of Clotrimazole is primarily fungistatic. In-vitro activity is limited to proliferating fungal elements; fungal spores are slightly sensitive.

In addition to its antimycotic action, Clotrimazole also acts on Trichomonas vaginalis, gram-positive microorganisms (Streptococci/Staphylococci) and gram-negative microorganisms (Bacteroides/Gardnerella vaginalis).

In-vitro Clotrimazole inhibits the multiplication of Corynebacteria and gram-positive cocci-with the exception of Enterococci in concentrations of 0.5-10 mg/ml substrate and exerts a trichomonacidal action at 100 mg/ml.

Primary resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

Pharmacokinetics:

Absorption of Clotrimazole is less than 0.5% after topical application.

Contraindication:

Hypersensitivity to the Clotrimazole or to any of the excipients.

Warning and Precaution:

For irritation or sensitivity caused with the use of Clotrimazole; treatment should be discontinued and appropriate therapy instituted. The patient should be advised to use the medication for the full treatment time even though the symptoms may have improved.

This medicine should be used in the first trimester of pregnancy after consideration of the benefit/risk relation and by physician decision. It is not known whether Clotrimazole is excreted in human being milk. As many drugs are excreted in human being milk, caution should be exercised with Clotrimazole used by nursing mothers.

Keep out of reach of children.
Jauhkan daripada capaian kanak-kanak.

Effects on Ability to Drive and Use Machines:

It is unlikely to impair a patient's ability to drive or use machinery.

Interactions with Other Medicaments:

Not known.

Pregnancy & Lactation:

Weigh the potential benefits of drug treatment against potential risks before prescribing this drug during pregnancy and breastfeeding.

Side Effects:

Local reactions include irritation, burning sensation, contact dermatitis, blister, edema, erythema, peeling of skin, pruritis, urticaria and stinging of skin.

Symptoms of Overdosage and Treatment:

Acute overdosage with topical application of Clotrimazole is unlikely and would not be expected to lead to a life-threatening situation.

Route of Administration:

Topical

Packing Size:

30ml

Storage Condition:

Store below 30°C.

Shelf Life:

Please refer to the packaging.
Do not use beyond the expiry date.

The information contained in this leaflet is limited.

For further information, please consult your doctor or pharmacist.

Product Registration Holder:
Biofizz Marketing Sdn Bhd
No. 30, Jalan Kilang 3,
Kawasan Perindustrian Jelapang,
30100 Ipoh, Perak.

Manufacturer:
Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru,
Johor, Malaysia.

Date of revision: January 2025

