

Front

(L)297mm

pharmaniaga®  
OMEPRAZOLE

OMEPRAZOLE CAPSULE 20 MG

COMPOSITION

Each capsule contains Omeprazole 20 mg.

DESCRIPTION

White to light brown pellets filled in size '2' dark pink/ pink capsule, printed with 'Pharmaniaga logo' and '20' marking.

PHARMACODYNAMICS

Pharmacotherapeutic group: Proton pump inhibitors, ATC code: A02BC01

Mechanism of action

Omeprazole, a racemic mixture of two enantiomers reduces gastric acid secretion through a highly targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. It is rapidly acting and provides control through reversible inhibition of gastric acid secretion with once daily dosing.

Omeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the intracellular canaliculi within the parietal cell, where it inhibits the enzyme H<sup>+</sup> K<sup>+</sup> -ATPase - the acid pump. This effect on the final step of the gastric acid formation process is dose-dependent and provides for highly effective inhibition of both basal acid secretion and stimulated acid secretion, irrespective of stimulus.

Pharmacodynamic effects

All pharmacodynamic effects observed can be explained by the effect of omeprazole on acid secretion.

Effect on gastric acid secretion

Oral dosing with omeprazole once daily provides for rapid and effective inhibition of daytime and night-time gastric acid secretion with maximum effect being achieved within 4 days of treatment. With omeprazole 20 mg, a mean decrease of at least 80% in 24-hour intragastric acidity is then maintained in duodenal ulcer patients, with the mean decrease in peak acid output after pentagastrin stimulation being about 70% 24 hours after dosing.

Oral dosing with omeprazole 20 mg maintains an intragastric pH of ≥ 3 for a mean time of 17 hours of the 24-hour period in duodenal ulcer patients.

As a consequence of reduced acid secretion and intragastric acidity, omeprazole dose-dependently reduces/normalizes acid exposure of the esophagus in patients with gastro-esophageal reflux disease.

The inhibition of acid secretion is related to the area under the plasma concentration-time curve (AUC) of omeprazole and not to the actual plasma concentration at a given time.

No tachyphylaxis has been observed during treatment with omeprazole.

Effect on H. pylori

*H. pylori* is associated with peptic ulcer disease, including duodenal and gastric ulcer disease. *H. pylori* is a major factor in the development of gastritis. *H. pylori* together with gastric acid are major factors in the development of peptic ulcer disease. *H. pylori* is a major factor in the development of atrophic gastritis which is associated with an increased risk of developing gastric cancer.

Eradication of *H. pylori* with omeprazole and antimicrobials is associated with, high rates of healing and long-term remission of peptic ulcers.

Dual therapies have been tested and found to be less effective than triple therapies. They could, however, be considered in cases where known hypersensitivity precludes use of any triple combination.

Other effects related to acid inhibition

During treatment with antsecretory medicinal products, serum gastrin increases in response to the decreased acid secretion. Also, chromogranin A (CgA) increases due to decreased gastric acidity. The increased CgA level may interfere with investigations for neuroendocrine tumours. Available published evidence suggests that proton pump inhibitors should be discontinued between 5 days and 2 weeks prior to CgA measurements. This is to allow CgA levels that might be spuriously elevated following PPI treatment to return to reference range.

An increased number of ECL cells possibly related to the increased serum gastrin levels, have been observed in some patients (both children and adults) during long term treatment with omeprazole. The findings are considered to be of no clinical significance.

During long-term treatment gastric glandular cysts have been reported in a somewhat increased frequency. These changes are a physiological consequence of pronounced inhibition of acid secretion, are benign and appear to be reversible.

Decreased gastric acidity due to any means including proton pump inhibitors, increases gastric counts of bacteria normally present in the gastrointestinal tract. Treatment with acid-reducing drugs may lead to slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* and, in hospitalized patients, possibly also *Clostridium difficile*.

PHARMACOKINETICS

Absorption

Omeprazole and omeprazole magnesium are acid labile and are therefore administered orally as enteric-coated granules in capsules or tablets. Absorption of omeprazole is rapid, with peak plasma levels occurring approximately 1-2 hours after dose. Absorption of omeprazole takes place in the small intestine and is usually completed within 3-6 hours. Concomitant intake of food has no influence on the bioavailability. The systemic availability (bioavailability) from a single oral dose of omeprazole is approximately 40%. After repeated once-daily administration, the bioavailability increases to about 60%.

Distribution

The apparent volume of distribution in healthy subjects is approximately 0.3 l/kg body weight. Omeprazole is 97% plasma protein bound.

Metabolism

Omeprazole is completely metabolised by the cytochrome P450 system (CYP). The major part of its metabolism is dependent on the polymorphically expressed CYP2C19, responsible for the formation of hydroxyomeprazole, the major metabolite in plasma. The remaining part is dependent on another specific isoform, CYP3A4, responsible for the formation of omeprazole sulfone. As a consequence of high affinity of omeprazole to CYP2C19, there is a potential for competitive inhibition and metabolic drug-drug interactions with other substrates for CYP2C19. However, due to low affinity to CYP3A4, omeprazole has no potential to inhibit the metabolism of other CYP3A4 substrates. In addition, omeprazole lacks an inhibitory effect on the main CYP enzymes.

Approximately 3% of the Caucasian population and 15-20% of Asian populations lack a functional CYP2C19 enzyme and are called poor metabolisers. In such individuals the metabolism of omeprazole is probably mainly catalyzed by CYP3A4. After repeated once-daily administration of 20 mg omeprazole, the mean AUC was 5 to 10 times higher in poor metabolisers than in subjects having a functional CYP2C19 enzyme (extensive metabolisers). Mean peak plasma concentrations were also higher, by 3 to 5 times. These findings have no implications for the posology of omeprazole.

Excretion

The plasma elimination half-life of omeprazole is usually shorter than one hour both after single and repeated oral once-daily dosing. Omeprazole is completely eliminated from plasma between doses with no tendency for accumulation during once-daily administration. Almost 80% of an oral dose of omeprazole is excreted as metabolites in the urine, the remainder in the faeces, primarily originating from bile secretion.

The AUC of omeprazole increases with repeated administration. This increase is dose-dependent and results in a non-linear dose-AUC relationship after repeated administration. This time- and dose-dependency is due to a decrease of first pass metabolism and systemic clearance probably caused by an inhibition of the CYP2C19 enzyme by omeprazole and/or its metabolites (e.g. the sulfone).

No metabolite has been found to have any effect on gastric acid secretion.

Special populations

Hepatic impairment

The metabolism of omeprazole in patients with liver dysfunction is impaired, resulting in an increased AUC. Omeprazole has not shown any tendency to accumulate with once-daily dosing.

Renal impairment

The pharmacokinetics of omeprazole, including systemic bioavailability and elimination rate, are unchanged in patients with reduced renal function.

Elderly

The metabolism rate of omeprazole is somewhat reduced in elderly subjects (75-79 years of age).

INDICATIONS

Pharmaniaga Omeprazole Capsule 20mg are indicated for treatment of:

- Duodenal ulcer
- Gastric ulcer
- NSAID associated gastric and duodenal ulcers or erosions
- *Helicobacter pylori* eradication in peptic ulcer disease
- Reflux oesophagitis
- Symptomatic gastro-oesophageal reflux disease
- Acid related dyspepsia
- Zollinger-Ellison syndrome

CONTRAINDICATIONS

Hypersensitivity to omeprazole, substituted benzimidazoles or to any of the excipients.

Omeprazole like other proton pump inhibitors must not be used concomitantly with nelfinavir (see section Drug Interactions).

ADVERSE REACTIONS

The most common side effects (1-10% of patients) are headache, abdominal pain, constipation, diarrhoea, flatulence and nausea/vomiting.

The following adverse drug reactions have been identified or suspected in the clinical trials programme for omeprazole and post-marketing. None was found to be dose-related. Adverse reactions listed below are classified according to frequency and System Organ Class (SOC). Frequency categories are defined according to the following convention: common ≥1/100 to <1/10; uncommon ≥1/1,000 to <1/100; rare ≥ 1/10,000 to <1/1,000; very rare <1/10,000.

| SOC/frequency                                               | Adverse reaction                                                                                                                                                                                                           |
|-------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Blood and lymphatic system disorders</b>                 |                                                                                                                                                                                                                            |
| Rare                                                        | Leukopenia, thrombocytopenia, agranulocytosis, pancytopenia                                                                                                                                                                |
| <b>Immune system disorders</b>                              |                                                                                                                                                                                                                            |
| Rare                                                        | Hypersensitivity reactions e.g. fever, angioedema and anaphylactic reaction/shock                                                                                                                                          |
| <b>Metabolism and nutrition disorders</b>                   |                                                                                                                                                                                                                            |
| Rare                                                        | Hyponatraemia                                                                                                                                                                                                              |
| Very rare                                                   | Hypomagnesaemia; severe hypomagnesaemia may result in hypocalcaemia. Hypomagnesaemia may also result in hypokalaemia.                                                                                                      |
| <b>Psychiatric disorders</b>                                |                                                                                                                                                                                                                            |
| Uncommon                                                    | Insomnia                                                                                                                                                                                                                   |
| Rare                                                        | Agitation, confusion, depression, aggression, hallucinations                                                                                                                                                               |
| <b>Nervous system disorders</b>                             |                                                                                                                                                                                                                            |
| Common                                                      | Headache                                                                                                                                                                                                                   |
| Uncommon                                                    | Dizziness, paraesthesia, somnolence                                                                                                                                                                                        |
| Rare                                                        | Taste disturbance                                                                                                                                                                                                          |
| <b>Eye disorders</b>                                        |                                                                                                                                                                                                                            |
| Rare                                                        | Blurred vision                                                                                                                                                                                                             |
| <b>Ear and labyrinth disorders</b>                          |                                                                                                                                                                                                                            |
| Uncommon                                                    | Vertigo                                                                                                                                                                                                                    |
| <b>Respiratory, thoracic and mediastinal disorders</b>      |                                                                                                                                                                                                                            |
| Rare                                                        | Bronchospasm                                                                                                                                                                                                               |
| <b>Gastrointestinal disorders</b>                           |                                                                                                                                                                                                                            |
| Common                                                      | Abdominal pain, constipation, diarrhoea, flatulence, nausea/vomiting, fundic gland polyps (benign)                                                                                                                         |
| Rare                                                        | Dry mouth, stomatitis, gastrointestinal candidiasis, microscopic colitis                                                                                                                                                   |
| <b>Hepatobiliary disorders</b>                              |                                                                                                                                                                                                                            |
| Uncommon                                                    | Increased liver enzymes                                                                                                                                                                                                    |
| Rare                                                        | Hepatitis with or without jaundice, hepatic failure, encephalopathy in patients with pre-existing liver disease                                                                                                            |
| <b>Skin and subcutaneous tissue disorders</b>               |                                                                                                                                                                                                                            |
| Uncommon                                                    | Dermatitis, pruritus, rash, urticaria                                                                                                                                                                                      |
| Rare                                                        | Alopecia, photosensitivity, erythema multiforme, Stevens-John son syndrome, toxic epidermal necrolysis (TEN), acute generalized exanthematous pustulosis (AGEP), drug rash with eosinophilia and systemic symptoms (DRESS) |
| Not known                                                   | Subacute cutaneous lupus erythematosus (SCLE)                                                                                                                                                                              |
| <b>Musculoskeletal and connective tissue disorders</b>      |                                                                                                                                                                                                                            |
| Uncommon                                                    | Fracture of the hip, wrist or spine                                                                                                                                                                                        |
| Rare                                                        | Arthralgia, myalgia, muscular weakness                                                                                                                                                                                     |
| <b>Renal and urinary disorders</b>                          |                                                                                                                                                                                                                            |
| Rare                                                        | Interstitial nephritis                                                                                                                                                                                                     |
| <b>Reproductive system and breast disorders</b>             |                                                                                                                                                                                                                            |
| Very rare                                                   | Gynaecomastia                                                                                                                                                                                                              |
| <b>General disorders and administration site conditions</b> |                                                                                                                                                                                                                            |
| Uncommon                                                    | Malaise                                                                                                                                                                                                                    |
| Rare                                                        | Increased sweating, peripheral oedema                                                                                                                                                                                      |

Other possible side effects include *Clostridium difficile* associated diarrhoea and Vitamin B<sub>12</sub> deficiency.

WARNINGS AND PRECAUTIONS

In the presence of any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, dysphagia, haematemesis or melena) and when gastric ulcer is suspected or present, malignancy should be excluded, as treatment may alleviate symptoms and delay diagnosis.

Co-administration of atazanavir with proton pump inhibitors is not recommended. If the combination of atazanavir with a proton pump inhibitor is judged unavoidable, dose clinical monitoring (e.g virus load) is recommended in combination with an increase in the dose of atazanavir to 400 mg with 100 mg of rilonavir; omeprazole 20 mg should not be exceeded.

Omeprazole, as all acid-blocking medicines, may reduce the absorption of vitamin B<sub>12</sub> (cyanocobalamin) due to hypo- or achlorhydria. This should be considered in patients with reduced body stores or risk factors for reduced vitamin B<sub>12</sub> absorption on long-term therapy.

Omeprazole is a CYP2C19 inhibitor. When starting or ending treatment with omeprazole, the potential for interactions with drugs metabolised through CYP2C19 should be considered. An interaction is observed between clopidogrel and omeprazole. The published clinical relevance of this interaction is uncertain. As a precaution, concomitant use of omeprazole and clopidogrel should be discouraged.

Pharmaniaga Omeprazole Capsule 20mg contain sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

Treatment with proton pump inhibitors may lead to slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* and, in hospitalised patients, possibly also *Clostridium difficile*.

Some published observational studies suggest that proton pump inhibitor (PPI) therapy may be associated with a small increased risk for osteoporosis related fractures. However, in other similar observational studies no such increased risk was found.

In randomized, double-blind and controlled published clinical studies on omeprazole and esomeprazole (including two open long-term studies of up to more than 12 years) there are no indications that PPIs are associated with osteoprototic fractures.

As in all long-term treatments, especially when exceeding a treatment period of 1 year, patients should be kept under regular surveillance.

Subacute Cutaneous Lupus Erythematosus (SCLE)

Proton pump inhibitors are associated with very infrequent cases of subacute cutaneous lupus erythematosus (SCLE). If lesions occur, especially in sun exposed areas of the skin, and if accompanied by arthralgia, the patient should seek medical help promptly and the health care professional should consider stopping esomeprazole. SCLE after previous treatment with a proton pump inhibitor may increase the risk of SCLE with other proton pump inhibitors.

Hypomagnesaemia

Severe hypomagnesaemia has been reported in patients treated with PPI like esomeprazole for at least three months, and in most cases for a year. Serious manifestations of hypomagnesaemia such as fatigue, tetany, delirium, convulsions, dizziness and ventricular arrhythmia can occur but they may begin insidiously and be overlooked. In most affected patients, hypomagnesaemia improved after magnesium replacement and discontinuation of the PPI.

For patients expected to be on prolonged treatment or who take PPI with digoxin or drugs that may cause hypomagnesaemia (e.g., diuretics), health care professionals should consider measuring magnesium levels before starting PPI treatment and periodically during treatment.

Fracture

Proton pump inhibitors, especially if used in high doses and over long durations (>1 year), may modestly increase the risk of hip, wrist and spine fracture, predominantly in the elderly or in presence of other recognised risk factors. Observational studies suggest that proton pump inhibitors may increase the overall risk of fracture by 10 -40%. Some of this increase may be due to other risk factors. Patients at risk of osteoporosis should receive care according to current clinical guidelines and they should have an adequate intake of vitamin D and calcium.

Interference with laboratory tests

Increased Chromogranin A (CgA) level may interfere with investigations for neuroendocrine tumours. To avoid this interference, Omeprazole treatment should be stopped for at least 5 days before CgA measurements. If CgA and gastrin levels have not returned to reference range after initial measurement, measurement should be repeated 14 days after cessation of proton pump inhibitor treatment.

Effects on Ability to Drive and Use Machines

Omeprazole is not likely to affect the ability to drive or use machines. Adverse drug reactions such as dizziness and visual disturbances may occur (see Adverse Reactions). If affected, patients should not drive or operate machinery.

Pregnancy and lactation

Results from three published prospective epidemiological studies (more than 1000 exposed outcomes) indicate no adverse effects of omeprazole on pregnancy or on the health of the foetus/newborn child. Omeprazole can be used during pregnancy.

Omeprazole is excreted in breast milk but it not likely to influence the child when therapeutic doses are used.

INTERACTIONS WITH OTHER MEDICAMENTS

Effects of omeprazole on the pharmacokinetics of other drugs

**Absorption:** The gastric acid suppression during treatment with omeprazole and other PPIs might decrease or increase the absorption of drugs with a gastric pH dependent absorption. Like with other drugs that decrease the intragastric acidity, the absorption of drugs, such ketoconazole, itraconazole and erlotinib can decrease while the absorption of drugs such as digoxin can increase during treatment with omeprazole. Concomitant treatment with omeprazole (20mg daily) and digoxin in healthy subjects increased the bioavailability of digoxin by 10% (up to 30% in two out of ten subjects).

**Metabolism:** Omeprazole inhibits CYP2C19, the major omeprazole metabolizing enzyme. Thus, the metabolism of concomitant drugs also metabolized by CYP2C19, such as diazepam, phenytoin, warfarin (R-warfarin) or other vitamin K antagonists and cilostazol, may be delayed. Monitoring of patients receiving phenytoin is recommended and a reduction of the phenytoin dose may be necessary. However, concomitant treatment with Pharmaniaga Omeprazole Capsule 20mg daily did not change the blood concentration of phenytoin in patients on continuous treatment with this drug. In patient receiving of the warfarin (or other vitamin K antagonist) dose may be necessary. Concomitant treatment with Pharmaniaga Omeprazole Capsule 20mg daily did, however, not change coagulation time in patients on continuous treatment with warfarin. Omeprazole, given in doses of 40mg to healthy subjects in a cross-over study, increased C<sub>max</sub> and AUC for cilostazol by 18% and 26% respectively, and one of its active metabolites by 29% and 69% respectively.

Result of studies in healthy subjects have shown a pharmacokinetic/pharmacodynamic interaction between clopidogrel (300mg loading dose/75mg daily maintenance dose) and omeprazole (80mg p.o daily, i.e., four times the recommended dose) resulting in decreased exposure to the active metabolite of clopidogrel by an average of 46% and resulting in decreased maximum inhibition of (ADP induced) platelet aggregation by an average of 16%.

|                                                                                                               |  |                                                  |  |                                            |  |
|---------------------------------------------------------------------------------------------------------------|--|--------------------------------------------------|--|--------------------------------------------|--|
| attn                                                                                                          |  | customer <b>Pharmaniaga Manufacturing Berhad</b> |  | date <b>22.01.2025</b>                     |  |
| <b>1</b> colours                                                                                              |  | <b>Black</b>                                     |  | <b>Please Chop &amp; Sign For Approval</b> |  |
| size (L)297 x (W)210mm                                                                                        |  | material 60g Simili                              |  |                                            |  |
| description <b>Omeprazole 20mg Insert - Front</b>                                                             |  | finishing                                        |  |                                            |  |
|  <b>FocusPrint</b> SDN BHD |  | t: 603-8766 6030 f: 603-8766 6033 (Factory)      |  |                                            |  |
| website: www.focusprint.com.my                                                                                |  |                                                  |  |                                            |  |
| artwork prepared by: cynthia gap                                                                              |  | email: graphic@focusprint.info (Graphic Dept)    |  | <b>date:</b>                               |  |

| ARTWORK LOG |            |                                           |
|-------------|------------|-------------------------------------------|
| Version no. | Date       | Reason for Change                         |
| 01          | 28.03.2022 | - New artwork received and amend wording. |
| 02          | 16.01.2023 | - Amend wording and date of revision.     |
| 03          | 18.01.2023 | - Relocate grey box.                      |
| 04          | 10.05.2023 | - Amend wording and date of revision.     |
| 05          | 16.05.2023 | - Add new paragraph.                      |
| 06          | 17.05.2023 | - Amend wording.                          |
| 07          | 25.05.2023 | - Amend wording and date of revision.     |
| 08          | 24.11.2023 | - Amend wording.                          |
| 09          | 27.11.2023 | - Amend wording.                          |
| 10          | 27.12.2024 | - Amend wording.                          |
| 11          | 07.01.2025 | - Amend wording.                          |
| 12          | 20.01.2025 | - Amend wording.                          |
| 13          | 22.01.2025 | - Amend wording.                          |

**IMPORTANT ! Please note that this colour print is for visual purpose only. Output colour might differ in actual production.**

Back

(L)297mm

(W)210mm

It is, however, uncertain to what extent this interaction is clinically important. One prospective, randomized (but incomplete) study (in over 3760 patients comparing placebo with omeprazole 20 mg in patients treated with clopidogrel and ASA) and non-randomized, post-hoc analyses of data from large, prospective, randomized clinical outcome studies (in over 47000 patients) did not show any evidence of an increased risk for adverse cardiovascular outcome when clopidogrel and PPIs, including omeprazole, were given concomitantly.

Results from a number of observational studies are inconsistent with regard to increased risk or no increased risk for CV thromboembolic events when clopidogrel is given together with a PPI.

When clopidogrel was given together with a fixed dose combination of esomeprazole 20 mg + ASA 81 mg compared to clopidogrel alone in a study in healthy subjects there was a decreased exposure by almost 40% of the active metabolite of clopidogrel. However, the maximum levels of inhibition of (ADP induced) platelet aggregation in these subjects were the same in the clopidogrel and the clopidogrel + the combined (esomeprazole + ASA) product groups, likely due to the concomitant administration of low dose ASA.

Omeprazole is partly metabolised also by CYP3A4, but omeprazole does not inhibit this enzyme. Thus, omeprazole does not affect the metabolism of drugs metabolised by CYP3A4, such as cyclosporin, lidocaine, quinidine, estradiol, erythromycin, and budesonide. Results from a range of interaction studies with omeprazole versus other drugs demonstrate that omeprazole, 20-40 mg daily, has no significant influence on any other CYP enzymes relevant for drug metabolism, as shown by the lack of metabolic interaction with substrates for CYP1A2 (such as caffeine, theophylline), CYP2C9 (such as S-warfarin, piroxicam, diclofenac, naproxen), CYP2D6 (such as metoprolol, propranolol), CYP2E1 (such as ethanol).

Unknown mechanism: Concomitant administration of omeprazole has been reported to increase the serum levels of tacrolimus.

When given together with proton pump inhibitors, methotrexate levels have been reported to increase in some patients. In high-dose methotrexate administration a temporary withdrawal of omeprazole may need to be considered.

Omeprazole has been reported to interact with some antiretroviral drugs. The clinical importance and the mechanisms behind these interactions are not always known. Increased gastric pH during omeprazole treatment may change the absorption of the antiretroviral drug. Other possible interaction mechanisms are via CYP 2C19. For some antiretroviral drugs, such as atazanavir and nelfinavir, decreased serum levels have been reported when given together with omeprazole. Concomitant administration with omeprazole and drugs such as atazanavir and nelfinavir is therefore not recommended. For other antiretroviral drugs, such as saquinavir, elevated serum levels have been reported. There are also some antiretroviral drugs of which unchanged serum levels have been reported when given with omeprazole.

Effects of other drugs on the pharmacokinetics of omeprazole

Metabolism: Since omeprazole is metabolised by CYP2C19 and CYP3A4, drugs known to inhibit CYP 2C19 or CYP 3A4 or both (such as clarithromycin and voriconazole) may lead to increased omeprazole serum levels by decreasing the rate of omeprazole's metabolism.

Concomitant voriconazole treatment resulted in more than doubling of the omeprazole exposure. Since high doses of omeprazole have been well-tolerated, adjustment of the omeprazole dose is not required during temporary concomitant use. Drugs known to induce CYP 2C19 or CYP 3A4 or both (such as rifampicin and St John's wort) may lead to decreased omeprazole serum levels by increasing omeprazole's rate of metabolism.

RECOMMENDED DOSAGE

Posology and method of administration

The tablets must be swallowed whole with at least 1/2 glass of liquid. They must not be chewed or crushed. The tablets may also be dispersed in water or any acidic drink (e.g. juice). The dispersion must then be taken within 30 minutes.

Duodenal ulcer

The recommended dosage in patients with an active duodenal ulcer is Pharmaniaga Omeprazole Capsule 20mg once daily. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 2 week treatment period.

In patients with poorly responsive duodenal ulcer Pharmaniaga Omeprazole Capsule 20mg, 40 mg once daily is recommended and healing is usually achieved within 4 weeks.

For the prevention of relapse in patients with duodenal ulcer disease the recommended dose is Pharmaniaga Omeprazole Capsule 20mg, 20mg once daily. If needed the dose can be increased to Pharmaniaga Omeprazole Capsule 20mg, 20-40mg once daily.

For NSAID associated duodenal ulcers see NSAID associated gastroduodenal lesions.

For eradication of *Helicobacter pylori* see *Helicobacter pylori* (Hp) eradication regimens in peptic ulcer disease.

Gastric ulcer

The recommended dosage is Pharmaniaga Omeprazole Capsule 20mg, 20mg once daily. Symptom resolution is rapid and in most patients healing occurs within 4 weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 4 weeks' treatment period.

In patients with poorly responsive gastric ulcer Pharmaniaga Omeprazole Capsule 20mg, 40mg once a daily is recommended and healing is usually achieved within 8 weeks.

For the prevention of relapse in patients with poorly responsive gastric ulcer the recommended dose is Pharmaniaga Omeprazole Capsule 20mg, 20mg once daily. If needed the dose can be increased to Pharmaniaga Omeprazole Capsule 20mg, 40mg once daily.

For eradication of *Helicobacter pylori* see *Helicobacter pylori* (Hp) eradication regimens in peptic ulcer disease.

NSAID associated ulcers or gastroduodenal erosions

NSAID associated gastric ulcers, duodenals ulcers or gastroduodenal erosions in patients with or without continues NSAID treatment the recommended dosage of Pharmaniaga Omeprazole Capsule is 20 mg once daily. Symptom resolution is rapid and in most patients healing occurs within 4 weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 4 weeks treatment period.

For the prevention of NSAID associated gastric ulcers, duodenal ulcers, gastroduodenal erosions and dyspeptic symptoms the recommended dosage of Pharmaniaga Omeprazole Capsule is 20mg once daily.

*Helicobacter pylori* (Hp) eradication regimens in peptic ulcer disease

Triple therapy regimens:

Pharmaniaga Omeprazole Capsule 20mg, amoxicillin 1g and clarithromycin 500mg, all twice a day for one week  
or  
Pharmaniaga Omeprazole Capsule 20mg, metronidazole 400mg (or tinidazole 500mg) and clarithromycin 250mg, all twice a day for one week  
or  
Pharmaniaga Omeprazole Capsule 20mg, 40mg once daily with amoxicillin 500mg and metronidazole 400mg both three times a day for one week.

Dual therapy regimens:

Pharmaniaga Omeprazole Capsule 20mg, 40mg - 80mg daily with amoxicillin 1.5g daily in divided doses for two weeks. In published clinical studies daily doses of 1.5g-3g of amoxicillin have been used  
or  
Pharmaniaga Omeprazole Capsule 20mg, 40mg once daily and clarithromycin 500mg three times a day for two weeks.

To ensure healing in patients with active peptic ulcer disease, see further dosage recommendations for Duodenal and Gastric ulcer.

In each regimen if the patient is still Hp positive, therapy may be repeated.

Reflux oesophagitis

The recommended dosage is Pharmaniaga Omeprazole Capsule 20mg, once daily. Symptom resolution is rapid and in most patients healing occurs within 4 weeks. For those patients who may not be fully healed after the initial course, healing usually occurs during a further 4 weeks' treatment period.

In patients with severe reflux oesophagitis Pharmaniaga Omeprazole Capsule 20mg, 40mg once daily is recommended and healing is usually achieved within 8 weeks.

For the long-term management of patients with healed reflux oesophagitis, the recommended dose is Pharmaniaga Omeprazole Capsule 20mg once daily. If needed, the dose can be increased to 40mg once daily.

Symptomatic gastro-oesophageal reflux disease

The recommended dosage is Pharmaniaga Omeprazole Capsule 20mg, 20mg daily. Symptom relief is rapid. Patients may respond adequately to 20mg daily, and therefore individual dose adjustment should be considered.

If symptom control has not been achieved after 4 weeks treatment with Pharmaniaga Omeprazole Capsule 20mg, 20mg daily, further investigation is recommended.

Acid related dyspepsia

In the relief of symptoms in patients with epigastric pain/discomfort with or without heartburn the recommended dosage is Pharmaniaga Omeprazole Capsule 20mg, 20mg once daily.

If symptom control has not been achieved after 4 weeks treatment with Pharmaniaga Omeprazole Capsule 20mg, daily, further investigation is recommended.

Zollinger-Ellison syndrome

In patients with Zollinger-Ellison syndrome the dosage should be individually adjusted and treatment continued as long as is clinically indicated. The recommended initial dosage is Pharmaniaga Omeprazole Capsule 20mg, 60mg daily. All patients with severe disease and inadequate response to other therapies have been effectively controlled and more than 90% of the patients maintained on doses of omeprazole 20-120mg daily. When doses exceed Pharmaniaga Omeprazole Capsule 20mg, 80mg daily, the dose should be divided and given twice daily.

Impaired renal function

Dose adjustment is not needed in patients with impaired renal function.

Impaired hepatic function

As bioavailability and plasma half-life of omeprazole are increased in patients with impaired hepatic function a daily dose of 20mg may be sufficient.

Elderly

Dose adjustment is not needed in the elderly.

Children

Pharmaniaga Omeprazole Capsule 20mg, is not approved for use in children.

ROUTE OF ADMINISTRATION

Oral

OVERDOSE AND TREATMENT

There is limited information available on the effects of overdoses of omeprazole in humans. Nausea, vomiting, dizziness, abdominal pain, diarrhea and headache have been reported. Also, apathy depression and confusion have been described in single cases.

The symptoms described in connection to omeprazole overdose have been transient, and no serious outcome has been reported. The rate of elimination was unchanged (first order kinetics) with increased doses. Treatment, if needed, is symptomatic.

STORAGE CONDITION

Store below 30°C.  
Protect from moisture and light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

DOSAGE FORMS AND PACKAGING AVAILABLE

7 capsules per Alu/Alu blister.  
In a box of 2 blisters consist of 14 capsules.

Date of Revision

22nd January 2025

PRODUCT REGISTRATION HOLDER/MANUFACTURER:

PHARMANIAGA MANUFACTURING BERHAD (199001006232)

11A, Jalan P/1, Kawasan Perusahaan Bangi,  
43650 Bandar Baru Bangi, Selangor, Malaysia

|                                                                                                               |  |                                                                               |  |                                            |  |
|---------------------------------------------------------------------------------------------------------------|--|-------------------------------------------------------------------------------|--|--------------------------------------------|--|
| attn                                                                                                          |  | customer <b>Pharmaniaga Manufacturing Berhad</b>                              |  | date <b>22.01.2025</b>                     |  |
| <b>1</b> colours                                                                                              |  | <b>Black</b>                                                                  |  | <b>Please Chop &amp; Sign For Approval</b> |  |
| size (L) <b>297</b> x (W) <b>210mm</b>                                                                        |  | material <b>60g Simili</b>                                                    |  |                                            |  |
| description <b>Omeprazole 20mg Insert - Back</b>                                                              |  | finishing                                                                     |  |                                            |  |
|  <b>FocusPrint</b> SDN BHD |  | t: 603-8766 6030 f: 603-8766 6033 (Factory)<br>website: www.focusprint.com.my |  |                                            |  |
| artwork prepared by: <b>cynthia yap</b>                                                                       |  | email: <b>graphic@focusprint.info</b> (Graphic Dept)                          |  | <b>date:</b>                               |  |

**IMPORTANT ! Please note that this colour print is for visual purpose only. Output colour might differ in actual production.**

| ARTWORK LOG |            |                                           |
|-------------|------------|-------------------------------------------|
| Version no. | Date       | Reason for Change                         |
| 01          | 28.03.2022 | - New artwork received and amend wording. |
| 02          | 16.01.2023 | - Amend wording and date of revision.     |
| 03          | 18.01.2023 | - Relocate grey box.                      |
| 04          | 10.05.2023 | - Amend wording and date of revision.     |
| 05          | 16.05.2023 | - Add new paragraph.                      |
| 06          | 17.05.2023 | - Amend wording.                          |
| 07          | 25.05.2023 | - Amend wording and date of revision.     |
| 08          | 24.11.2023 | - Amend wording.                          |
| 09          | 27.11.2023 | - Amend wording.                          |
| 10          | 27.12.2024 | - Amend wording.                          |
| 11          | 07.01.2025 | - Amend wording.                          |
| 12          | 20.01.2025 | - Amend wording.                          |
| 13          | 22.01.2025 | - Amend wording.                          |