

Remeco

PEPTICON

Liquid
Oral Suspension

COMPOSITION
Each 10ml of suspension contains 500mg sodium alginate, 267mg sodium bicarbonate and 160mg calcium carbonate.

PRODUCT DESCRIPTION
Remeco PEPTICON Liquid Oral Suspension is an opaque, off-white colored, peppermint odour, viscous suspension.

PHARMACODYNAMICS
Remeco PEPTICON Liquid Oral Suspension relieves the pain & discomfort of heartburn and acid indigestion by forming a protective layer on top of the stomach contents to prevent stomach acid escaping up into the food pipe where it hurts.

PHARMACOKINETICS
Absorption:
The mechanism of action of the medicinal product is physical and does not depend on absorption into the systemic circulation.

Distribution:
No data available.

Biotransformation:
No data available.

Elimination:
No data available.

Linear /Non-linear situation:
No data available.

INDICATION
Treatment of symptoms resulting from the reflux of acid into the oesophagus such as acid regurgitation, heartburn and indigestion which, for example, can occur after meals or during pregnancy.

RECOMMENDED DOSAGE AND ADMINISTRATION
For Oral Use.

Shake well before use.

Adults and children over 12 years: Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime, up to four times a day.

Children under 12 years: Not to be given unless recommended by your doctor.

CONTRAINDICATIONS
Remeco PEPTICON Liquid Oral Suspension should not be used in patients with known or suspected hypersensitivity to any of the active ingredients or other excipients (parabens) contained in the suspension.

WARNINGS AND PRECAUTIONS
Each 10 ml dose has a sodium content of 149.45 mg. This should be taken into account when a highly restricted salt diet is recommended, e.g. in some cases of congestive cardiac failure and renal impairment.

Each 10 ml contains 64.06 mg calcium. Care needs to be taken in treating patients with hypercalcaemia, nephrocalcinosis and recurrent calcium renal calculi.

This medicinal product contains methyl paraben (E 218) and propyl paraben (E 216), which may cause allergic reactions (possibly delayed).

If symptoms do not improve after seven days, the clinical situation of patient should be reviewed by doctor. Avoid to use more than you should.

Like as other antacids, antacid-alginate combinations can mask more serious underlying situations.

Do not use PEPTICON in following situations:
• Patients with severe / partial renal function / failure
• Hypophosphatemic patients

It is possible for this medicine to reduction in activity for the patient who has very low gastric acid levels.

Children with renal failure or gastroenteritis are under a serious hypernatrem-ia risk

Except medical advice, it is generally not recommended in the treatment of children under 12 years.

INTERACTIONS WITH OTHER MEDICAMENTS
It is recommended to wait 2 hours before using other medicinal products (especially H2 antihistamines, tetracyclines, digoxin, fluoroquinolone, iron salts, ketoconazole, neuroleptics, penicillamine, beta blockers (atenolol, metoprolol, propranolol), glucocorticoid, chloroquine, bisphosphonates and estramustine (for prostate gland cancer)) since the calcium carbonate in the product has antacid properties.

Since the absorption of medicine containing levothyroxine is impaired when taken together with PEPTICON, it is recommended to wait minimum 2 hours between taking these products.

Additional information on special populations:
No interaction studies have been conducted.

Paediatric population:
No interaction studies have been conducted.

PREGNANCY AND LACTATION
Women of childbearing potential / birth control (contraception)
No data is available that proposes additional contraceptive method while Remeco PEPTICON Liquid Oral Suspension treatment. There is no evidence that the effects on contraception methods.

Pregnancy
More than 500 pregnant women have shown that the active substances have no effect on the fetus / newborn poisoning or deformity. When clinically necessary, Remeco PEPTICON Liquid Oral Suspension can be used during pregnancy.

Breast feeding
These active substances had no negative effects on the infants / newborns of the mothers receiving treatment and breastfeeding. Remeco PEPTICON Liquid Oral Suspension can be used during breastfeeding.

Fertility
Alginate has no negative effect on reproductive ability and fertility. Remeco PEPTICON Liquid Oral Suspension has no adverse effect on human fertility.

SIDE EFFECTS
Adverse reactions have been ranked under headings of frequency using the following convention: very common (≥ 1/10), common (<1/10 and ≥1/100), uncommon (<1/100 and ≥1/1000), rare (<1/1000 and ≥1/10,000), very rare (<1/10,000) and not known (cannot be estimated from the available data).

Immune System Disorders
Very rare: Allergic symptoms such as urticaria, anaphylactic or anaphylactoid reactions may develop very rarely in patients sensitive to the ingredients of the product.

Metabolism and Nutrition Disorders
Not known: Excess amount of calcium carbonate consumption can cause alkalosis, hypercalcemia, acid rebound, milk alkali syndrome or constipation. Generally, these side effects can occur when taken more than normal dose.

Respiratory, Thoracic and Mediastinal Disorders
Very rare: Respiratory effects such as bronchospasm

Gastrointestinal Disorders
Not Known: Constipation

OVERDOSAGE
In the event of overdose, symptomatic treatment should be given. The patient may notice abdominal distension.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE
It has no negative effect on ability to drive and machine operation.

STORAGE CONDITION
Store below 30°C. Do not refrigerate or freeze. Check that the cap seal is unbroken before first taking this product.

PACKING
100ml or 200ml of amber colour glass bottle.

PRODUCT REGISTRATION HOLDER
Abio Marketing Sdn Bhd
No. 2, Jalan SS 13/5, 47500 Subang Jaya, Malaysia.

MANUFACTURER
Vefa İlaç Sanayi ve Ticaret Anonim Şirketi
Beylikdüzü OSB Mah. 2. Cadde No: 3/2 Beylikdüzü/İSTANBUL/Turkey.

REVISION DATE
May 2025

Y X Kesim / Cut-line			Remeco PEPTICON Liquid Oral Suspension KT Malezya VEFA İLAÇ			
			Barkod		Pharma Code	
Kağıt Cinsi			60gr. 1.Hamur			
Ölçüler (mm)			Ek Özellikler		KIRIMSIZ	
X	Y	Z	Renk	black	K	
170	240					
			Versiyon	Açıklama	Malzeme Kodu	
			REMPEPLİ-MAL-OS-200-KT.00	KT Tasarımı	12011924	