

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

MAVLAB KETAMAV 100 ANAESTHETIC IMMOBILISER SOLUTION FOR INJECTION

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

Active substance: Ketamine (as hydrochloride) 100 mg/mL

Excipient(s): Chlorocresol 2 mg/mL as preservative

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection

Clear, colorless liquid, particle free

4. CLINICAL PARTICULARS

4.1 Target species

Cats, Equine, Birds, Echidna, Marsupials, Reptiles and Zoo Felidae

Indications

For anaesthesia & immobilisation of cats, horses, birds, echidnas, marsupials, reptiles, zoo felidae

4.2 Contraindications

Do not use in animals presenting with:

- severe hypertension,
- cardio-respiratory deficiency,
- hepatic or renal dysfunction.

Do not use in animals with glaucoma.

Do not use in animals with eclampsia or pre-eclampsia.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use the product as a sole anaesthetic agent in any of the target species.

Do not use for surgical intervention on pharynx, larynx, trachea or bronchial tree, if sufficient relaxation is not ensured by administration of a muscle relaxant (intubation obligatory).

Do not use in ocular surgical interventions.

Do not use in animals undergoing a myelogram procedure.

Do not use in patients with pheochromocytoma or untreated hyperthyroidism.

4.3 Warnings and Precautions

Do not use in horses intended for human consumption.

4.4 Recommended Dosage/Instructions for Use

Cats- Minor surgery 11-22mg/kg bodyweight (BW) IM, Castration- 22-33mg/kg BW IM, Major surgery- 33-44mg/kg BW IM.

Horses- 2.2mg/kg BW IM, precede with xylazine 1.1mg/kg.

Birds- weighing less than 100g BW, give 0.1-0.2 mg/g BW IM, weighing 250-500g BW, give 0.05-0.1mg BW IM, weighing 500-1000g BW, give 0.02-0.1 mg/g BW IM, weighing more than 1kg, give 0.02-0.05mg/g BW IM.

Echidna- 20mg/kg BW

Marsupials- 15mg/kg BW



Reptiles- 22-88mg/kg BW

Zoo Felidae- By IM Injection, mixture of xylazine 1m/kg and Ketamav 5-15mg/kg BW

4.5 Side Effects

Emergence reactions - ataxia, hypersensitivity to stimuli, excitation – have been reported in horses rarely.

Salivation in cats has been reported very rarely. Increased tonus of skeletal muscles in cats, horses has been reported very rarely. Dose-related respiratory depression, which may lead to respiratory arrest in cats has been reported very rarely. Combination with respiratory depressant products may increase this respiratory effect.

Increased heart rate in cats has been reported very rarely.

Eyes that remain open with mydriasis and nystagmus have been reported in cats very rarely.

Pain on intramuscular injection in cats has been reported very rarely. All adverse reactions and frequencies are from spontaneous reports.

The frequency of adverse reactions is defined using the following convention: - very common (more than 1 in 10 animals treated displaying adverse reaction(s)) - common (more than 1 but less than 10 animals in 100 animals treated) - uncommon (more than 1 but less than 10 animals in 1,000 animals treated) - rare (more than 1 but less than 10 animals in 10,000 animals treated) - very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

4.6 Pregnancy and Lactation

Ketamine passes the blood placenta barrier very well to enter the fetal blood circulation in which 75 to 100 % of the maternal blood levels can be reached. This partially anesthetizes neonates delivered by caesarean section. The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

4.7 Interaction with Other Medications

Neuroleptics, tranquillisers, cimetidine and chloramphenicol potentiate ketamine anaesthesia. Barbiturates, opiates and diazepam may prolong the recovery time. Effects may be potentiated. A decrease of the dose of one or both agents may be necessary. There is a possibility of an increased risk of cardiac arrhythmia when ketamine is used in combination with thiopental or halothane. Halothane prolongs the half-life of ketamine. The simultaneous intravenous administration of a spasmolytic agent may provoke a collapse. Theophylline, when given with ketamine, may provoke an increase of epileptic crises. When detomidine is used together with ketamine, the recovery is slower than when ketamine is used alone.

4.8 Route of Administration

Intramuscular (IM)

4.9 Symptoms and Treatment of Overdose

Respiratory depression at high doses.

4.10 Withholding period

Do Not use in horses intended for human consumption.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: anaesthetics, other general anaesthetics, ketamine

ATC vet code: QN01AX03.

5.1 Pharmacodynamic properties

Ketamine is a dissociative anaesthetic agent. Ketamine induces a state of catalepsy with amnesia and analgesia; muscle tone is maintained including the pharyngeal and laryngeal reflexes. The heart rate,



blood pressure and cardiac output are increased; respiratory depression is not a noticeable feature. All these characteristics may be modified if the product is used in combination with other agents.

5.2 Pharmacokinetics properties

Ketamine is distributed rapidly. Distribution to tissues is variable with highest concentrations found in the liver and kidneys. Plasma protein binding is approximately 50%. Hepatic metabolism varies between species: for example, ketamine undergoes extensive hepatic biotransformation in dogs and horses, but the majority of the drug is excreted via the kidneys in cats. Recovery from ketamine anaesthesia after an intravenous bolus occurs by rapid redistribution from the CNS to other tissues, primarily fat, lung, liver and kidney.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Chlorocresol
RO water
Sodium Hydroxide BP
Hydrochloric acid BP

6.2 Major incompatibilities

Due to a chemical incompatibility, do not mix barbiturates or diazepam with ketamine in the same syringe.

6.3 Shelf life

Shelf Life: 2 years

Shelf Life after first opening of container: Use within 24 hours after first broaching

6.4 Storage Conditions

Store below 30°C (room temperature). Protect from light.

6.5 Nature and composition of immediate packaging

Amber type I glass vials containing 20 ml and 50 ml product, closed with an aseptic yellow flip top injection seal septa 20mm water/auto-hard

6.6 Disposal

Dispose of empty container by wrapping with paper and putting in garbage.

7. Product Registration Holder

Wiltop Enterprise
No 8, Jalan Silc 1/12,
Kawasan Perindustrian Silc,
79200 Iskandar Puteri, Johor

8. Manufacturer

Mavlab Pty Ltd
29-33 Rowland Street, Stacks Creek,
Queensland Australia

9. Date of Revision of Package

Revision 6, 10/02/2025

