



RefluxD⁺⁺

Suspension

Heartburn / Indigestion / Gastroesophageal reflux



1. Name of the medicinal product
RefluxD Suspension

2. Qualitative and quantitative composition
Each 10 ml contains sodium alginate 500mg, sodium bicarbonate 213mg and calcium carbonate 325mg.

For full list of excipients, see Section 6.1.

3. Product Description
Pale yellow to cream colored oral suspension with peppermint flavor in sachets.

4. Clinical particulars
4.1 Therapeutic indications
Treatment of acid related symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, for example following meals or during pregnancy.

4.2 Posology and method of administration
For oral administration.
Adults and children 12 years and over: One to two sachets (10-20ml) after meals and at bedtime, up to four times per day.
Children under 12 years: Should be given only on medical advice.

4.3 Contraindications
Patient with history of hypersensitivity to any of the ingredients in this medicine.

4.4 Special warnings and precautions for use
Patient listed below should consult with doctor or pharmacist before administration:

1. Patient with congestive heart failure or renal failure who must care about sodium (This medicine contains 254.5mg (11.06mmol) of sodium per 20mL) .
2. Patient with hypercalcemia, nephrocalcinosis and recurrent calcium containing renal calculi (This medicine contains 260mg (6.5mmol) of calcium per 20mL) .
3. In administration of this drug to the children under 12 years, should get diagnosis from doctor.

4.5 Interaction with other medicinal products and other forms of interaction
Due to the presence of calcium carbonate which act as an antacid, a time-interval of 2 hours should be considered between RefluxD Suspension intake an the administration of other medicinal products, especially H2-antihistaminics tetracyclines, digoxine, fluoroquinolone, iron salt, ketoconazole, neuroleptics, thyroxine, penicilamine, beta-blockers (atenolol, metoprolol, propanolol), glucocorticoid, chloroquine, and diphosphonates.

4.6 Pregnancy and lactation
There is no any significant adverse effects of the product known on the course of pregnancy or on the health of the foetus/new-born child. This medicinal product may be used during pregnancy and lactation.

4.7 Effects on ability to drive and use machines
Not relevant.

4.8 Undesirable effects
Adverse events which have been associated with sodium alginate, sodium bicarbonate and calcium carbonate are given below, tabulated by system organ class and frequency. Frequencies are defined as: Very common (>1/10); Common (>1/100 and <1/10); Uncommon (>1/1000 and <1/100); Rare (>1/10,000 and <1/1000); Very rare (< 1/10,000); Not known (cannot be estimated from the available data). Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

System Organ Class	Frequency	Adverse Events
Immune System Disorders	Very Rare	Anaphylactic reaction, anaphylactoid reaction. Hypersensitivity reactions such as urticaria.
Metabolism and Nutritional Disorders	Not Known	Alkalosis ¹ , Hypercalcaemia ¹ , Milk-alkali Syndrome ¹
Respiratory, Thoracic and Mediastinal Disorders	Not Known	Respiratory effects such as bronchospasm.
Gastrointestinal Disorders	Very Rare	Abdominal pain, acid rebound, diarrhoea, nausea, vomiting
	Not Known	Constipation ¹
Skin and Subcutaneous Tissue Disorders	Very Rare	Rash Pruritic

1 Usually occurs following larger than recommended dosages.

4.9 Overdose
In the event of overdose, symptomatic treatment should be given. The patient may notice abdominal distension.

5. Pharmacological properties
5.1 Pharmacodynamic properties
Pharmacotherapeutic classification: A02BX, other drugs for peptic ulcer and gastro-oesophageal reflux disease.
The medicinal product is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate.
On ingestion, the medicinal product reacts rapidly with gastric acid to form a raft of alginic acid gel having a near neutral pH and studies have shown that the raft interacts with and caps the acid pocket in the stomach, reducing oesophageal acid exposure. The raft floats on the stomach contents effectively impeding gastro-oesophageal reflux, for up to 4 hours, and protecting the oesophagus from acid, pepsin and bile. In severe cases the raft itself may be refluxed into the oesophagus, in preference to the stomach contents, and exert a demulcent effect. In addition, in vitro evidence has shown that the raft has a secondary action and is able to entrap bile and pepsin within it structure, further protecting the oesophagus from these gastric components. Calcium carbonate neutralises gastric acid to provide fast relief from indigestion and heartburn. This effect is increased by the addition of sodium bicarbonate which also has a neutralising action.

5.2 Pharmacokinetic properties
The mode of action of the medicinal product is physical and does not depend on absorption into the systemic circulation.

6. Pharmaceutical particulars
6.1 List of excipients
D-Sorbitol solution, Saccharin sodium hydrate, Enzymatically Modified Stevia, Steviol glucoside, Carbomer 934P, Sodium Hydroxide, Methylparaben, Propylparaben, Citrus flavor 91069, Peppermint Oil and purified water.

6.2 Incompatibilities
Not applicable.

6.3 Shelf life
Two years from manufacturing date.

6.4 Storage condition
Store at below 30°C.

6.5 Nature and contents of container
A cardboard outer carton containing unit dose stick pack style sachets.
Pack sizes: 5, 24 sachets /Box
The sachets are composed of polyester, aluminium and polyethylene.
Each sachet contains 10ml of RefluxD Suspension.

6.6 Special precautions for disposal and other handling
None required.

Product Registration Holder :


AiPharma Healthcare Sdn Bhd (1304105-X)
No. 83, Jalan I-Park 1/9, Kawasan Perindustrian I-Park,
Bandar Indahpura, 81000 Kulai, Johor, Malaysia.

Manufactured by :


GENUONE SCIENCES INC.
Manufacturing site : 245, Sandan-gil,
Jeonui-myeon, Sejong-si, Republic of Korea