

KETOVID CREAM 2%

DESCRIPTION

White to off white, opaque cream.

COMPOSITION

Ketoconazole 2% w/w.

ACTIONS AND PHARMACOLOGY

Ketoconazole is a synthetic imidazole dioxolane derivative which has a potent antimycotic activity against dermatophytes such as *Trichophyton sp.*, *Epidermophyton floccosum* and *Microsporum sp.* and against yeasts. The effect on *Pityrosporum sp.* is very pronounced. Ketoconazole cream usually acts very rapidly on pruritus which is commonly seen in dermatophyte and yeast infections as well as skin conditions related with the presence of *Pityrosporum sp.* This symptomatic improvement is observed before the first signs of healing are observed.

PHARMACOKINETICS

Ketovid Cream does not produce detectable blood levels after topical application.

INDICATIONS

Treatment of dermatophyte infections of the skin-tinea corporis, tinea cruris, tinea manus and tinea pedis caused by *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Microsporum canis* and *Epidermophyton floccosum* as well as in the treatment of cutaneous candidiasis, pityriasis versicolor and seborrhoeic dermatitis.

CONTRAINDICATIONS

Ketoconazole cream is contraindicated in individuals who have shown hypersensitivity to ketoconazole and any of its ingredients.

PRECAUTIONS

Ketovid cream is not for ophthalmic use. Irritation may occur when ketoconazole cream is used immediately after prolonged treatment with topical corticosteroids. Therefore, it is recommended to continue applying the topical corticosteroid in the mornings and to apply ketoconazole cream in the evenings, and to subsequently and gradually withdraw the steroid therapy over a period of 2-3 weeks. In case a potent steroid is used, replace it by a mild steroid and withdraw therapy over the same period.

PREGNANCY AND LACTATION

Not relevant since Ketovid Cream applied topically is not absorbed.

VIKET20-PC (MY)

MAIN SIDE / ADVERSE EFFECTS

A few instances of irritation and burning sensation have been observed during treatment with ketoconazole cream. In rare circumstances allergic local skin phenomena, such as contact dermatitis, have been associated with ketoconazole cream or its components like propylene glycol.

DRUG INTERACTIONS

No known drug interaction since Ketovid Cream applied topically is not absorbed.

OVERDOSE

Exaggerated topical application may lead to erythema, oedema and burning sensation, which will disappear upon discontinuation of the treatment.

Treatment for overdose: No special measures have to be taken in case of accidental ingestion.

DOSAGE AND ADMINISTRATION

Topical.

Cutaneous candidiasis, tinea corporis, tinea cruris, tinea manus, tinea pedis or tinea (pityriasis) versicolor: Apply to the affected skin and immediate surrounding areas, once a day.

Seborrhoeic dermatitis: Apply to the affected skin or areas, once or twice a day depending on the severity of infection.

Treatment should be continued for a sufficient period, at least until a few days after disappearance of all symptoms. The diagnosis should be reconsidered if no clinical improvement is noted after 4 weeks of treatment. General measures in regard to hygiene should be observed to control sources of infection or reinfection.

The usual duration of treatment is: tinea versicolor 2-3 weeks, yeast infections 2-3 weeks, tinea cruris 2-4 weeks, tinea corporis 3-4 weeks, tinea pedis 4-6 weeks. The usual duration of treatment for seborrhoeic dermatitis is 2 to 4 weeks. Maintenance therapy is applied once or twice weekly in seborrhoeic dermatitis.

Note: The information given here is limited. For further information, kindly consult your doctor or pharmacist.

Storage : Store below 30°C.
Presentation/Packing : 15g aluminium tube.

Product Registration Holder / Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: October 2017