

TIDACT® CAPSULE 300MG

Ingredient(s):

Each capsule contains:

Clindamycin Hydrochloride 300mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetics):

Pharmacodynamics

Clindamycin inhibits protein synthesis in susceptible bacteria by binding to the 50s subunits of bacterial ribosomes and preventing peptide bond formation. It is usually considered bacteriostatic, but may be bactericidal in high concentrations or when used against highly susceptible organisms.

Pharmacokinetics:

Clindamycin is active against most aerobic Gram-positive bacteria including *Streptococci*, *Staphylococci*, *Bacillus anthracis*, and *Corynebacterium diphtheriae*. It has good activity against a wide range of anaerobic bacteria. Susceptible Gram-positive anaerobes include *Eubacterium*, *Propionibacterium*, *Peptococcus*, *Peptostreptococcus*, and most strain of *Clostridium perfringens* and *Clostridium tetani*. The Gram-negative anaerobes that are susceptible to Clindamycin are *Fusobacterium sp.* and *Bacteriodes sp.*, including *B. fragilis* group.

About 90% of a dose of Clindamycin Hydrochloride is absorbed from the gastrointestinal tract and peak plasma concentration is achieved more rapidly than with Lincomycin. Concentrations of about 2.5µg/ml occur within 1 hour after a 150mg dose with average concentrations of about 0.7µg/ml after 6 hours. After a dose of 300 and 600mg, peak plasma concentration of 4 and 8 µg/ml, respectively, have been reported. The biological half-life is about 1.5 hours. Absorption is not significantly diminished by food but the rate of absorption may be reduced. Between 80% to 90% of Clindamycin in the circulation is bound to plasma proteins.

Clindamycin is widely distributed in body fluids and tissues including bone but does not reach the cerebrospinal fluid in significant concentrations. It diffuses across the placenta into the fetal circulation and has been reported to appear in breast milk. High concentration occurs in the bile.

Clindamycin undergoes metabolism, presumably in the liver, to the active N-dimethyl and sulfoxide metabolites and also some inactive metabolites. About 10% of a dose is excreted in the urine as active drug or metabolites, and about 4% is excreted in the faeces. It is not effectively removed from the blood by dialysis.

Indication(s):

Serious infections caused by susceptible anaerobic bacteria or strain of streptococci, pneumococci and staphylococci. Anaerobes: Serious respiratory tract infections such as empyema, anaerobic pneumonitis and lung abscess, serious skin and soft tissue infections; septicemia; intra-abdominal infections such as peritonitis and intra-abdominal abscess (typically resulting from anaerobic organisms resident in the normal gastrointestinal tract); infections of the female pelvis and genital tract such as endometritis, non-gonococcal tubo-ovarian abscess, pelvic cellulitis and post-surgical vaginal-cuff infection.

Streptococci: Serious respiratory tract infections, serious skin and soft tissue infections.

Pneumococci: Serious respiratory tract infections.

Dosage and Administration:

Adults: Serious infections ~ 150 to 300mg every 6 hours.

More severe infections ~ 300 to 450mg every 6 hours.

Children: Serious infections ~ 8 to 16mg/kg/day divided to 3 or 4 equal doses.

More severe infections ~ 16 to 20mg/kg/day divided to 3 or 4 equal doses.

Route of administration: Oral

Contraindication(s):

History of hypersensitivity to preparations containing clindamycin or lincomycin.

Item Code	: 204380 - XX
PM Code	: Ins.P TDC3
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: R 193 - R 195 mm
CURSOR (ATTACHMENT 2)	: L 205 - L 210 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC
Draft	: 14 Dec 2022

L
260mm
±
2mm

W
110mm ± 2mm

Side Effect(s) / Adverse Reaction(s):

Clindamycin may cause diarrhea which can be severe and persistent, nausea, vomiting, abdominal cramps, and abnormality of taste. Severe pseudomembranous colitis has occurred in some patients and occasionally has been fatal. Hypersensitivity reaction including skin rashes and urticaria may occur, and transient leucopenia and eosinophilia, abnormalities in liver function test, and jaundice have been reported. Agranulocytosis, thrombocytopenia, and erythema multiforme have been observed.

Precaution(s) / Warning(s):

1. Tidact Capsule should be used with caution in individuals with a history of gastrointestinal disease, particularly colitis.
2. It should not be used in patients with diarrheal states.
3. Caution should be exercised in patients with impaired liver and renal functions.
4. Since Clindamycin is reported to possess neuromuscular blocking activity, it should be used cautiously with other drugs having similar activity.
5. Safety for use in pregnancy has not been established. Clindamycin has been reported to appear in breast milk in ranges of 0.7-3.8 mcg/ml.

WARNINGS

1. Clindamycin therapy has been associated with severe colitis which may end fatally.
2. It should be reserved for serious infections where less toxic antimicrobial agents are inappropriate.
3. It should not be used in patients with nonbacterial infections, such as most upper respiratory tract infections.
4. Its use in newborns is contraindicated.

Interaction with Other Medicaments:

1. Antagonism has been demonstrated between clindamycin and erythromycin *in vitro*. Because of possible clinical significance, these 2 drugs should not be administered concurrently.
2. The concurrent use of kaolin-containing anti-diarrhoeal preparations markedly reduces the rate of absorption of Clindamycin but not the extent of absorption.

Pregnancy and Lactation:

Safety for use in pregnancy has not been established. Clindamycin has been reported to appear in breast milk in ranges of 0.7-3.8µg/mL.

Symptoms and Treatment for Overdosage, and Antidote(s):

There is no specific antidote for its overdose; therefore management of the patient consists of symptomatic and supportive therapy.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Product Description & Packing(s):

A size #0 light blue color hard gelatin capsule with the mark " YSP " on both cap and body. TDC₃

Blister packing of 10's x 10, 10's x 50 and 10's x 100.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
 Lot 3, 5 & 7, Jalan P/7, Section 13,
 Kawasan Perindustrian Bandar Baru Bangi,
 43000 Kajang, Selangor Darul Ehsan, Malaysia.
 Ordering Line: 1 800 88 3027
 Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 14 Dec 2022

Item Code	: 204380 - XX
PM Code	: Ins.P TDC3
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: R 193 - R 195 mm
CURSOR (ATTACHMENT 2)	: L 205 - L 210 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC
Draft	: 14 Dec 2022

L
260mm
±
2mm

W
110mm ± 2mm