

FROTIN[®] PESSARIES

Ingredient(s):

Each pessary contains:

Metronidazole 250 mg

Pharmacodynamics:

1. Metronidazole is a trichomonacidal agent. It has antiprotozoal and antibacterial actions and is effective against *Trichomonas vaginalis* and other protozoa including *Entamoeba histolytica* and against anaerobic bacteria.
2. Metronidazole is effective against most obligate anaerobes by undergoing intracellular chemical reduction via mechanisms unique to anaerobic metabolism. Reduced metronidazole, which is cytotoxic but short-lived, interact with DNA to cause a loss of helical structure, strand breakage and result inhibition of nucleic acid synthesis and cell death.
3. Metronidazole is a synthetic antibacterial agent with an average elimination half-life in healthy humans of eight hours.

Pharmacokinetics:

1. Metronidazole is widely distributed in body tissue. Maximum concentrations occur in the serum after about 1 hour and traces are detected after 24 hours.
2. When metronidazole is given as a pessary, the bioavailability is approximately 25%. Peak serum levels after an intra-vaginal dose is 15% of the equivalent oral dose.
3. At least half the dose is excreted in the urine as metronidazole and its metabolites, including an acid oxidation product, a hydroxy derivative and a glucuronide. Metronidazole diffuses across the placenta, and is found in breast milk of nursing mother in concentrations equivalent to those in the serum.

Indication(s):

For the treatment of vaginitis caused by *Trichomonas vaginalis*.

Dosage and Administration:

One pessary to be placed deeply into the vagina daily for 10 - 21 days.

To be used as directed by a physician.

Route of Administration:

Vaginal

Contraindication(s):

1. Known sensitivity to metronidazole.
2. Concurrent use of ethyl alcohol.
3. Active organic disease of CNS, including seizures, peripheral neuropathy, as metronidazole can cause CNS toxicity.
4. Blood dyscrasias, as metronidazole can cause leukopenia.
5. Women using metronidazole should not breast feed their babies.
6. It also should not be used during pregnancy.

Side Effect(s) / Adverse Reaction(s):

Side effects of metronidazole include anorexia, nausea, coated tongue, dry mouth and unpleasant taste, headache, pruritus and less frequently vomiting, diarrhoea, weakness, vertigo, ataxia, depression, insomnia, drowsiness, urethral discomfort and darkening of the urine. Occasionally there may be temporary moderate leukopenia. Peripheral neuropathy has been reported in patients on prolonged therapy.

Warnings and Precautions:

1. Regular clinical and biological surveillance are advised if administration of metronidazole for more than 10 days is considered to be necessary.
2. There is a possibility that after *Trichomonas vaginalis* has been eliminated a gonococcal infection might persist.
3. Metronidazole is mainly metabolised in liver. Hepatic dysfunction may lead to decrease plasma clearance and accumulation of metronidazole and its metabolites, dosage may need to reduce in severe hepatic impairment.
4. The elimination half-life of metronidazole remains unchanged in the presence of renal failure. Therefore, the dosage of metronidazole needs no reduction. However, its metabolites may retain in that patient. The clinical significance of this is not known at present.

5. Cases of severe hepatotoxicity/acute hepatic failure, including cases with a fatal outcome with very rapid onset after treatment initiation in patients with Cockayne syndrome have been reported with products containing metronidazole for systemic use. In this population, metronidazole should therefore be used after careful benefit-risk assessment and only if no alternative treatment is available. Liver function tests must be performed just prior to the start of therapy, throughout and after end of treatment until liver function is within normal ranges, or until the baseline values are reached. If the liver function tests become markedly elevated during treatment, the drug should be discontinued.

Patients with Cockayne syndrome should be advised to immediately report any symptoms of potential liver injury to their physician and stop taking metronidazole.

Pregnancy and Lactation:

There is inadequate evidence of the safety of metronidazole in pregnancy. This product should not be given during pregnancy or during lactation unless the physician considers it essential; in these circumstances the short, high-dosage regimens are not recommended.

Interaction with Other Medicaments:

1. Metronidazole should not be used concurrently with, or for at least 1 day following, ingestion of alcohol. This is because alcohol may interact with metronidazole to produce a disulfiram-like reaction in some individuals.
2. Metronidazole enhances the anti-coagulant effect of warfarin.
3. When cimetidine is used concurrently with metronidazole, hepatic metabolism of metronidazole may be decreased, resulting in increased serum concentration. Monitoring of serum concentration and dosage adjustment of metronidazole may be necessary.
4. Metronidazole may impair the clearance of phenytoin, increasing phenytoin's plasma concentration.
5. Phenobarbital may induce microsomal liver enzyme, increasing metabolism of metronidazole and resulting in a decrease in half life and plasma concentration.

Symptoms and Treatment for Overdosage, and Antidote(s):

Symptoms of overdose include nervous system toxicity such as ataxia, confusion, tremors, neuropathies and epileptiform seizures. Since there is no specific antidote treatment for metronidazole overdose should be symptomatic and supportive.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 25°C. Protect from light and moisture.

Product Description:

A pale orange pessary.

Packing(s):

10's X 10.

FROTIN® PESARI

Bahan Aktif:

Setiap pesari mengandungi:

Metronidazole 250 mg

Indikasi:

Untuk rawatan vaginitis yang disebabkan oleh *Trichomonas vaginalis*.

Dos:

Satu pesari dimasukkan ke dalam vagina setiap hari untuk selama 10 - 21 hari.

Digunakan seperti yang diarahkan oleh doktor.



Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,

Kawasan Perindustrian Bandar Baru Bangi,

43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679

FBO V1 - 02

Date of revision: 12 Nov 2025