

93 mm

Funginox Solution 2%

Description

Funginox Solution is a clear, pink liquid packed in an opaque glass bottle which is closed with a white plastic cap. A white plastic sprayer which is packed separately from the glass container is enclosed within the same cardboard box.

Name and Strength of Active Ingredient

Funginox Solution contains Ketoconazole 2%

Pharmacodynamics

Ketoconazole, a synthetic imidazole dioxolane derivative, is a broad-spectrum antifungal agent. It has a potent antimycotic activity against dermatophytes namely, Trichophyton sp., Epidermophyton floccosum and microsporum sp. and yeasts. The effect on Pityrosporum sp.is especially very distinctive.

Pharmacokinetics

Percutaneous absorption of Ketoconazole topical solution spray is negligible since blood levels cannot be detected even after prolonged use. There should not be any systemic effect.

Indication

Topical treatment and prophylaxis of dermatophyte infections of the skin: Tinea Corporis, tinea cruris, tinea menus and tinea pedis due to Trichophyton rubrum, Trichophyton mentagrophytes, Microsporum canis, Epidermophyton floccosum, cutaneous candidosis and tinea (pityriasis) versicolor, seborrhoeic dermatitis (a skin condition related with the presence of Pityrosporum ovale). Topical treatment and prophylaxis of infections of the scalp in which the yeast pityrosporum is involved e.g. pityriasis versicolor (localized), seborrhoeic dermatitis and pityriasis capitis (dandruff associated with fungal infections of the scalp).

Recommended Dose

For topical treatment of dermatophyte infections of the skin: Tinea corporis, tinea cruris, tinea manus, tinea pedis, cutaneous candidosis and pityriasis versicolor : it is recommended that Funginox Solution should be gently sprayed or applied once daily to cover the affected and immediate surrounding area.

Seborrheic Dermatitis: Funginox Solution should be sprayed or applied to the affected area once or twice daily depending on the severity of infection.

Dandruff: Funginox Solution to be sprayed or applied on the scalp twice a week. Treatment should be continued for a sufficient period, at least until a few days after disappearance of all symptoms. The diagnosis should be reassessed if no clinical improvement is observed after 4 weeks of treatment. General requirements in terms of personal hygiene should be observed to control sources of infection or reinfection.

The usual duration of treatment is : Tinea corporis 3-4 weeks, tinea cruris and tinea manus 2-4 weeks, tinea pedis 4-6 weeks, yeast infections 2-3 weeks, pityriasis versicolor 2-3 weeks.

The usual duration of treatment in seborrhoeic dermatitis is 4 weeks and its maintenance therapy is application of once or twice weekly.

For topical treatment of infections of the scalp: Pityriasis versicolor (localized): Once daily for 5 days, Seborrhoeic Dermatitis and Pityriasis capitis: Twice weekly for 3-4 weeks.

For prophylaxis of infections of the scalp: Pityriasis versicolor: Once daily for 3 days during a single treatment course before exposure to sunshine. Seborrhoeic Dermatitis and Pityriasis capitis: Once every 1 or 2 weeks.

Route of Administration

Spray or apply topically on affected areas as directed.

Contraindication

Known hypersensitivity to Ketoconazole or any of the excipients of Funginox Solution.

Warning and Precautions

For external use only. Avoid contact with eyes. Discontinue if sensitivity or chemical irritation occurs. This preparation contains Sodium Metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to any bisulphites.

Interactions with other Medicaments

No known drug interaction has been reported since Funginox Solution applied topically is not absorbed systemically.

Pregnancy and Lactation

Since Ketoconazole is not absorbed systemically after topical application, it is not contraindicated in pregnancy and lactation.

Side Effects

Irritation of the skin may occur.

Symptoms and Treatment of Overdose

Overdosage is not expected as Funginox Solution is intended for external use only. In the event of accidental ingestion, only supportive measures should be carried out. In order to avoid aspiration, neither emesis nor gastric lavage should be performed.

Storage Condition

Store below 30°C. Avoid direct sunlight or heat.

Shelf Life

3 years from manufacturing date.

Packing: 25ml packed in an opaque glass bottle is closed with a white plastic cap. A white plastic sprayer in the plastic bag which is packed separately from the glass container is enclosed within the same individual cardboard box.

Registration No.: MAL05090003AZ

Manufacturer

Charoen Bhaesaj Lab Co., Ltd.
32 Soi Phetkasem 21, Phetkasem Road., Pakklong Phasi Charoen, Phasi Charoen, Bangkok 10160, Thailand.

Name of Importer and Distributor:

Propharm (M) Sdn.Bhd. No.8, Jalan Udang Harimau 2, Medan Niaga Kepong,51200 Kuala lumpur, Malaysia.
Tel: 03-62436389
Fax: 03-62436385 e-mail: propharm@gmail.com

Date of Revision : April 15, 2021

AW-PI-EO-001 REV003

296 mm

Black 100

