



ADAZOL suspension

Description: White, orange flavoured suspension.

Content:

Each 5ml contains albendazole 200mg

Indication:

It is used in the treatment of single and mixed infestations of roundworm, pinworm, hookworm, threadworm (strongyloidiasis), whipworm and tapeworm.

Pharmacology:

Albendazole, a benzimidazole carbamate, which exhibits vermifugal, ovicidal and larvacidal activity and exerts its anthelmintic effect by inhibiting tubulin polymerization. This causes the disruption of the helminth metabolism, incl. energy depletion, which immobilises and kills the susceptible helminth.

It is poorly absorbed from the gastrointestinal tract but rapidly undergoes extensive first-pass metabolism. The principal metabolite albendazole sulphoxide has anthelmintic activity and plasma half-life of about 8.5 hours. It is excreted in the urine together with other metabolites; albendazole sulphone, 2-amino sulphoxide and 2-amino sulphone.

Contraindications:

It is contraindicated in pregnant ladies, children less than 1 year old and patients hypersensitive to albendazole.

Side effects/Adverse reactions:

Most common side effects are gastrointestinal disturbances like nausea, vomiting and diarrhoea. The other side effects are like headache, fever and fatigue.

Precautions:

It should not be administered during confirmed or suspected pregnancy. If symptoms still persist after treatment, please consult the doctor. Albendazole should only be used in the treatment of hydatid disease if there is constant medical supervision with regular monitoring of serum transaminase concentrations and of leukocyte and platelet counts.

Use in pregnancy and lactation:

It is contraindicated in pregnant ladies. Albendazole can be absorbed from breast milk. Therefore, breast feeding should be stopped while taking albendazole and for at least 5 days after finishing treatment.

Drug Interactions:

Nil.

Dosage and administration:

Adult and children over 2 years:

Ascariasis, hookworm infections and trichuriasis: 400mg as a single dose.

Strongyloidiasis, taeniasis when mixed with other worm infections: 400mg given daily for 3 consecutive days.

May be repeated after 3 weeks, if necessary.

Children 1 - 2 years: Half of adult dose.

Treatment of over dosage

Symptoms: Gastrointestinal disturbances i.e. nausea, vomiting and diarrhoea.

Treatment: Symptomatic and supportive.

Storage conditions : Keep container tightly closed. Store in a dry place at 30°C. Protect from light. Keep away from children.

Shelf life : 3 years

Presentation : 10ml

Manufactured & : **PRIME PHARMACEUTICAL SDN. BHD.**

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

17/03/2021