

Item Code	: 204xxx draft 05 August 2021
PM Code	: Ins.P. CMS (TW)
Measurement(mm) ± 2mm	: 180(L) x 150(W)
Colour of Artwork	: K
Source of Artwork	: MAC 3 / CC

Clamentin® Powder for Syrup

31.25mg/mL

Ingredient(s):

Each mL contains:

Amoxicillin(asTrihydrate).....25mg
Clavulanic Acid (as Potassium Salt)6.25mg

Pharmacodynamics:

Resistance to many antibiotics is caused by bacterial enzymes which destroy the antibiotic before it can act on the pathogen. The clavulanate in Clamentin anticipates this defense mechanism by blocking the β -lactamase enzymes, thus rendering the organisms sensitive to Amoxicillin's rapid bactericidal effect at concentrations readily attainable in the body.

Clamentin is bactericidal to a wide range of organisms including:

Gram-positive:

Aerobes: **Enterococcus faecalis*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus viridans*, **Staphylococcus aureus*, *Coagulase-negative staphylococci (including *Staphylococcus epidermidis*), *Corynebacterium* sp, **Bacillus anthracis*, *Listeria monocytogenes*, *Enterococcus faecium*, *Streptococcus agalactiae*, *Streptococcus* sp, Anaerobes: *Clostridium*, *Peptococcus* spp, *Peptostreptococcus*.

Gram-negative:

Aerobes: **Haemophilus influenzae*, **Escherichia coli*, **Proteus mirabilis*, **Proteus vulgaris*, **Klebsiella* sp, **Moraxella catarrhalis* (*Branhamella catharralis*), **Salmonella* sp, **Shigella* sp, *Bordetella pertussis*, *Brucella* sp, **Neisseria gonorrhoeae*, **Neisseria meningitidis*, *Vibrio cholerae*, **Yersinia enterocolitica*, *Pasteurella multocida*, *Gardnerella vaginalis*, *Helicobacter pylori*, *Legionella* sp.

Anaerobes: **Bacteroides* sp (including *Bacteroides fragilis*), **Fusobacterium* sp.

* Including β -lactamase producing strains resistant to ampicillin and Amoxicillin.

Pharmacokinetics:

Amoxicillin and clavulanate potassium are well absorbed from the gastrointestinal tract after oral administration. Dosing in the fasted or fed state has minimal effect on the pharmacokinetics of Amoxicillin. While Clamentin can be given without regard to meals, absorption of clavulanate acid when taken with food is greater relative to the fasted state. Amoxicillin serum concentrations achieved with Clamentin are similar to those produced by the oral administration of equivalent doses of amoxicillin alone. The half-life of amoxicillin after oral administration is 1.3 hours and that of clavulanate acid is 1.0 hour. Concurrent administration of probenecid delays Amoxicillin excretion but does not delay renal excretion of clavulanate acid. Clavulanate acid has been found to be approximately 25% bound to human serum and Amoxicillin approximately 18% bound. Amoxicillin diffuses readily into most body tissues and fluids with the exception of the brain and spinal fluid. The results of experiments involving the administration of clavulanate acid to animals suggest that this compound, like Amoxicillin, is well distributed in body tissues.

Indication(s):

For the treatment of infections caused by staphylococci, *streptococcus pneumonia*, *streptococcus meningitis* and other susceptible pathogens, Clamentin is indicated for short-term treatment of bacterial infections at the following sites: Upper Respiratory Tract Infections e.g. Sinusitis, tonsillitis, otitis media. Lower Respiratory Tract Infections e.g. Acute and chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, Skin and Soft Tissue Infections e.g. Boils, abscesses, cellulitis, wound infections, intra-abdominal sepsis. Genito-Urinary Tract Infections e.g. Cystitis, urethritis, pyelonephritis, septic abortion, puerperal sepsis, pelvic infections, chancroid, gonorrhoea. Other Infections e.g. Osteomyelitis, septicaemia, peritonitis, post-operative infections.

Dosage and Administration:

Children

7-12 years: 10ml three times a day.

2-7 years: 5ml three times a day.

Children 9 months-2 years: 25mg/kg/day, in three divided doses.

Severe infections: Increase frequency to 6 hourly intervals.

Renal impairment:

For children with a creatinine clearance >30 mL/min, no changes in dosage is required.

For children with a creatinine clearance <30mL/min, Clamentin syrup is not recommended.

Hepatic impairment:

Dose with caution; monitor hepatic function at regular intervals. There is, as yet, insufficient data on which to base a dosage recommendation.

Directions for mixing:

Prepare syrup at time of dispensing as follows: Tap bottle until all powder flows freely. Add approximately 2/3 of the total amount of water for reconstitution and shake vigorously to suspend powder. Add remaining water to 100mL and again shake vigorously.

Shake syrup well before using. Reconstituted syrup must be stored under refrigeration and discarded after 7 days.

Route of Administration

For oral use.

Contraindication(s):

1. Penicillin hypersensitivity.
2. Attention should be paid to possible cross-sensitivity with other β -lactam antibiotics, e.g. cephalosporins.
3. A previous history of Clamentin or penicillin-associated jaundice/hepatic dysfunction.

Precaution(s) / Warning(s):

1. Clamentin should be used with caution in patients with evidence of hepatic dysfunction.
2. Cholestatic jaundice, which may be severe, but is usually reversible, has been reported rarely. Signs and symptoms may not become apparent for up to six weeks after treatment has ceased.
3. In patients with moderate or severe renal impairment, dosage should be adjusted.
4. Serious and occasionally fatal hypersensitivity (anaphylactoid) reactions have been reported in patients on penicillin therapy. These reactions are more likely to occur in individuals with a history of penicillin hypersensitivity.

L
180mm

W
150mm

Item Code	: 204xxx draft 05 August 2021
PM Code	: Ins.P. CMS (TW)
Measurement(mm) ± 2mm	: 180(L) x 150(W)
Colour of Artwork	: K
Source of Artwork	: MAC 3 / CC

5. Erythematous rashes have been associated with glandular fever in patients following the use of Amoxicillin. Clamentin should be avoided if glandular fever is suspected.
6. During the administration of high doses, adequate fluid intake and urinary output should be maintained to minimize the possibility of crystalluria.
7. Prolonged use may also occasionally result in overgrowth of non-susceptible organisms.
8. Clamentin syrup contains aspartame, therefore it is unsuitable for phenylketonurics.
9. Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients on penicillin therapy.

Interaction with Other Medicaments

In common with other broad-spectrum antibiotics, Clamentin may reduce the efficacy of oral contraceptives and patients should be warned accordingly. Concomitant use of Probenecid is not recommended. Probenecid decreases the renal tubular secretion of Amoxicillin. Concomitant use with Clamentin may result in increased and prolonged blood levels of Amoxicillin but not of clavulanic acid. Concomitant use of allopurinol during treatment with Amoxicillin can increase the likelihood of allergic skin reactions. There are no data on the concomitant use of Clamentin and allopurinol.

Pregnancy and Lactation

1. Use in pregnancy
Reproduction studies in animals (mice and rats) with orally and parenterally administered Clamentin powder for syrup 31.25mg/ml have shown no teratogenic effects. As with all medicines, use should be avoided in pregnancy, especially during the first trimester, unless considered essential by the physician.
2. Use in lactation
Clamentin powder for syrup 31.25mg/ml may be administered during the period of lactation. With the exception of the risk of sensitization associated with the excretion of trace quantities in breast milk, there are no detrimental effects for the infant.

Side Effect(s) / Adverse Reaction(s):

Side effects, as with Amoxicillin, are uncommon and mainly of a mild and transitory nature.

Gastrointestinal reactions

Diarrhea, indigestion, nausea, vomiting, candidiasis and antibiotic-associated colitis have been reported rarely. Nausea, although uncommon, is more often associated with higher oral dosages.

Hepatic effects

A moderate rise in AST and/or ALT has been noted in patients with beta-lactam class antibiotics, but the significance of these findings is unknown. Hepatitis and cholestatic jaundice have been reported rarely. They may, however, be severe and continue for several months. They are reported as occurring predominantly in adult or elderly patients and slightly more frequently in males. Signs and symptoms may occur during treatment but are more frequently reported after cessation of therapy with a delay of up to six weeks. The hepatic events are usually reversible. However, in extremely rare circumstances, deaths have been reported.

Hypersensitivity reactions

Urticarial and erythematous rashes sometimes occur. Rarely erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis and exfoliative dermatitis have been reported. Treatment should be discontinued if one these types of rash appear. In common with others β -lactam antibiotics, angioedema, oedema, anaphylaxis, serum sickness-like syndrome and hypersensitivity vasculitis have been reported. Interstitial nephritis can occur rarely.

Skin and subcutaneous tissue disorders

Frequency 'very rare': Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

Haematological effects

As with other β -lactams, reversible leucopenia, reversible thrombocytopenia and haemolytic anemia have been reported rarely.

CNS effects

Hyperactivity, dizziness, and headache are rare. Convulsions may occur with impaired renal functions or in those receiving high doses.

Local effects

Phlebitis at the site of injection.

Symptoms and Treatment of Overdose

Cases of overdosage with Amoxicillin-clavulanic acid are usually asymptomatic. If encountered, gastrointestinal symptoms and disturbance of the fluid and electrolyte balances may be evident. They may be treated symptomatically with attention to the water electrolyte balance. Clamentin powder for syrup 31.25mg/ml can be removed from the circulation by haemodialysis.

Shelf-Life:

2 years from the date of manufacture.

Reconstituted Solution: 7 days after reconstitution.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture. Reconstituted Solution: Store at temperature 2°C ~ 8°C.

Product Description and Packing(s):

A white to off-white powder.

Appearance after reconstitution:

An off-white suspension.

Glass bottle of 100ml.



Manufacturer:
YUNG SHIN PHARMACEUTICAL IND. CO., LTD.
Taichung Youth Factory III
No. 26, Kung Liu Rd.,
Tachia, Taichung, Taiwan, R.O.C.
Importer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD.
Lot 3, 5 & 7, Jln. P/7, Sec. 13,
Kaw. Perindustrian Bdr. Baru Bangi,
43000 Kajang, Selangor D.E., Malaysia
Ordering Line : 1 800 88 3027
Product Info : 1 800 88 3679
xxxxxx printing code xxxxxx

Date of revision: 05 August 2021

L
180mm

W
150mm