

70 mm

70 mm

Pharmacode 298

Pharmacode 298

55 mm

220mm

Pharmacode 298

Pharmacode 298

Pharmacode 298

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory.

Myclav 625

Amoxicillin and Clavulanate Potassium Tablets USP

COMPOSITION:

Each film coated tablet contains:

Amoxicillin Trihydrate

equivalent to Amoxicillin..... 500 mg

Clavulanate potassium

equivalent to Clavulanic acid.... 125 mg

DESCRIPTION OF PRODUCT:

Myclav - 625: White, film-coated, oval shaped, plain biconvex tablets with thickness-6.5 mm to 7.0 mm, Length-19.50 mm to 19.7mm, width-9.5 mm to 9.7 mm.

CLINICAL PHARMACOLOGY:

PHARMACODYNAMICS:

Resistance to many antibiotics is caused by bacterial enzymes which destroy the antibiotic before it can act on the pathogen. The clavulanate in **Myclav** anticipates this defence mechanism by blocking the β-lactamase enzymes, thus rendering the organisms sensitive to amoxicillin's rapid bactericidal effect at concentrations readily attainable in the body. Clavulanate by itself has little antibacterial activity; however, in association with amoxicillin as **Myclav**, it produces an antibiotic agent of broad spectrum with wide application in hospital and general practice.

Myclav is bactericidal to a wide range of organisms including:

Gram-positive

Aerobes: *Enterococcus faecalis**, *Enterococcus faecium**, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus viridans*, *Staphylococcus aureus**, *Coagulase negative staphylococci** (including *Staphylococcus epidermidis**), *Corynebacterium species*, *Bacillus anthracis**, *Listeria monocytogenes*.

Anaerobes: *Clostridium species*, *Peptococcus species*, *Peptostreptococcus*.

Gram-negative

Aerobes: *Haemophilus influenzae**, *Moraxella catarrhalis** (*Branhamella catarrhalis*), *Escherichia coli**, *Proteus mirabilis**, *Proteus vulgaris**, *Klebsiella species**, *Salmonella species**, *Shigella species**, *Bordetella pertussis*, *Brucella species*, *Neisseria gonorrhoeae**, *Neisseria meningitidis**, *Vibrio cholerae*, *Pasteurella multocida*.

Anaerobes: *Bacteroides species** including *B. fragilis*.

* Some members of these species of bacteria produce beta-lactamase, rendering them insensitive to amoxicillin alone.

PHARMACOKINETICS:

The pharmacokinetics of the two components of **Myclav** are closely matched. Peak serum levels of both occur about one hour after oral administration. Absorption of **Myclav** is optimised at the start of a meal. Both clavulanate and amoxicillin have low levels of serum binding; about 70% remains free in the serum.

Doubling the dosage of **Myclav** approximately doubles the serum levels achieved.

INDICATIONS:

Amoxicillin/clavulanate potassium is indicated in the treatment of infections caused by susceptible strains of the designated organisms in the conditions listed below:

Upper respiratory tract infections (including ENT) e.g. Tonsillitis, sinusitis, otitis media.

Lower respiratory tract infections e.g. Acute exacerbation of chronic bronchitis, lobar and bronchopneumonia.

Genito-urinary tract infections e.g. Cystitis, urethritis, pyelonephritis.

Skin and soft tissue infections, e.g. boils, abscesses, cellulitis, wound infections.

Bone and joint infections e.g. Osteomyelitis.

Dental infections e.g. dento alveolar abscess.

Other infections e.g. septic abortion, puerperal sepsis, intra-abdominal sepsis.

RECOMMENDED DOSE:

Usual dosages for the treatment of infection:

Adults and children over 12 years	One Myclav 625 mg Tablet two times daily; Severe infections : One Myclav 625 mg Tablet three times daily.
-----------------------------------	--

Not recommended in children of 12 years and under.

Dosage in renal impairment:

Adults:

Mild impairment (Creatinine clearance>30 ml/min)	Moderate impairment (Creatinine clearance 10-30 ml/min)	Severe impairment (Creatinine clearance<10 ml/min)
No change in dosage	One 625 mg tablet 12 hourly.	Not recommended.

Dosage in hepatic impairment:

Dose with caution; monitor hepatic function at regular intervals.

ADMINISTRATION:

Oral Tablets: To minimise potential gastrointestinal intolerance, administer at the start of a meal. The absorption of **Myclav** is optimised when taken at the start of a meal.

Duration of therapy should be appropriate to the indication and should not exceed 14 days without review.

CONTRAINDICATIONS:

Patients with penicillin allergy should not be given Amoxicillin/clavulanic acid, as the Amoxicillin component is likely to precipitate a hypersensitivity reaction.

Patients with suspected glandular fever, or pharyngitis who may be suffering from this condition, should not be given Amoxicillin/clavulanic acid, as the Amoxicillin component is likely to cause a maculopapular rash. This should not prove a

therapeutic problem as there is no reason to use this combination for the treatment of pharyngitis.

Contraindicated in patients with a previous history of Amoxicillin/ clavulanate potassium-associated cholestatic jaundice/hepatic dysfunction.

WARNING:

Before initiating therapy with Amoxicillin/clavulanic acid, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, or other allergens. If an allergic reaction occurs, Amoxicillin should be discontinued and appropriate therapy instituted. Serious anaphylactic reactions require immediate emergency treatment with epinephrine. Oxygen, intravenous steroids, and airway management, including intubation, should also be administered as indicated.

Pseudomembranous colitis may occur with Amoxicillin/clavulanic acid, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of Amoxicillin/clavulanic acid.

Treatment with Amoxicillin/clavulanic acid alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by Clostridium difficile is a primary cause of "antibiotic-associated colitis."

After the diagnosis of pseudomembranous colitis has been established, appropriate therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial drug clinically effective against Clostridium difficile colitis.

Amoxicillin/clavulanate potassium should be used with caution in patients with evidence of hepatic dysfunction. Hepatic toxicity associated with the use of Amoxicillin/clavulanate potassium is usually reversible. On rare occasions, deaths have been reported (less than one death reported per estimated 4 million prescriptions worldwide). These have generally been cases associated with serious underlying diseases or concomitant medications.

PRECAUTIONS:

While Amoxicillin; clavulanate potassium possesses the characteristic low toxicity of the penicillin group of antibiotics, periodic assessment of organ system functions, including renal, hepatic and hematopoietic function, is advisable during prolonged therapy.

A high percentage of patients with mononucleosis who receive ampicillin develop an erythematous skin rash. Thus, Amoxicillin/clavulanic acid should not be administered to patients with mononucleosis.

The possibility of super infections with mycotic or bacterial pathogens should be kept in mind during therapy. If super infections occur (usually involving Pseudomonas or Candida), Amoxicillin/clavulanic acid should be discontinued and/or appropriate therapy instituted.

Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients receiving therapy with beta-lactams. Before initiating therapy with Myclav 625, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, carbapenems or other beta-lactam agents. If an allergic reaction occurs, Myclav 625 must be discontinued immediately and appropriate alternative therapy instituted.

INTERACTION WITH OTHER MEDICAMENTS:

Probenecid decreases the renal tubular secretion of Amoxicillin. Concurrent use with Amoxicillin; clavulanate potassium may result in increased and prolonged blood levels of Amoxicillin. Co-administration of probenecid cannot be recommended.

There is no data available on the concurrent administration of Amoxicillin/clavulanate potassium and allopurinol.

In common with other broad-spectrum antibiotics, Amoxicillin/clavulanate potassium may reduce the efficacy of oral contraceptives.

PREGNANCY AND LACTATION:

Caution should be exercised when amoxicillin; clavulanate potassium is administered to pregnant women and nursing mothers.

SIDE EFFECTS:

The adverse drug reactions with Amoxicillin/clavulanic acid are diarrhea/loose stools, nausea, skin rashes and urticaria, vomiting and vaginitis. The overall incidence of side effects, and in particular diarrhea, increased with the higher recommended dose.

Skin and subcutaneous tissue disorders:

Frequency 'very rare': Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS).

SYMPTOMS AND TREATMENT OF OVERDOSE:

Overdosage may include asymptomatic symptoms such as gastrointestinal symptoms including stomach and abdominal pain, vomiting, and diarrhea. Rash, hyperactivity, or drowsiness may also be observed.

In case of overdosage, discontinue medication, treat symptomatically, and institute supportive measures as required. If the overdosage is very recent and there is no contraindication, an attempt at emesis or other means of removal of the drug from the stomach may be performed.

Interstitial nephritis resulting in oliguric renal failure may occur after overdosage with Amoxicillin. Renal impairment appears to be reversible with cessation of drug administration. High blood levels may occur more readily in patients with impaired renal function because of decreased renal clearance of both Amoxicillin and clavulanic acid. Both Amoxicillin and clavulanic acid may be removed from circulation by hemodialysis.

STORAGE:

Store below 30°C.

Keep medicines out of reach of children.

SHELF LIFE:

24 Months.

PRESENTATION:

Each pack contains: 10 Strips of 10 Tablets each.
2 Strips of 7 Tablets each.

Manufactured in India by:

UNICHEM LABORATORIES LTD.

Village Bhatauli Kalan, Baddi, Dist. Solan (H.P.) - 173 205.

PRODUCT REGISTRATION HOLDER:

HEALOL PHARMACEUTICALS Sdn Bhd.
(Reg.No. 298878-K)
74-3, Jalan Wangsa Delima 6,
KLSC Wangsa Maju,
53300 Kuala Lumpur, Malaysia

13016596

Date of Publication of Insert : October, 2025

140 mm

140 mm

Company Name : UNICHEM LABORATORIES LTD.		Date: 03-10-2025
Agency Name: Unicorn Creation	Pharmacode: 298	Open Size: 140 x 220 mm
Product: Myclav-625 - Leaflet	Packing: -	Pre-folded size: 70 x 55 mm
Item Code: 13016596	Location: Baddi I	No. of folds: 2 Horizontal & 1 Vertical
Specification: 54 gsm Maplitho Paper (Front back printing)		Market: Malaysia with (30°C Storage)
Old Item Code: 13008352		Text: Malaysia specific
Reason: Pharmacode added and text alignment modified.		