

REMACYCLINE L.A. 20%w/v Solution For Injection

Composition: Oxytetracycline 20g/100ml. Sodium formaldehyde sulphoxylate (as preservative) 0.5g/100ml. Long acting excipient.

Product Description: Clear yellow solution in glass bottle.

Pharmacodynamics : Inside the cell, oxytetracycline binds reversibly to receptors on the 30S sub-unit of the bacterial ribosome where it interferes with the binding of the aminoacyl-transfer RNA to the acceptor site on the messenger RNA ribosome complex. This effectively prevents the addition of amino acids to the elongating peptide chain, inhibiting protein synthesis and leads consequently to a blockade of the growth of bacteria. Oxytetracycline has mainly a bacteriostatic activity. This bacteriostatic activity involves a penetration of the substance in the bacterial cell. Oxytetracycline enters into the cell by both passive and active diffusion. The principal mode of resistance to oxytetracycline is linked with the eventual presence of a R factor responsible for a decrease of the active transport of oxytetracycline. Oxytetracycline is a broad-spectrum antibiotic. It is mainly active against Gram-negative and Gram-positive organisms, aerobian or anaerobian, and against mycoplasma, Chlramydiae and Riscckettiae.

Acquired resistance to oxytetracycline has been noted. Such resistance is usually plasmid mediated. Cross-resistance to other tetracyclines occurs. Continuous treatment with low doses of oxytetracycline can also result in increased resistance to other antibiotics.

Pharmacokinetics : After administration, oxytetracycline is rapidly absorbed and it is well distributed throughout the body, with highest concentrations found in liver, spleen, kidney and lung. Oxytetracycline passes through the placental barrier. The excipient of the veterinary medicinal product as well as the oxytetracycline form allow a plasma oxytetracycline concentration more than 0.5 µg/ml during about 72 hours, following an intramuscular injection at the dose of 20 mg/kg. Oxytetracycline binds to plasma proteins differently according to species (20-40%). Oxytetracycline is excreted under unchanged form, mainly by urinary route. It is also eliminated by biliary route but an important part of oxytetracycline is reabsorbed by small intestine (enterohepatic cycle).

Indication : In cattle and swine: Treatment of septicaemia, respiratory diseases, digestive or genito-urinary diseases, foot rot due to organisms sensitive to oxytetracycline. In Sheep: for the control of enzootic abortion and pneumonia caused by oxytetracycline-sensitive organisms. The product may be an aid in the treatment of foot rot, acute severe mastitis, infectious ovine keratonconjunctivitis (pink-eye).

Routes of administration: Intramuscular route in cattle, swine and sheep. Sub-cutaneous route in piglets.

Dosage: 20 mg of oxytetracycline per kg of live weight in one single injection, i.e. 1 ml of solution for injections per 10 kg of live weight. When clinical signs are persistent, a second injection of 20 mg of oxytetracycline per kg can be given 72 hours after the first administration.

Contraindications: Do not use in case of intolerance to oxytetracycline or to any other substance belonging to the tetracyclines group. Do not use in cases of known resistance to tetracyclines.

Warnings and Precautions:
Special warnings for each target species: None.
Special precautions for use in animals: None.
Special precautions to be taken by the person administering the veterinary medicinal product to animals: People with known hypersensitivity to tetracyclines should avoid contact with this product. In the case of reaction after contact with the product (cutaneous eruption for example), consult a doctor.
Other precautions: None.
Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products: Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Interactions with other medicaments: Oxytetracycline should not be administered simultaneously with penicillins and cephalosporins. Divalent or trivalent cations (Mg, Fe, Al, Ca) can chelate tetracyclines.

Statement on usage during pregnancy and lactation: Oxytetracycline showed no evidence of embryotoxicity or teratogenicity in laboratory animals. In mammals, oxytetracycline passes through the placental barrier, leading to a coloration of the teeth and foetal growth retardation. Tetracyclines are found in maternal milk. The safety of the product was not evaluated in pregnant cows. The use of the product in pregnant cows should be the subject of a risk/benefit assessment report issued by the veterinary.

Adverse Effects: Intolerance reactions (from injection site pain to lesions of muscular necrosis) can be observed locally following intramuscular or sub-cutaneous administration. For all the tetracyclines, general adverse reactions have been reported as gastro-intestinal troubles, and, less often, hypersensitivity reactions and photosensitivity.

Overdose and Treatment: Intolerance reactions (from injection site pain to lesions of muscular necrosis) can be observed locally following intramuscular or sub-cutaneous administration. For all the tetracyclines, general adverse reactions have been reported as gastro-intestinal troubles, and, less often, hypersensitivity reactions and photosensitivity.

Incompatibilities : None known

Storage Conditions: Store below 25 °C, protect from light.

Withdrawal Period: Meat and offal: 21 days. Milk: 7 days.

Packing: 50ml, 100ml, 250ml, 500ml in glass bottles.

Manufactured by :

DOPHARMA FRANCE, 23 rue du Prieuré, Saint Herblon, 44150 Vair sur Loire, France

Registration holder and repacker:

RHONE MA MALAYSIA SDN BHD (200001023279)
Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor
Tel: 03-7873 7355 Fax: 03-2770 0119
E-mail: customercareline@rhonema.com

Date of revision: January 2020
5LREMAC3-100 03/XXX

W10cm H14cm

LABEL CODE: 5LREMAC3-100 PRODUCT: REMACYCLINE LA PACK SIZE: 100 ML COUNTRY: MY DIMENSIONS: cm = 10 x 14	VERSION Version: 3 MMM/YY: XXX	COLORS Black ⚠ Please use the official PANTONE® (P) matching ● system for an accurate color representation.
Marketing approval: COMPLETE NAME: DATE + SIGNATURE:	Regulatory approval: COMPLETE NAME: DATE + SIGNATURE:	GMP approval: COMPLETE NAME: DATE + SIGNATURE: