

Clinacin 25 mg, 75 mg and 150 mg
Tablets For Dogs



Composition:

Active Ingredient

Clinacin 25mg tablets contain 25mg Clindamycin (as Clindamycin Hydrochloride) 100 tablets for oral administration
Clinacin 75mg tablets contain 75mg Clindamycin (as Clindamycin Hydrochloride) 50 or 100 tablets for oral administration
Clinacin 150mg tablets contain 150mg Clindamycin (as Clindamycin Hydrochloride) 50 or 100 tablets for oral administration
Each tablet also contains the following inactive ingredients: Ludipress (Lactose monohydrate, Povidone and Crospovidone), Microcrystalline Cellulose, Sodium lauryl sulphate, Colloidal silicon dioxide, Magnesium stearate.

Product Description:

Clinacin 25mg: A small round white tablet, with a break-line on one side and the letter J embossed on both side of the breakline
Clinacin 75mg: White circular tablets with a breakline on one side
Clinacin 150mg: An oblong white tablet with a breakline on one side.

Pharmacodynamics: Clindamycin is a chlorinated analogue of lincomycin. The antibiotic activity of clindamycin is based on the inhibition of bacterial synthesis. Reversible coupling to the 50s subunit of the bacterial ribosome inhibits inter alia the translation of tRNA-bound amino acids, thereby preventing elongation of the peptide chain. Because of this, the mode of action of clindamycin is predominantly bacteriostatic. Clindamycin has been shown to have in-vitro activity against the following organisms: *Staphylococcus spp*; *Streptococcus spp*; *Bacteroides spp*; *Fusobacterium spp*; *Clostridium spp*. Clindamycin and lincomycin show cross-resistance, which is common also to erythromycin and other macrolid-antibiotics. Acquired resistance can occur, by methylation of the ribosomal binding site via chromosomal mutation in gram positive organisms, or by plasmid-mediated mechanisms in gram negative organisms.

Pharmacokinetics: Clindamycin is almost completely absorbed after oral administration. Peak serum concentrations are attained approximately 1 hour after administration at a dose rate of 10 mg/kg, Cmax 3.3 µg/ml (non-fasting) – 5.0 µg/ml (fasting). Clindamycin penetrates well and may concentrate in some tissues. The t ½ of clindamycin is approximately 4 hours. Approximately 70% clindamycin is excreted in the faeces and approximately 30% in the urine.
Indication: Clinacin tablets are antibiotics indicated for the treatment of infected wounds, abscesses and oral cavity/dental infections caused by or associated with clindamycin-sensitive species of; *Staphylococcus spp*, *Streptococcus spp*, *Bacteroides spp*, *Fusobacterium necrophorum*, *Clostridium perfringens* and osteomyelitis caused by *Staphylococcus aureus*.

Routes of administration: Oral

Dosage:

Infected wounds, abscesses and oral cavity/dental infections:

5.5mg/kg clindamycin every 12 hours for 7-10 days (ie 1 tablet per 4.5kg bodyweight twice daily). If no improvement is seen within 4 days the sensitivity of the pathogen involved should be redetermined.

Dental and periodontal infections: - in the case of dental/surgical treatment due to dental infection, treatment may be started before the dental/surgical treatment.
Osteomyelitis

11mg/kg clindamycin every 12 hours for at least 4 weeks (ie 2 tablets per 4.5kg bodyweight twice daily). If no improvement is seen within 14 days the sensitivity of the pathogens involved should be redetermined.

Treatment with Clinacin should be based on susceptibility testing.

Contraindications: Do not use in cases of known hypersensitivity to clindamycin or

lincomycin. Do not administer to rabbits, hamsters, guinea pigs, chinchillas, horses or ruminants because ingestion of clindamycin by these species may result in severe gastro-intestinal disturbance.

Warnings and Precautions:

Special precautions for use

During prolonged therapy of one month or greater, periodic liver and kidney function tests and blood counts should be performed. Patients with severe renal and/or hepatic disturbances accompanied by severe metabolic aberrations should be dosed with caution and should be monitored by serum examination during clindamycin therapy.

Special Warnings for each target species

Not known

Special Precautions to be taken by the person administering the product to animals

Wash hands after the administration of the product. Persons with known hypersensitivity to lincosamides (lincomycin, clindamycin) should not handle the product.

Keep out of reach of children

Special precautions for the disposal of unused product or waste material, if any

Any unused product or waste material should be disposed of in accordance with national requirements.

Interactions with other medicaments: Neuromuscular blocking effects have been observed with clindamycin possibly leading to an increase of efficacy of other neuromuscular blocking agents. The simultaneous use of such drugs must be handled with care. Clindamycin should not be used simultaneously with chloramphenicol or macrolides because their action site is also the 50s subunit and antagonistic effects can possibly occur. When clindamycin and aminoglycoside antibiotics (eg gentamycin) are used simultaneously adverse interactions (acute renal failure) cannot be fully excluded.

Statement on usage during pregnancy and lactation: While high dose studies in rats suggest that clindamycin is not teratogen and does not significantly affect the breeding performance of males and females, safety during pregnancy and lactation or in breeding male dogs has not been established.

Adverse Effects: Vomiting and diarrhea are observed occasionally. Clindamycin sometimes causes the overgrowth of non sensitive organisms such as resistant *Clostridia* and yeasts. In cases of superinfection, appropriate measures should be taken according to the clinical situation.

Overdose and Treatment: Doses of 300mg/kg have been tolerated by dogs with no adverse reactions. Occasional vomiting, inappetence, diarrhoea, leukocytosis and increases in liver enzymes (AST, ALT) have been observed. In such cases, treatment should be stopped immediately and the animals treated symptomatically.

Storage Conditions: Store below 25° C.

Withdrawal Period: N/A

Packing: 100 tablets per bottle.

FOR VETERINARY USE ONLY

Manufactured by :



Chanelle Pharmaceuticals Manufacturing Ltd.,
Loughrea, Co. Galway, Ireland.

Registration holder and repacker:



RHONA MA MALAYSIA SDN BHD (200001023279)
Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor, Malaysia
Tel: 03-7873 7355 Fax: 03-2770 0119
E-mail: customercareline@rhonema.com

Date of revision: XXX
5LCL0253-100 03/XXX

W10cm H14cm

LABEL CODE:	VERSION	COLORS
5LCL0253-100		Black
PRODUCT: CLINACIN 25MG TABLETS FOR DOGS	Version: 3	
PACK SIZE: 100 TABLETS	MMM/YY: XXX	
COUNTRY: MY		
DIMENSIONS: cm = 10 x 14		⚠ Please use the official PANTONE® (P) matching system for an accurate color representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME:	COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE:	DATE + SIGNATURE: