

Important Information. Please read carefully.

Evofen

Suspension 100mg/5ml



COMPOSITION

Each 5ml contains Ibuprofen 100mg

PHARMACODYNAMIC

Ibuprofen is a phenylpropionic acid derivative which has analgesic, antipyretic and anti-inflammatory actions.

PHARMACOKINETIC

Ibuprofen is absorbed from the gastrointestinal tract and peak plasma concentration occurs about 1 to 2 hours after ingestion. It is extensively bound to plasma proteins and is rapidly excreted in the urine, mainly as metabolites and their conjugates. Approximately 1% is excreted in the urine as unchanged ibuprofen and about 14% as conjugated ibuprofen.

INDICATIONS

Evofen is indicated for the relief of fever such as fever associated with cold and flu, and pain such as pain from teething and toothache, earache, sore throats, headache and minor aches and sprains. Also indicated for the relief of inflammation of muscles and joints.

DOSAGE AND ADMINISTRATION

The recommended daily dose, unless otherwise prescribed by the physician is given below. Preferably to be taken with food or milk.

The daily dosage of Evofen Suspension is 20-30mg per kilogram body-weight in divided doses.

The doses should be given approximately every 6-8 hours (or with a minimum of 4 hours between each dose if required).

Recommended Daily Dose		Daily Frequency
Adults (Maximum daily dose: 2400mg Ibuprofen in divided doses)		
Adults	2 teaspoons (10ml) or 4 teaspoons (20ml) (equivalent to 200mg or 400mg of Ibuprofen)	4 times
Children (Maximum daily dose: 30mg Ibuprofen/kg body weight in divided doses) Children weighing less than 30kg should not take more than 500mg Ibuprofen within a day. Not suitable for children under 3 months of age unless advised by your doctor.		
10 - 12 years old	3 teaspoons (15ml) (equivalent to 300mg of Ibuprofen)	3 times
7 - 9 years old	2 teaspoons (10ml) (equivalent to 200mg of Ibuprofen)	3 times
4 - 6 years old	1½ teaspoons (7.5ml) (equivalent to 150mg of Ibuprofen)	3 times
1 - 3 years old	1 teaspoon (5ml) (equivalent to 100mg of Ibuprofen)	3 times
6 months - 1 year old	½ teaspoon (2.5ml) (equivalent to 50mg of Ibuprofen)	3 - 4 times
3 - 6 months (weighing more than 5kg)	½ teaspoon (2.5ml) (equivalent to 50mg of Ibuprofen)	3 times

Post-immunisation

May be given to baby under 6 months old, on the advice of a doctor. Initial dose of 2.5ml followed by a further 2.5ml 6 hours later if necessary. Not more than 5.0ml in 24 hours.

CONTRAINDICATIONS

Evofen should not be given to patients who are sensitive to aspirin, ibuprofen or other non-steroidal anti-inflammatory drugs (NSAIDs). It should not be given to patients in whom aspirin and other NSAIDs induce the symptoms of asthma, rhinitis or urticaria. It should not be given to patients with active or history of peptic ulceration.

WARNINGS AND PRECAUTIONS

Use with caution in gastric ulceration, allergic disorders particularly in asthma and hypersensitivity to salicylates (eg. aspirin) and renal (kidney) and hepatic (liver) impairment.

It is best avoided in pregnancy as its safety for such use has not been absolutely established.

WARNING**Risk of GI ulceration, bleeding and perforation with NSAID**

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms, in patients treated with NSAID therapy. Although minor upper GI problems (e.g. dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract symptoms.

Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious GI events and other risk factors associated with peptic ulcer disease (e.g. alcoholism, smoking and corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events.

Measures such as the use of physical therapy and mild analgesics like paracetamol (when inflammation is not a major factor) should be instituted prior to initiation of therapy with NSAID. NSAIDs should only be used after proper appraisal of potential risks to patients. It should be used with lowest effective dose for only as long as needed.

This drug should not be co-administered with other NSAIDs. Prescribers should inform patients about the signs and/or symptoms of serious GI toxicity and what steps to take if they occur. In considering the use of relatively large doses (within the recommended dosage range), sufficient benefit should offset the potential increased risk of GI toxicity.

DRUG INTERACTIONS

Ibuprofen may reduce the antihypotensive effect of antihypertensives. It may potentiate the effect of lithium and digoxin when given concomitantly. Diuretics may increase the risk of nephrotoxicity of NSAIDs.

SIDE EFFECTS

Gastrointestinal discomfort and occasional gastro-intestinal bleeding and nausea may occur. These side effects may be reduced if the medication is taken with food or milk.

Other side effects may include hypersensitivity reactions such as asthma and rashes; headache; vertigo; tinnitus and in rare cases fluid retention.

PREGNANCY AND LACTATION**USE DURING PREGNANCY**

Whilst no teratogenic effects have been demonstrated in animal experiments, the use of ibuprofen during pregnancy should, if possible, be avoided.

USE DURING LACTATION

Ibuprofen appears in the breast milk in very low concentration, and is unlikely to affect the breast-fed infant adversely.

OVERDOSAGE AND TREATMENT

There is no specific antidote to overdose with ibuprofen. Gastric lavage may be performed and if necessary blood electrolytes corrected.

SHELF-LIFE

The expiry date is indicated on the packaging.

PRESENTATION

Orange colour suspension with orange flavour.

DOSAGE FORM AND PACKAGING

Oral Suspension. Available in pack of 90ml.

STORAGE

Store below 30°C. Protect from light.

FOR ORAL ADMINISTRATION

SHAKE WELL BEFORE USE

KEEP OUT OF THE REACH OF CHILDREN/JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date : 03-July-2017

Manufacturer

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia

Distributor

Apex Pharmacy Marketing Sdn Bhd
No. 2, Jalan SS13/5, 47500 Subang Jaya, Selangor, Malaysia