

ERYTHRON 300MG Water Soluble Powder

FOR ANIMAL USE ONLY

Brand or Product Name :

ERYTHRON 300MG Water Soluble Powder

Name and Strength of Active Substance

Each gram of ERYTHRON 300MG water soluble powder contains Erythromycin Thiocyanate 368mg which is equivalent to Erythromycin 300mg/g (potency)

Product Description

Soluble fine white powder with characteristic odour and bitter taste.

Pharmacodynamics & Pharmacokinetics

The main effect of erythromycin thiocyanate comes from erythromycin. Erythromycin is usually a bacteriostatic agent but in high concentration or against highly susceptible organisms it may be bactericidal. Erythromycin is believed to act by binding to the 50S ribosomal subunit of susceptible bacteria thereby inhibiting peptide bond formation.

Antimicrobial activity : Erythromycin has activity against gram-positive cocci (staphylococci, streptococci), gram-positive bacilli (Bacillus anthracis, Corynebacterium, Clostridium sp., (not C. difficile), Listeria, Erysipelothrix), some strains of Gram-negative bacilli, including Haemophilus, Pasteurella and Brucella. Some strains of Actinomyces, Mycoplasma, Chlamydia, Ureaplasma and Rickettsia are also inhibited by erythromycin. **Absorption :** Erythromycin is absorbed after oral administration in the upper small intestine.

Distribution : Erythromycin is distributed throughout the body into most fluids and tissues including the prostate, macrophages and leukocytes. CSF levels are poor. Erythromycin will cross the placenta and levels of 5 – 20% of those found in the mother's serum can be found in the fetal circulation. Erythromycin levels of about 50% of those found in the serum can be detected in milk. The volume of distribution in dogs is reportedly 2L/kg, 3.7-7.2L/kg in foals, 2.3L/kg in mares, and 0.8L/kg in cattle. **Protein Binding :** Erythromycin may be 73 – 81% bound to serum proteins. **Metabolism and excretion :** Erythromycin is primarily excreted unchanged in the bile, but is also partly metabolized by the liver via N-demethylation to inactive metabolites. Some of the drug is reabsorbed after biliary excretion. Only about 2-5% of a dose is excreted unchanged in the urine. The reported elimination half-life of erythromycin in various species are 60-90 minutes in dogs and cats, 60-70 minutes in foals and mares, and 190 minutes in cattle.

Indication

For the treatment of Chronic Respiratory Disease (CRD) in chickens caused by Mycoplasma organisms.

Recommended Dosage

Poultry :

100 g ERYTHRON 300MG per 116 litres of drinking water for 1 to 5 consecutive days, depending on response (approximately 25.5mg erythromycin / kg bodyweight/day).

150 g ERYTHRON 300MG per 175 litres of drinking water for 1 to 5 consecutive days, depending on response (approximately 25.5mg erythromycin / kg bodyweight/day).

500 g ERYTHRON 300MG per 582 litres of drinking water for 1 to 5 consecutive days, depending on response (approximately 25.5mg erythromycin / kg bodyweight/day).

Mode of Administration

To be administered via animal drinking water.

Mixing Directions :

To assure thorough dissolution, place the ERYTHRON 300MG powder in a small mixing container and add 25 litres of water. Stir until all the powder completely dissolved. Mix this concentrated stock solution with water to make the final desired volume of treated animal drinking water. Prepare fresh medicated solution daily.

Contraindications

Any medicated drinking water which is not consumed within 24 hours should be discarded.

Warnings and Precautions

Due to a lack of information on the fate and effects of erythromycin in the environment, veterinary surgeons prescribing this product under the cascade should satisfy themselves that there will be no adverse impact in the environment. Do not eat, drink or smoke while preparing the diluted formulation or while animals are being treated. Wash hands and exposed skin after use. The water system to which the product is added must be in good working order and the header tanks and troughs must be free of dust, algae and other particulate matter.

Interactions with Other Medicaments

None reported.

Statement on usage during pregnancy and lactation

Not applicable.

Adverse Effects / Undesirable Effects

None reported.

Overdose and Treatment

No information available.

Withdrawal Period(s)

Chickens : slaughter 3 days , eggs 6 days.

Shelf-life

3 years from the date of production.

Maximum Residual Limit (MRL)

Milk (mammalian) is 40 mcg/kg, Edible offal, muscle, egg (mammalian and poultry) is 300 mcg/kg. [Ref. REGOVP, Version 2, dated Aug 2008]

Storage Conditions

Store in a dry place below 25°C. Keep out of reach of children. Any unused sachet or medicated water to be discarded after 24 hours.

Dosage Forms and packaging available

Available in 100g, 150g, 500g and 1000g powder per sachet which are packed in 10, 20, 50 and 100 sachets per box.

Name and Address of manufacturing

Life Biopharma Sdn. Bhd. No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2010-03-18