

AMOXICILLIN

SOLUBLE POWDER 500mg/g

Ingredient(s):

Each g contains:

Amoxicillin (as Amoxicillin Trihydrate) 500mg

Preservative(s):

Each g contains :

Sodium Benzoate 1.0mg

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

Amoxicillin is an aminopenicillin. It is bactericidal against susceptible bacteria and act by inhibiting mucopeptide synthesis in the cell wall resulting in a defective barrier and an osmotically unstable spheroplast. Beta-lactam antibiotics have been shown to bind to several enzymes (carboxypeptidases, transpeptidases, endopeptidases) within the bacterial cytoplasmic membrane that are involved with cell wall synthesis. The different affinities that various beta-lactam antibiotics have for these enzymes (also known as penicillin-binding proteins; PBPs) help explain the differences in spectrums of activity the drugs have that are not explained by the influence of beta-lactamases. Like other beta-lactam antibiotics, penicillins are generally considered to be more effective against actively growing bacteria. The aminopenicillins, have increased activity against many strains of G(-) aerobes not covered by either the natural penicillins or penicillinase-resistant penicillins.

Amoxicillin trihydrate is relatively stable in the presence of gastric acid. After oral administration, it is about 74~92% absorbed in humans and animals (monogastric). Food will decrease the rate, but not the extent of oral absorption. Amoxicillin serum levels will generally be 1.5~3 times greater than those of ampicillin after equivalent oral doses.

After absorption the volume of distribution for amoxicillin is approximately 0.3L/kg in humans and 0.2L/kg in dogs. The drug is widely distributed to many tissues, including liver, lungs, prostate (human), muscle, bile, and ascitic, pleural and synovial fluids. Amoxicillin will cross into the CSF when meninges are inflamed in concentrations that may range from 10~60% of those found in serum. Very low levels of the drug are found in the aqueous humor, and low levels found in tears, sweat and saliva. Amoxicillin crosses the placenta, but it is thought to be relatively safe to use during pregnancy. It is approximately 17~20% bound to human plasma proteins, primarily albumin. Protein binding in dogs is approximately 13%. Milk levels of amoxicillin are considered to be low.

Amoxicillin is eliminated primarily through renal mechanisms, principally by tubular secretion, but some of the drug is metabolized by hydrolysis to penicilloic acids (inactive) and then excreted in the urine. Elimination half-lives of amoxicillin have been reported as 45~90 minutes in dogs and cats, and 90 minutes in cattle. Clearance is reportedly 1.9ml/kg/min in dogs. Hepatic inactivation and biliary secretion is a minor route of excretion.

Indication(s):

For the treatment of infections caused by bacteria susceptible to amoxicillin, including:

Chicken:

- i Colibacillosis, peritonitis and respiratory tract infections caused by non-β-lactamase-producing *E. coli*
- ii Salmonellosis caused by non-β-lactamase-producing *Salmonella enterica* subsp. *Enteric*

Swine:

- i Respiratory infections caused by *Actinobacillus pleuropneumoniae*
- ii Arthritis and meningitis caused by *S. suis*
- iii Secondary infections due to ear and tail-biting

Target Species

Chicken and swine.

Dosage and Administration:

Chicken:

Recommended dosage: 14~28mg per kg body weight per day, for a period of 3~5 consecutive days.

Birds 0~4 weeks of age: Dissolve 8~17g into every 100L of drinking water.

Birds > 4 weeks of age: Dissolve 14~28g into every 100L of drinking water.

Swine:

Recommended dosage: 28mg per kg body weight per day, for a period of 3~5 consecutive days.

Pigs < 4 months of age: Dissolve 28g into every 100L of drinking water.

Pigs > 4 months of age: Dissolve 42g into every 100L of drinking water.

Orally via drinking water.

Contraindications:

1. Penicillins are contraindicated in patients who have a history of hypersensitivity to them. Because there may be cross-reactivity, use penicillins cautiously in patients who are documented hypersensitive to other beta-lactam antibiotics (e.g., cephalosporins, cefamycins, carbapenems).
2. Do not use in rabbits, hamsters, gerbils and guinea pigs.

Precaution(s) / Warning(s):

Do not administer systemic antibiotics orally in patients with septicemia, shock, or other grave illnesses as absorption of the medication from GI tract may be significantly delayed or diminished.

Special precautions for use in animals:

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Special precaution for be taken by the person administering the veterinary medicinal product to animals:

Avoid inhalation of dust. Wash hands after use.

L
260mmW
110mm

243011 - 02

260 x 110

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

- 1) Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.
- 2) Handle this product with great care to avoid exposure, taking all recommended precautions.
- 3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Interaction with Other Medicaments

- 1) In vitro studies have demonstrated that penicillins can have synergistic or additive activity against certain bacteria when used with aminoglycosides or cephalosporins.
- 2) Use of bacteriostatic antibiotics (e.g., chloramphenicol, erythromycin, tetracyclines) with penicillins is generally not recommended, particularly in acute infections where the organism is proliferating rapidly as penicillins tend to perform better on actively growing bacteria.
- 3) Probenecid competitively blocks the tubular secretion of most penicillins, thereby increasing serum levels and serum half-lives.

Pregnancy and Lactation

Penicillins have been shown to cross the placenta and safe use of them during pregnancy has not been firmly established, but neither have there been any documented teratogenic problems associated with these drugs. However, use only when the benefits outweigh the risks.

Side Effect(s) / Adverse Reaction(s):

- 1) Hypersensitivity reactions unrelated to dose can occur with these agents and can be manifested as rashes, fever, eosinophilia, neutropenia, agranulocytosis, thrombocytopenia, leukopenia, anemias, lymphadenopathy, or full blown anaphylaxis. In humans, it is estimated that up to 15% of patients hypersensitive to cephalosporins will also be hypersensitive to penicillins. The incidence of cross-reactivity in veterinary patients is unknown.
- 2) When given orally, penicillins may cause GI effects (anorexia, vomiting, diarrhea). Because the penicillins may also alter gut flora, antibiotic-associated diarrhea can occur, as well as selecting out resistant bacteria maintaining residence in the colon of the animal (superinfections).
- 3) High doses or very prolonged use has been associated with neurotoxicity (e.g., ataxia in dogs).
- 4) Although the penicillins are not considered to be hepatotoxic, elevated liver enzymes have been reported.
- 5) Other effects reported in dogs include tachypnea, dyspnea, edema and tachycardia.

Environmental Property

None has been mentioned in animal.

Symptoms and Treatment for Overdosage, and Antidote(s):

Acute oral penicillin overdoses are unlikely to cause significant problems other than GI distress, but other effects are possible.

Shelf-Life:

3 years.

Use within 24 hours after reconstitution.

Storage Condition(s):

Store at temperature below 30°C.

Product Description and Packing(s):

A white to pale yellow color powder, and without other foreign matter.
Al. Pouch 1kg and 100g.

For food producing animals product:

Withdrawal Periods

Broiler poultry: 2 days before slaughter.

Swine: 3 days before slaughter.

Egg: Do not use in birds producing eggs for human consumption.

Maximum Residual Limit (MRL):

Swine and chicken edible tissues: 0.01mg/kg.



Made by YSP

Manufacturer:

Y.S.P INDUSTRIES VIETNAM CO., LTD

Road No.6, Nhon Trach 3 Industrial zone, Nhon Trach,
Dong Nai province, Vietnam Tel: +8461 3569 993

Product Registration Holder :

Y.S.P INDUSTRIES (M) SDN. BHD.

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 06 Feb 2018

To update date of revision.