

SULTRIM POWDER

Ingredient(s):

Each g contains :

Sulfadiazine	400mg
Trimethoprim	80mg

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

Sulfadiazine is a bacteriostatic antibiotic belonging to the sulphonamide group which acts by interference with the synthesis of nucleic acids. Trimethoprim is a dihydrofolate reductase inhibitor which also interferes with the synthesis of bacterial nucleic acids. Sulfadiazine and trimethoprim act on the same metabolic pathway, resulting in potentiation of antibacterial activity.

Following oral administration of sulfadiazine and trimethoprim to chickens, $t_{1/2\alpha}$ and $t_{1/2\beta}$ values of 0.756 and 7.07 hours (sulfadiazine) and 0.680 and 6.24 hours (trimethoprim) were obtained. Values for T_{max} were 2.46 and 2.44 hours, values for C_{max} were 86.45 and 3.65mcg/ml and values of AUC were 620.50 and 19.87 mcg.hour/ml, respectively, for sulfadiazine and trimethoprim.

Indication(s):

Chickens and turkeys: For use in the treatment of diseases caused by bacteria sensitive to potentiated sulphonamides including infections due to *Salmonella* infection and pasteurellosis.

Pigs: For the treatment of atrophic rhinitis when associated with *Bordetella bronchiseptica* and streptococcal meningitis caused by *Streptococcus suis* type II.

Target Species

Swine, non-laying chickens and turkeys.

Dosage and Administration:

To ensure thorough dispersion, the product should first be mixed with a small quantity of feed before incorporation in the final mix.

Swine: Incorporate **0.625kg** of the product per tonne of feed to provide 300 mg/kg of combined actives. The recommended dose is 15 mg combined actives/kg bodyweight. This is equivalent to **30mg** of the product/kg bodyweight daily. Feed for 5 consecutive days.

Chickens: Incorporate **0.625kg** of the product per tonne of feed to provide 300 mg/kg combined actives. The recommended dose is 30 mg combined actives/kg bodyweight daily. This is equivalent to **62mg** of the product/kg bodyweight daily. For example, this dosage will be achieved in chickens of approximately 28 days of age (1.1 kg bodyweight) by the addition of **0.625kg** Sultrim Powder per tonne of feed to provide 300 mg/kg of combined actives. Feed for 5 to 10 consecutive days.

Turkeys: The recommended dose is 15 mg combined actives/kg bodyweight daily. Feed for 10 consecutive days.

Contraindications:

- 1) Hypersensitivity to diaminopyrimidines or sulfonamides: Animals that have had a previous reaction may be much more likely to react on subsequent administration.
- 2) Risk-benefit should be considered when the following medical problems exist:
 - a. Hepatic function impairment: Delayed biotransformation may increase the risk of adverse effects.
 - b. Renal function impairment: Delayed elimination could cause accumulation of sulfonamide and metabolites, increasing the risk of adverse effects.

Precaution(s) / Warning(s):

- 1) Trimethoprim/sulfadiazine should not be used in dogs or horses showing marked liver parenchymal damage, blood dyscrasias, or in those with a history of sulfonamide sensitivity.
- 2) An idiosyncratic sulfonamide toxicosis can occur in any breed of dog, but the reaction has been reported more frequently in the Doberman Pinscher than in other breeds. This specific type of drug reaction includes blood dyscrasias, nonseptic polyarthritis, and skin rash.
- 3) To avoid the possibility of crystalluria, adequate water intake is essential.
- 4) Special precautions to be taken by the person administering the veterinary medicinal product to animals. Incorporation into the feed must be performed by a suitably approved manufacturer. Persons handling this product should avoid inhalation of any dust and contact with skin. Wear either a disposable half-mask respirator or a non-disposable respirator with filter when mixing or handling this product. Rubber gloves should be worn when mixing or handling this product. Hands should be washed thoroughly after use. Sulfonamides may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to sulphonamides may lead to cross reactions with other antibiotics. Allergic reactions to these substances may occasionally be serious.
 - i) Do not handle this product if you know you are sensitive to sulphonamides.
 - ii) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning.

Interaction with Other Medicaments

- 1) Trimethoprim/ sulfadiazine may prolong the clotting times in patients receiving coumarin (warfarin) anticoagulants.
- 2) Sulfonamides may displace other highly bound drugs, such as methotrexate, phenylbutazone, thiazide diuretics, salicylates, probenecid and phenytoin.
- 3) Antacids may decrease the bioavailability of sulfonamides if administered concurrently.
- 4) Trimethoprim may decrease the therapeutic effect of cyclosporine (systemic) and increase the risk of nephrotoxicity developing.
- 5) A trimethoprim and sulfonamide combination administered to a detomidine-anesthetized horse can lead to arrhythmias, hypotension, and death; it is suspected that the antimicrobial potentiates the cardiac changes reported with detomidine.

Pregnancy and Lactation

Data on the exact level below which no effects on foetal development were observed are not available. However extensive use of the product in different species over many years, have not shown adverse effects on the foetus.

When administered to lactating females, small amounts of trimethoprim and sulfadiazine are present in the maternal milk. Since no studies have been reported of the effect on the development of new born young of the ingestion of this milk, it would be prudent not to feed very young animals with milk obtained from the mother.

Do not feed to laying poultry.

Side Effect(s) / Adverse Reaction(s):

Hypersensitivity to sulphonamides has been rarely reported. However this remains a potential undesirable effect. Crystalluria caused by precipitation of insoluble sulphonamide is a theoretical risk although rarely reported.

Environmental Property

None has been mentioned in animal.

Symptoms and Treatment for Overdosage, and Antidote(s):

No information available. As there is no specific antidote, treatment should be symptomatic.

Shelf-Life:

4 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C.

Product Description and Packing(s):

A white to pale-yellow color powder, and without other foreign matter.

Paper bag of 20kg

For food producing animals product:

Withdrawal Periods

Animals and birds must not be slaughtered for human consumption during treatment.

Pigs may be slaughtered for human consumption only after 10 days from the last treatment.

Chickens may be slaughtered for human consumption only after 5 days from the last treatment

and turkeys only after 2 days from the last treatment.

Not to be used in chickens or turkeys producing eggs for human consumption

Maximum Residual Limit (MRL):

Trimethoprim (mg/kg):

Cattle milk	0.05
Edible offal (mammalian)	0.05
Eggs	T*0.02
Meat [mammalian]	0.05
Poultry, Edible offal of	0.05
Poultry meat	0.05

Sulfadiazine (mg/kg):

Cattle milk	0.1
Edible offal (Mammalian)	0.1
Eggs	T*0.02
Meat [mammalian]	0.1
Poultry, Edible offal of	0.1
Poultry meat	0.1



Manufacturer : **Y.S.P. INDUSTRIES (M) SDN. BHD.**

Factory : Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.
Tel: 03-89251215 (4 Lines) Fax: 03-89251298

Office : No. 18, Jalan Wan Kadir, Taman Tun Dr. Ismail,
60000 Kuala Lumpur, Malaysia.
Tel: 03-77276390 (12 Lines)
Ordering Line: 1800 88 3027
Product Info: 1 800 88 3679