

TRISULMIX LIQUIDE

Solution For Oral Use

Composition: Each Trisulmix Liquide 100ml contains sodium sulphadimethoxine 20.00g (equivalent to sulphadimethoxine 18.67g); trimethoprim 4.00g, methyl parahydroxybenzoate 0.10g and propyl parahydroxybenzoate 0.01g.

Product Description: Clear light yellow solution, with alkaline taste, miscible with water.

Pharmacodynamics: Sulphadimethoxine is a long acting sulfamide with a wide antimicrobial spectrum including anti coccidia properties. This substance is active against Gram-positive and Gram-negative bacteria and certain protozoan such as coccidia. Trimethoprim belongs to diaminopyrimidine group. It is active against streptococci and most of Gram-negative bacteria. In association, the two active substances are synergic. Sulphadimethoxine is potentiated by a diaminopyrimidine, trimethoprim. The combination of these two active substances allows a sequential blockade on the folic acid biosynthesis. These two active substances act by sequential way on the synthesis route of tetrahydrofolic acid: the sulphamide inhibits the PAB acid incorporation in folic acid and trimethoprim inhibits specifically the microbial dihydrofolate reductase. The theoretical activity spectrum includes Gram-positive organisms (*Staphylococcus*, *Listeria* ...) and Gram-negative organisms (*Echerichia coli*, *Salmonella*, *Proteus*, *Enterobacter*, *Bordetella* ...).

Pharmacokinetics: Sulphadimethoxine is considered as a sulphonamide with a long persistence in plasma rate. Sulphadimethoxine is highly protein bound. Drug diffuses widely into all tissues and organs. Trimethoprim is rapidly absorbed after oral administration. It is widely distributed in the body. The two active substances are partly metabolized by liver. Their elimination is mainly renal.

Indication: In poultry, rabbits: Infections caused by or associated with organisms sensitive to sulphadimethoxine and trimethoprim. Curative and preventive treatment of respiratory and digestive infections. Curative and preventive treatment of digestive coccidiosis.

Routes of administration: oral route: in drinking water

Dosage: 18.68 to 37.36 mg of sulfadimethoxine and 4 to 8 mg of trimethoprim per kg of live weight and per day, during 5 consecutive days, by oral route, i.e. 1 to 2 ml of TRISULMIX LIQUIDE per 10 kg of live weight and per day during 5 consecutive days, in drinking water. Adjust dosage to the real consumption of the animals.

Contraindications:
Do not use in case of intolerance to sulphonamides or trimethoprim.
Do not use in animals in case of severe hepatic or renal dysfunction.

Warnings and Precautions:
Special precautions for use in animals: Water abundantly during treatment.
Special precautions to be taken by the person administering the veterinary medicinal product to animals: Handle this product with care and avoid direct contact. Take all recommended precautions by wearing protective clothing, glasses, gloves and mask. Avoid contact with skin and eyes.
In case of symptoms as cutaneous eruption, consult a doctor and show him this

information. An oedema at the level of the face, the eyelids, the lips or respiratory difficulties constitute severe symptoms and require urgent medical cares. Avoid to handle this product or the medicated water in case of allergy to trimethoprim or sulphonamides.

Other precautions: None.
Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products: Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Interactions with other medicaments: None known

Statement on usage during pregnancy and lactation: Teratogenic and foetotoxic effects have been reported in laboratory animals at doses more important than the recommended therapeutic doses.
Do not use this product during pregnancy or lactation.

Adverse Effects: No adverse events have been described.

Overdose and Treatment: None known

Storage Conditions: The product should be stored at a temperature not exceeding 25°C, protected from light.

Withdrawal Period:
Poultry: 6 days after the end of treatment
Rabbits: 8 days after the end of treatment
Eggs: As no withdrawal period for the eggs has been established, do not use in layers when eggs are delivered to human consumption (4 weeks prior and during laying).

Packing: 1L/5L HDPE bottle

FOR VETERINARY USE ONLY

Manufactured by :

DOPHARMA FRANCE, 23 rue du Prieuré, Saint Herblon, 44150 Vair sur Loire, France
Registration holder and repacker:

RHONE MA MALAYSIA SDN BHD (200001023279)
Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor
Tel: 03-7873 7355 Fax: 03-2770 0119
E-mail: customercareline@rhonema.com

Date of revision: January 2020
5LTRISU3-01L 03/XXX

W10cm H14cm

LABEL CODE:	VERSION	COLORS
5LTRISU3-01L		
PRODUCT: TRISULMIX LIQUIDE SOLUTION FOR ORAL USE	Version: 3	BLACK
PACK SIZE: 1L, 5L	MMM/YY: XXX	
COUNTRY: MY		
DIMENSIONS: cm = 10 x 14		! Please use the official PANTONE® (P) matching system for an accurate color representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME:	COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE:	DATE + SIGNATURE: