



Tucoprim® 150 (Premix)
Reg No. G2021 (Act 36/1947)

Category / Sub Category:
Sulphonamides

Active Ingredients:

Tucoprim® 150 premix contains 125g sulphadiazine and 25g trimethoprim per kg.

General Description:

- ☐ Broad spectrum anti-microbial treatment for pigs and poultry.

Indications:

- ☐ For the prevention and treatment of sulphonamide-sensitive diseases in pigs and non-laying poultry.
- ☐ **Growing Pigs:** For the treatment of streptococcal infection, atrophic rhinitis, diarrhoea and other sulphonamide sensitive diseases.
- ☐ **Sows:** For the prevention and treatment of MMA and other sulphonamide-sensitive diseases.
- ☐ **Poultry:** For the prevention and treatment of E-coli, Salmonella spp and other infections caused by sulphonamide-sensitive diseases.

Contra-indications:

- ☐ None

Directions For Use:

- ☐ Use only as directed.
- ☐ The product must be incorporated into animal feed at the recommended dose rates, mixed thoroughly and administered for the recommended period.

Pigs:

- ☐ Dosage: 15-30 mg combined activity per kg live mass. For streptococcal infection and E. coli diarrhoea use at
- ☐ 30 mg/kg.
- ☐ Guideline to recommended feed inclusion rates of Tucoprim Premix according to dosage required and feed intake of pigs.

Daily Feed Intake g/kg Pig Live Mass (= % of Body Mass)	15 mg/kg	30 mg/kg
	Inclusion rate of Tucoprim Premix per 1000 kg Feed	
20 g/kg LM (2%)	5,0 kg	10,0 kg
30 g/kg LM (3%)	3,3 kg	6,6 kg
40 g/kg LM (4%)	2,5 kg	5,0 kg
50 g/kg LM (5%)	2,0 kg	4,0 kg
60 g/kg LM (6%)	1,6 kg	3,2 kg
70 g/kg LM (7%)	1,4 kg	2,8 kg

Dosage Schedule:

☐ Growing Pigs: Feed daily for 5 days.

☐ Sows: General Infection: Feed daily for 5 days

☐ MMA: Feed daily for 3 days prior to parturition and continue for 2 days after parturition.

Poultry:

☐ Dosage: Use 25-30 mg combined activity per kg live mass. Mix 2 kg Tucoprim Premix per 1000 kg feed.

Dosage Schedule: Feed daily for 10 days.

☐ The product will remain stable when mixed with feed for 4 weeks from mixing.

Precautions:

☐ Keep out of reach of children and uninformed persons.

Warnings:

☛ For use in pigs and poultry only. Not for use in laying hens.

☛ The use of sulphonamides for prolonged periods can result in side-effects. Use only as directed.

☛ In rare cases animals may be allergic to the drug. If this occurs, stop treatment immediately and consult a veterinarian.

☛ Do not use in animals with liver damage or blood abnormalities.

☛ Treated animals must have free access to drinking water.

☛ Animals must not be slaughtered for human consumption during treatment and for the following period after treatment: Pigs: 5 days; Poultry: 24 hours (1 day).

Storage:

Store in cool, dry place.

Pack Sizes:

Boxes of 10 x 2 kg sachets.

DISCLAIMER: ALTHOUGH GREAT CARE HAS BEEN TAKEN IN THE COMPILATION OF THIS DOCUMENT, READERS ARE REQUESTED TO REFER TO THE PACKAGE INSERT FOR COMPLETE DETAILS BEFORE USING THE SPECIFIC PRODUCT.

Package Insert
Voubiljet