

PREGNANCY AND LACTATION

Pregnancy or breastfeeding patients should seek advice of doctors before taking this medication.

INTERACTION WITH OTHER MEDICATIONS

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Ask your doctor before taking pambrom if you also take any other diuretic (often used as blood pressure medication).

Pambrom is a diuretic. There are potential additive effects if combined with other diuretic drugs.

SIGNS AND SYMPTOMS OF OVERDOSE

Early symptoms of paracetamol overdose include nausea, vomiting, abdominal pain, hypotension, sweating and weakness. The most serious adverse effect of acute overdose of paracetamol is hepatotoxicity.

PACKING

Box of 20's and 180's.

SYMPTOMS AND TREATMENT OF OVERDOSE

Early symptoms (nausea, vomiting, and weakness) usually occur after ingestion of paracetamol overdose large enough to cause hepatic toxicity.

First empty stomach via induction of emesis or gastric lavage. It is recommended that acetylcysteine administration be instituted as soon as possible after ingestion of an overdose has been reported without waiting for the results of plasma paracetamol concentration or other laboratory tests. Acetylcysteine is most effective if treatment is started within 10 to 12 hours after ingestion of the overdose. However, it may be of some benefits if treatment is started within 24 hours.

For oral administration, the recommended adult dose of acetylcysteine is 140mg per kg of body weight initially, then 70mg per kg of body weight every 4 hours for 17doses. Each dose should be diluted to a 5% solution with cola, grapefruit juice, orange juice, or water prior to administration because of its unpleasant odour and its irritation and sclerosing properties. Any dose vomited may be given via duodenal intubation. Plasma acetaminophen concentration should be determined and liver function test should be performed to ascertain the status of the patient. Renal and cardiac function should be monitored and appropriate therapy instituted if necessary.

Supportive treatment consists of maintaining fluid and electrolyte balance, correcting hypoglycemia and administering vitamin K1 (if prothrombin time ratio exceeds 1.5) and fresh frozen plasma or clotting factor concentrate (if prothrombin ratio exceeds 3.0).

STORAGE CONDITIONS AND USER INSTRUCTIONS

Store in a dry place below 30°C. Protect from light.

Keep out of reach of children.

Jauh dari anak-anak.

Shelf life: Please refer to the outer package.



Product Registration Holder:
DUOPHARMA MARKETING SDN. BHD.
Lot No. 2A,6,8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate,
43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:
DUOPHARMA MANUFACTURING
(BANGI) SDN. BHD.
Lot No. 2 & 4, Jalan P/7, Section 13,
Bangi Industrial Estate,
43650 Bandar Baru Bangi,
Selangor, Malaysia.

Uphamol Menstrual

Tablet 500/25 mg



Uphamol Menstrual Tablet 500/25mg contains Paracetamol & Pambrom. Paracetamol is a known effective analgesic and antipyretic. Pambrom acts as a diuretic by increasing renal blood flow and glomerular filtration rate thus, promoting diuresis. Uphamol Menstrual Tablet 500/25mg provides symptomatic relief of water-weight gain, bloating, swelling, headache, backache, muscular aches, pains and cramps associated with premenstrual and menstrual periods.

DESCRIPTION

A light pink, oblong shape film coated tablet plain on the both side.

Each tablet contains Paracetamol500mg

Pambrom25mg

ROUTE OF ADMINISTRATION

Oral

PHARMACODYNAMICS

Analgesic: The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (CNS) and to lesser extent, through a peripheral action by blocking pain-impulse generation. The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis of action of other substances that sensitize pain receptors.

Antipyretic: Paracetamol probably produces antipyresis by acting centrally on the hypothalamic heat regulating center to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating and heat loss. The central action probably involves inhibition of prostaglandin synthesis in the hypothalamus.

Pambrom acts as a diuretic by increasing renal blood flow and glomerular filtration rate thus, promoting diuretics.

PHARMACOKINETICS

Paracetamol is rapidly and almost completely absorbed after oral administration.

Approximately 90-95% of a dose is metabolized in the liver, primarily by conjugation with glucuronic acid, sulfuric acid and cysteine. Peak concentration can be achieved within 0.5-2 hours. Paracetamol is excreted primarily as conjugates via the kidney.

DOSAGE

Menstruating females (12 years and above): Take 2 tablets every 4 to 6 hours. Do not take more than 8 tablets in 24 hours. May be taken with or without food.

Not recommended for children below 12 years.

If you miss a dose of Uphamol Menstrual Tablet 500/25mg and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is almost time for next dose, do not double the dose to catch up.

SIDE EFFECTS

When taken as directed, unwanted effects from Uphamol Menstrual are rare and usually mild. Skin rashes and other hypersensitivity reactions (eg. urticaria, erythema) have been reported occasionally.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

CONTRAINDICATION

Uphamol Menstrual Tablet 500/25mg should not be taken if you are allergic to paracetamol and/or pambrom.

This medication is contraindicated in hepatic failure patients.

Do not use this medication if you are unable to urinate.

WARNING/PRECAUTION

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time.

Do not use this adult product in children under 12 years of age; this will provide more than the recommended dose and may cause liver damage.

For patients with underlying liver disease and renal impairment, please seek medical advice before taking this medication.

It should be given with care to patients with alcohol dependence.

Before using Pambrom, tell your doctor if you take any other diuretic (water pill), or if you have high blood pressure, heart disease or kidney disease. If you have any of these conditions, you may need a dose adjustment or special tests to safely take pambrom.

Please consults your doctor or pharmacists if symptoms persists.

Allergy alert: Paracetamol may cause severe skin reactions.

Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition.

If these reactions occur, stop use and seek medical assistance right away.

Date of Revision: Feb 2026

1500010598 N.2

Pantone Black C

Product's Leaflet

Size: 196mm (W) x 107mm (H)