



RHONAMOX[®] 50 WSP

Powder for Oral Solution
50%w/w

Composition: Amoxicillin 50% w/w (as amoxicillin potency) (equivalent to 57.4% w/w of amoxicillin trihydrate)

Product Description: White to off-white powder

Pharmacodynamics: Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against Gram positive and Gram negative bacteria. It owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

Pharmacokinetics: Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

Indication:
Chickens: For the treatment of pasteurellosis and colibacillosis.
Turkeys: For the treatment of pasteurellosis.
Ducks: For the treatment of infections caused by *Streptococcus bovis*, *Pasteurella anatipestifer* and *Escherichia coli*.
Pigs: For the treatment of salmonellosis and pasteurellosis.

Routes of administration: Oral, in drinking water or liquid feed

Dosage:
Chickens
Dissolve 150g of the product in 450 litres of water immediately before use. This will provide medication for 5000 kg bodyweight of birds for 1 day (15 mg amoxicillin per kilogram bodyweight). The medicated drinking water should be provided on alternate days. The total period of treatment should be for 3 days (2 days of medication) or in severe cases for 5 days (3 days of medication).

Ducks
Administer in the drinking water to give 20 mg amoxicillin/kg bodyweight. Medication should be provided on alternate days for 3 days, i.e. 2 days of medication. Because of the variable water intake of ducks, depending on temperature, light and feeding regime, exact recommendation of dilution of the product cannot be given. In each case the quantities of the product to be used must be calculated taking into account the water intake and total bodyweight of the birds involved.

Turkeys
Administer in the drinking water to give 15 - 20 mg amoxicillin/kg bodyweight. Medication to be provided on alternate days for 5 days (medication on days 1, 3 and 5). Dosage should be calculated taking into account normal water intake and total bodyweight of birds.

W10cm H14cm

Pigs
Administer in the drinking water to give 20 mg amoxicillin/kg bodyweight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days.

Contraindications: The product should not be administered to rabbits, hamsters, gerbils and guinea pigs.
Do not use in animals with known hypersensitivity to the active substance.

Warnings and Precautions:
(i) Special precautions for use in animals
Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.
(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals
Avoid inhalation of dust.
Wash hands after use.
Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1) Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.
2) Handle this product with great care to avoid exposure, taking all recommended precautions.
3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Special precautions for disposal of unused products
Drug containers and any residual contents should be disposed of safely, according to the regulation in force.

Interactions with other medicaments: None known

Statement on usage during pregnancy and lactation: Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin. Use only according to the benefit/risk assessment of the responsible veterinarian.

Adverse Effects: Occasional allergies to the penicillins have been observed but these are rare.

Overdose and Treatment: No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

Storage Conditions: Store in dry place below 30°C.

Withdrawal Period:
Chickens: 1 day
Ducks: 9 days
Turkeys: 5 days
Pigs: 2 days
Not authorised for use in laying birds producing eggs for human consumption.

Packing: 100 g, 500 g and 1000g in aluminium sachets
FOR VETERINARY USE ONLY

Registration holder and manufactured by:



RHONE MA MALAYSIA SDN BHD (200001023279)

Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor, Malaysia.
Tel: 03-7873 7355 Fax: 03-2770 0119
Email: customercareline@rhonema.com

Date of revision: February 2020
5LRHONW3-500 04/FEB20

LABEL CODE:	VERSION	COLORS
5LRHONW3-500	Version: 4 MMM/YY: FEB 20	Black
PRODUCT: RHONAMOX 50 WSP POWDER FOR ORAL SOLUTION 50% w/w PACK SIZE: 100 G, 500 G and 1KG COUNTRY: MY DIMENSIONS: cm = 10 x 14		⚠ Please use the official PANTONE® (P) matching system for an accurate color representation.
Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME:	COMPLETE NAME:	COMPLETE NAME:
DATE:	DATE:	DATE:
SIGNATURE:	SIGNATURE:	SIGNATURE: