

FLUCOR DAY PE SOFT GELATIN CAPSULE

WIFLU06-1 (MY)

DESCRIPTION

Oblong shaped, pink transparent soft gelatin capsule printed with 'Flucor PE', filled with transparent pink liquid.

COMPOSITION

Each soft gelatin capsule contains:

Paracetamol	250.00 mg/cap
Dextromethorphan HBr	10.00 mg/cap
Phenylephrine HCl	5.00 mg/cap

ACTIONS AND PHARMACOLOGY

Paracetamol, as an analgesic, may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (CNS) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation. The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation. Paracetamol probably produces antipyresis by acting centrally on the hypothalamic heat-regulating center to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating, and heat loss. The central action probably involves inhibition of prostaglandin synthesis in the hypothalamus.

Dextromethorphan suppresses the cough reflex by a direct action on the cough center in the medulla of the brain.

Phenylephrine hydrochloride is a sympathomimetic with mainly direct effects on adrenergic receptors. It has predominantly alpha-adrenergic activity and is without significant stimulating effects on the CNS at usual dose. Following oral administration or topical application of phenylephrine to the mucosa, constriction of blood vessels in the nasal mucosa relieves nasal congestion associated with allergy or head colds.

PHARMACOKINETICS

Paracetamol is readily absorbed from the gastrointestinal tract with peak plasma concentrations occurring about 10 to 60 minutes after oral administration and is distributed into most body tissues. The elimination half-life of paracetamol varies from about 1 to 3 hours. It is metabolized predominantly in the liver and excreted in the urine mainly as the glucuronide and sulphate conjugates, less than 5% is excreted as unchanged paracetamol.

Dextromethorphan is rapidly absorbed from the gastrointestinal tract. It is metabolized in the liver and excreted in the urine as unchanged dextromethorphan and demethylated metabolites including dextrophan, which has some cough suppressant activity.

Phenylephrine has a low bioavailability owing to irregular absorption and first-pass metabolism by monoamine oxidase in the gut and liver.

INDICATIONS

For the temporary relief of nasal congestion, minor aches and pains, headache, fever and cough associated with the common cold.

CONTRAINDICATIONS

Risk-benefit should be considered when the following medical problems exist:

- Active alcoholism, or hepatic disease, or viral hepatitis
- Phenylketonuria
- Renal function impairment
- Sensitivity to acetaminophen or aspirin, or dextromethorphan or phenylephrine or other sympathomimetics.
- Asthma, chronic bronchitis, emphysema or productive cough
- Diabetes
- Cardiovascular diseases or hypertension

WARNING

When used for treatment of Cough and Cold:

- a) Not to be used in children less than 2 years of age.
- b) To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time.

Do not take Flucor Day PE if you have taken a monoamine oxidase inhibitor (MAOI) such as Isocarboxazid, Phenelzine, Tranylcypromine or Selegiline in the last 14 days.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

PRECAUTIONS

- Paracetamol should be given with care to patients with impaired kidney or liver function and to patients with alcohol dependence.
- Dextromethorphan should not be given to patient at risk of developing respiratory failure. Caution is needed in patients with a history of asthma and it should not be given during an acute attack.
- Phenylephrine should be avoided in severe hyperthyroidism and used with caution in severe ischaemic heart disease.
- Do not exceed recommended dosage. Consult general practitioner should symptoms persist.
- Patients with hearts disease, thyroid disease, high blood pressure, breathing problems, persistent or chronic cough, asthma, diabetes, chronic bronchitis, glaucoma or difficulty in urination due to enlargement of the prostate gland should consult a doctor before using this product.

PREGNANCY AND LACTATION

Safety and efficacy for use during pregnancy and lactation has not been established.

MAIN SIDE/ADVERSE EFFECTS

- Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.
- Haematological reactions including thrombocytopenia, leukopenia, pancytopenia, neutropenia and agranulocytosis have been reported as side effects of paracetamol with skin rashes and other hypersensitivity reactions occur occasionally.

- Adverse effects with dextromethorphan appear to be rare and may include dizziness and gastrointestinal disturbances.
- The commonest adverse effects of phenylephrine include anxiety, palpitations, arrhythmias, headache, cold extremities / gangrene, hypertension, nausea, vomiting, sweating, weakness, fear, restlessness, insomnia, confusion, irritability, psychotic states, dyspnoea, anorexia, extravasation causing tissue necrosis and sloughing, mydriasis, difficulty in micturition and urinary retention, piloerection, increased salivation, hyperglycaemia and lactic acidosis.

DRUG INTERACTIONS

Concurrent use of Flucor Day PE and the following drugs must be avoided or if necessary, use with care:

- Alcohol
- Hepatic enzyme inducers
- Hepatotoxic medications
- Anticoagulants, coumarin- or indandione-derivative
- Antihypertensive drugs
- Antidepressant drugs
- Monoamine oxidase inhibitor (MAOI)

OVERDOSE

Symptoms of overdosage include gastrointestinal upset (diarrhoea, loss of appetite, nausea and vomiting, stomach cramps or pains), increased sweating, hepatotoxicity (pain, tenderness, and/or swelling in upper abdominal area), ataxia, blurred vision, coma; confusion, drowsiness or dizziness, respiratory depression, severe unusual excitement, nervousness, restlessness or irritability, urinary retention, convulsions, fast breathing, hallucinations, paresthesia, hypertension and palpitations.

Treatment for overdose

- Emptying of stomach via induction of emesis or gastric lavage
- Assisted respiration
- Vital signs monitoring
- For convulsions, i.v. diazepam may be administered.

DOSAGE AND ADMINISTRATION

Adults and children 12 years of age and above: 2 softgels every 4 hours; not exceeding 8 softgels in 24 hours.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C. Protect from moisture.

Presentation/Packing:

Blister pack of 1 x 8's

Product Registration Holder /
Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman,
30010 Ipoh, Malaysia.

Revision date: January 2017