

# OXY 200g/kg Premix

## FOR ANIMAL USE ONLY

### Brand or Product Name :

OXY 200g/kg Premix

### Name and Strength of Active Substance(s)

Each kg contains 215.6g Oxytetracycline dehydrate, which is equivalent to 200g Oxytetracycline base..

### Product Description

OXY 200g/kg Premix is a yellowish to light brownish powder.

### Pharmacodynamics Properties

Oxytetracycline is a tetracycline derivative obtained from *Streptomyces rimosus*. Oxytetracycline generally acts as bacteriostatic antibiotic and inhibit protein synthesis by reversibly binding to 30S ribosomal subunits of susceptible organisms. This binding effectively prevents the addition of amino acids to the elongating peptide chain, inhibiting bacterial protein synthesis. Oxytetracycline also is believed to reversibly bind to 50S ribosomes and additionally alter cytoplasmic membrane permeability in susceptible organisms. Oxytetracycline is a broad-spectrum antibiotics. It has activity against Gram-positive and gram-negative aerobic and anaerobic bacteria, including *Brucella*, *Coxiella*, *Ehrlichia*, *Mycoplasmas*, *Chlamydiae*, *Rickettsias*, and protozoa.

### Pharmacokinetics Properties

Oxytetracycline is readily absorbed after oral administration to fasting animals. Bioavailability are approximately 60 – 80 %. The presence of food or dairy products can reduce the amount of oxytetracycline absorbed. Oxytetracycline is widely distributed in the body, including to the heart, kidney, lungs, muscle, pleural fluid, bronchial secretions, sputum, bile, saliva, urine, synovial fluid, ascetic fluid, and aqueous and vitreous humor. Only small quantities of oxytetracycline is distributed to the CSF. Oxytetracycline crosses the placenta, enter fetal circulation and are distributed into milk. The volume of distribution of oxytetracycline is approximately 2.1 L/kg in small animals. The amount of plasma protein binding of oxytetracycline is about 10 – 40%. Oxytetracycline is eliminated unchanged primarily via glomerular filtration. The drug is not metabolized, but is excreted into GI tract via biliary and nonbiliary routes and may become inactive after chelation with fecal materials. The elimination half-life of oxytetracycline is approximately 4 – 10 hours in dogs and cats, 4.3 – 9.7 hours in cattle, 10.5 hours in horses, 6.7 hours in swine, and 3.6 hours in sheep.

### Indication

For the control and treatment of ailments caused by microorganisms sensitive to oxytetracycline.

Chickens : For control of infectious synovitis caused by *Mycoplasma synoviae* ; control of fowl cholera caused by *Pasteurella multocida* susceptible to oxytetracycline ; control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to oxytetracycline ; reduction of mortality due to air sacculitis (air sac infection) caused by *Escherichia coli* susceptible to oxytetracycline.

Swine : For treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella chloreraesuis* susceptible to oxytetracycline and treatment of bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline ; for control and treatment of *Leptospirosis* caused by *Leptospira pomona* susceptible to oxytetracycline.

### Recommended Dosage

To be mixed thoroughly with animal feed.

| Indications for Use In Chickens                                                                                                                                         | Amount of Oxytetracycline base                        | Amount of Oxy 200g/kg Powder           |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|----------------------------------------|
| Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline    | 100 – 200g/tonne<br>Feed continuously for 7 – 14 days | 0.5 – 1.0 kg of Oxy 200g/kg Powder/ton |
| Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline | 400g/tonne<br>Feed continuously for 7 – 14 days.      | 2.0 kg of Oxy 200g/kg Powder/ton       |
| Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to oxytetracycline.                                      | 500g/tonne<br>Feed continuously for 5 days.           | 2.5 kg of Oxy 200g/kg Powder/ton       |

  

| Indications for Use in Swine                                                                                                                                                                                                           | Amount of Oxytetracycline                                                       | Amount of Oxy 200g/kg Powder                            |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|---------------------------------------------------------|
| Treatment of bacterial enteritis caused by <i>E.coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline. | 22.05mg/kg of body weight daily.<br>Feed continuously for 7 – 14 days           | 110.25 mg of Oxy 200g/kg Powder/kg of body weight daily |
| For control and treatment of <i>Leptospirosis</i> (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira Pomona</i> susceptible to oxytetracycline.                                                  | 22.05mg/kg of body weight daily.<br>Feed continuously for not more than 14 days | 110.25 mg of Oxy 200g/kg Powder/kg of body weight daily |

### Contraindications

Hypersensitivity to any tetracycline drug.

### Warnings and Precautions

Use with caution in patients with renal insufficiency and hepatic impairment. Lower than normal dosage is recommended with enhanced monitoring of renal and hepatic function. Chicken and animal intended for human consumption must not be slaughtered before the specified withdrawal periods.

### Interactions with Other Medicaments

When orally administered, tetracyclines can chelate divalent or trivalent cations which can decrease the absorption of the tetracyclines or the other drug if it contains these cations. Oral antacids, saline cathartics or other GI products containing aluminum, calcium, magnesium, zinc or bismuth cations are most commonly associated with this interaction. It is recommended that all oral tetracyclines be given at least 1-2 hours before or after the cation-containing product.

### Statement on usage during pregnancy and lactation

Because tetracyclines can retard fetal skeletal development and discolor deciduous teeth, they should only be used in the last half of pregnancy when the benefits outweigh the fetal risks. Do not administer to chickens producing eggs for human consumption

### Adverse Effects / Undesirable Effects

The most common adverse reactions include gastrointestinal upset : nausea, vomiting, diarrhea and anorexia.

### Overdose and Treatment

Stop medication and to seek for veterinarian advice if there is sign of overdose. Gastric lavage is beneficial for first hours after ingestion and milk slows down absorption. Tetracyclines are generally well tolerated after acute overdoses. Manifestations of an acute overdose would most likely be associated with GI distress (nausea, anorexia, vomiting, diarrhea). Should the patient develop severe emesis or diarrhea, fluid and electrolytes should be monitored and replaced if necessary.

### Withdrawal Period(s)

Swine and Chickens : 7 days.

### Shelf-life

Original packing expired 3 years from date of manufacture. Opened bag to be discarded after 24 hours. Unused medicated feed to be discarded after 24 hours.

### Allowable Maximum Residual Limit (MRL)

Fat (pig, chicken and turkey ) = 10µg/kg, egg (chicken) = 200µg/kg, muscle (pigs, chicken and turkey ) = 100µg/kg, liver (pigs, chicken and turkey ) = 300µg/kg, kidney (pig, chicken and turkey ) = 600µg/kg [Ref. REGOV, Version 2, dated Dec 2009]

### Storage Conditions

Store airtight and below 25°C. Keep out of reach of children. Jauhi ubat dari kanak-kanak.

### Disposal of Containers

Traces of medicated powder left in an opened bag can be diluted by pouring small amount of animal feed into the bag, mix well and use in animal treatment. Emptied bag can be safely disposed in trash according to local government regulations.

### Dosage Forms and packaging available

Available in 500 g and 1 kg aluminum sachet and 25 kg paper bag.

### Name and Address of manufacturer

Life Biopharma Sdn. Bhd. No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan Darul Khusus, Malaysia.

### Date of Revision of Package Insert

2012-06-08