

Pengesic SR Tablet 100 mg

Description

10 mm. White, round, transparent film-coated tablet, shallow convex faces and 'hovid' embossed on one face.

Composition

Each tablet contains:
Tramadol Hydrochloride 100 mg

Actions and Pharmacology

Tramadol is a centrally-acting analgesic that is not chemically related to opiates. The mechanism of action of tramadol has not been fully determined, but it may bind to mu-opioid receptors and inhibit the reuptake of norepinephrine (NE) and serotonin. The ability of tramadol to inhibit the neuronal uptake of monoamines in the same concentration range at which it binds to mu-opioid receptors differentiates it from typical opioids.

Pharmacokinetics

Absorption

Tramadol is readily absorbed after oral doses but is subject to first-pass metabolism. Mean absolute bioavailability of 100-mg dose is approximately 75%. The rate or extent of absorption is not significantly affected by administration with food.

Distribution

Tramadol is widely distributed, crosses the placenta, and appears in small amounts in breast milk.

Protein Binding

Low (20%)

Biotransformation

Tramadol is metabolized by N- and O-demethylation via the cytochrome P450 isoenzymes CYP3A4 and CYP2D6 and glucuronidation or sulfation in the liver. The metabolite O-des-methyltramadol is pharmacologically active. The inactive metabolites are formed by N-demethylation.

Elimination

Tramadol is excreted mainly in the urine, predominantly as metabolites. The elimination half life after oral doses is about 6 hours.

Indications

Pengesic SR tablet 100 mg is indicated for the management of moderate to severe pain.

Contraindications

It is generally contraindicated in the following conditions:

- Respiratory depression, especially in the presence of cyanosis and excessive bronchial secretion.
- In the presence of acute alcoholism, head injuries and conditions in which intracranial pressure is raised.
- It should not be used in patients receiving MAO inhibitors.
- Previous hypersensitivity to Tramadol.
- It should not be used during pregnancy.
- It should not be administered during lactation as tramadol and its metabolites have been detected in breast milk.

Precautions

Tramadol may only be used after careful consideration of the benefit/risk ratio and relevant precautions in:-

- dependence on opioids
- consciousness disorders of uncertain origin, shock
- disorders of the respiratory centre or function
- increased intracranial pressure due to head injuries or brain diseases

In patients sensitive to opiates the medicinal product should only be used with caution.

Convulsion have been reported in patients receiving tramadol at the recommended dose levels. The risk may be increased when doses of tramadol exceed the recommended upper daily dose limit (400mg). In addition, tramadol may increase the seizure risk in patients taking other medication that lowers the seizure threshold. Patients with epilepsy or those susceptible to seizures should be only treated with tramadol if there are compelling circumstances.

Tramadol has a low dependence potential. On long-term use tolerance, psychic and physical dependence, treatment with Tramadol should only be carried out for short periods under strict medical supervision.

Tramadol is not suitable as a substitute in opioid-dependent patients. Although tramadol is an opiate agonist, it cannot suppress morphine withdrawal symptoms.

Tramadol is not suitable for children below the age of 12 year.

During treatment with Tramadol, alcohol should be avoided as it may have an unfavourable influence on the effect of the drug.

Pregnancy and Lactation

- It should not be used during pregnancy.
- It should not be administered during lactation as tramadol and its metabolites have been detected in breast milk.

Interaction with Other Medicaments

- It should be given with extreme caution in patients taking monoamine oxidase inhibitors or within 14 days of stopping such treatment.
- The depressant effects of Tramadol are enhanced by depressants of the central nervous system such as alcohol, anaesthetics, hypnotics and sedatives, tricyclic anti depressants and phenothiazines.
- Opioid analgesics with some antagonist activity such as buprenorphine, butorphanol, nalbuphene or pentazocine may precipitate withdrawal symptoms in patients who have recently used pure agonists such as Tramadol.
- On the concomitant administration of Pengesic SR tablet 100 mg with substances which also act on the central nervous system (e.g. tranquillizers, hypnotics) the sedative effects (fatigue) may be intensified. At the same time, however, combining Pengesic SR tablet 100 mg with a tranquillizer, for example, will probably have a favourable effect on the pain sensation. Pengesic SR tablet 100 mg should not be used in patients receiving MAO inhibitors.

Main Side/Adverse Effects

- Tramadol can produce drug dependence of the mu-opioid type and may potentially be abused. Tolerance development, drug-seeking behavior and craving have been associated with the use of tramadol.
- The commonest side effects include abdominal or stomach pain, anorexia, asthenia, central nervous system (CNS) stimulation, constipation, diarrhea, dizziness or vertigo, drowsiness, dry mouth, dyspepsia, headache, nausea, pruritus, skin rash, sweating and vomiting.

Overdose

Clinical features:

Cold, clammy skin; confusion; convulsions; severe dizziness; severe drowsiness; pinpoint pupils of eyes; slow heartbeat; slow or troubled breathing; unconsciousness; severe weakness.

Treatment:

Intensive supportive therapy may be required to correct failure and shock. In addition, the specific antagonist naloxone hydrochloride is used to counteract very rapidly the severe respiratory depression and coma produced by excessive doses of opioid analgesics. A dose of 0.4 to 2 mg is given by intravenous injection, repeated at intervals of 2 to 3 minutes if necessary, up to 10 mg. Naloxone may also be given by subcutaneous or intramuscular injection or by intravenous infusion.

Dosage and Administration

The dosage should be adjusted to the intensity of the pain and the sensitivity of the individual patient. Unless otherwise prescribed, Pengesic SR tablet 100 mg should be taken as follows - with or without meals:

Single dose for adults and adolescents over 12 years of age:

1-2 tablets (100-200mg tramadol hydrochloride) should be swallowed whole (not divided or chewed) with sufficient liquid, preferably mornings and evenings. If in severe pain the analgesic demand is likely to be higher, 2 tablets (200mg tramadol hydrochloride) can be taken as initial dose.

The lowest analgesically effective dose should generally be selected. Daily doses of 400mg tramadol hydrochloride should not be exceeded, except in special clinical circumstances. The dosage interval should not be less than 8 hours.

Children:

On account of the dosage strength, Pengesic SR tablet 100 mg is not recommended for children below the age of 12 years.

Geriatric patients:

A dose adjustment is usually not necessary in elderly patients (up to 75 years) with no clinically manifest hepatic or renal insufficiency. In old patients (above the age of 75 years) elimination may be prolonged. Therefore, if necessary the dosage intervals may have to be extended according to the patient's requirements.

Hepatic and renal insufficiency/dialysis:

In acute pain a dosage adjustment is not necessary as, Pengesic SR tablet 100 mg is given only once or a few times. In patients with severe renal and/or hepatic insufficiency, Pengesic SR tablet 100 mg should not be administered. In less severe cases prolongation of the dosage interval should be considered.

Duration of Treatment:

Pengesic SR tablet 100 mg must not be given for longer than therapeutically absolutely necessary. If long-term pain treatment is necessary, checks should be carried out at regular and brief intervals (if necessary with breaks in treatment) as to whether and in what doses further treatment with Pengesic SR tablet 100 mg is necessary.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage	:	Store below 30°C. Protect from light and moisture.
Presentation/Packing	:	Blister pack of 1x10's., 3x10's.
Marketing Authorization holder	:	Hovid Bhd, 121 Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.
Manufactured by	:	Hovid Bhd., Lot 56442, 7½ Miles, Jalan Ipoh / Chemor, 31200 Chemor, Malaysia.
Date of Information	:	September 2012