

Product Name:

AMPIVET 50% Water Soluble Powder

Composition:

Each gram contains:

Ampicillin 500mg

Product Description:

White powder

Pharmacodynamics:

Ampicillin (a beta-lactam antibiotic) prevents formation of the bacterial cell wall by interfering with the final stage of peptidoglycan synthesis. It inhibits the activity of transpeptidase enzymes which catalyse cross-linkage of glycopeptide polymer units that form the cell wall.

Pharmacokinetics:

Ampicillin anhydrous and trihydrate are relatively stable in the presence of gastric acid. After oral administration, ampicillin is about 30-55% absorbed in humans (empty stomach) and monogastric animals. Food will decrease the rate and extent of oral absorption. When administered parenterally (IM, SC) the trihydrate salt will achieve serum levels of approximately 1/2 those of a comparable dose of the sodium salt. The trihydrate parenteral dosage form should not be used where higher MIC's are required for treating systemic infection. After absorption, the volume of distribution for ampicillin is approximately 0.3L/kg in humans and dogs, 0.167L/kg in cats, and 0.16-0.5 L/kg in cattle. The drug is widely distributed to many tissues, including liver, lungs, muscle, bile, and ascitic, pleural and synovial fluids. Ampicillin is approximately 20% bound to plasma protein. milk levels of ampicillin are considered low. In lactating dairy cattle, the milk to plasma ratio is about 0.3. Ampicillin is eliminated primarily through renal mechanism, principally by tubular secretion, but some of the drug is metabolized by hydrolysis to penicilloic acids (inactive) and then excreted in the urine. elimination half-lives of ampicillin have been reported as 45-80 minutes in dogs and cats, and 60 minutes in swine.

Indications:

Oral treatment of porcine colibacillosis (*E. coli*) and salmonellosis (*Salmonella* species) infections in swine, and bacterial pneumonia caused by *Pasteurella multocida*, *Staphylococcus* species *Streptococcus* species and *Salmonella* species.

Dosage and Administration:

Swine up to 75 pounds of body weight: 5 milligrams of ampicillin per pound of body weight twice daily for up to 5 days

Contraindications:

Do not administer to penicillin sensitive animals.

Do not administer to small herbivores such as gerbils, guinea pigs, hamsters and rabbits.

Interaction with Other Medicaments:

It is not recommended that bacteriostatic antimicrobials be administered concomitantly.

Pregnancy and Lactation:

Can be safely administered to pregnant or lactating animals.

Side effects:

None known

Symptoms and Treatment of Overdose:

No known symptoms of overdose.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1. Do not handle if you know you are sensitised, or if you have been advised not to work with such preparations.
2. Handle this product with great care to avoid exposure, taking all recommended precautions.
3. If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes, or difficulty with breathing are more serious symptoms and require urgent medical attention.

Wash hands after use.

Withdrawal Periods:

Swine: 24 hours before slaughter.

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,
Prai Industrial Estate MK1,
13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

DEXCHEM INDUSTRIES SDN BHD

No. 6 & 8, Jalan Industri Cherok To'Kun 2,
Taman Industri Cherok To'Kun,
14000 Bukit Mertajam, Pulau Pinang.

DATE OF REVISION: APRIL 2019