

Product Name:

AMOVET 50% Water Soluble powder

Composition:

Each gram contains:

Amoxicillin 500mg

Product Description:

White powder

Pharmacodynamics:

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against gram positive and gram negative bacteria. Its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall

Pharmacokinetics:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Indications:

Chickens: For the treatment of pasteurellosis and colibacillosis. Pigs: For the treatment of salmonellosis and pasteurellosis.

Dosage and Administration:

For administration via the drinking water, the solution should be prepared with fresh potable water.

In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst on treatment.

Chickens:

Dissolve 150g of the product in 450 litres (100 gallons) of water immediately before use.

This will provide medication for 5000kg bodyweight of birds for 1 day (15mg amoxicillin trihydrate per kilogram bodyweight).

The medicated drinking water should be provided on alternate days. The total period of treatment should be for 3 days (2 days of medication) or in severe cases for 5 days (3 days of medication).

Pigs:

Administer in the drinking water to give 20mg amoxicillin trihydrate/kg bodyweight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days. Dissolve 150g of the product in the requisite quantity of water immediately before use.

As a guide, 150g of the product will treat the following number of birds:

Broilers	
Age in weeks	No. of birds
1	30000
2	13500
3	7500
4	5000
6	3000
8	2000

Contraindications:

Should not be administered to rabbits, hamster, gerbils and guinea pigs. Do not use in animals with known hypersensitivity to the active substance

Interaction with Other Medicaments:

None known

Pregnancy and Lactation:

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin. Use only according to the benefit/risk assessment of the responsible veterinarian.

Side effects:

None known

Symptoms and Treatment of Overdose:

Treatment should be symptomatic and no specific antidote is available.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may cause cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

People with known hypersensitivity to penicillin or cephalosporin should avoid contact with the product.

Handle this product with great care to avoid exposure, taking all recommended precautions. Wear protective clothing, impervious gloves and either a disposable half mask respirator or a non-disposable respirator when mixing and handling the product. Wash hands immediately after handling the product.

If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Withdrawal Periods:

Chicken	: 1 day
Swine	: 2 days
Ducks	: 9 days
Calves	: 9 days
Turkeys	: 5 days

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.
2692, Tingkat Perusahaan Enam,
Prai Industrial Estate MK1,
13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

DEXCHEM INDUSTRIES SDN BHD
No. 6 & 8, Jalan Industri Cherok To’Kun 2,
Taman Industri Cherok To’Kun,
14000 Bukit Mertajam, Pulau Pinang.

DATE OF REVISION: APRIL 2019